



Full wwPDB X-ray Structure Validation Report ⓘ

May 14, 2020 – 07:00 am BST

PDB ID : 1SAR
Title : DETERMINATION AND RESTRAINED LEAST-SQUARES REFINEMENT OF THE CRYSTAL STRUCTURES OF RIBONUCLEASE SA AND ITS COMPLEX WITH 3'-GUANYLIC ACID AT 1.8 ANGSTROMS RESOLUTION
Authors : Sevcik, J.; Dodson, E.J.; Dodson, G.G.
Deposited on : 1990-12-13
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	96	
1	B	96	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 1735 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RIBONUCLEASE SA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	96	Total 745	C 465	N 122	O 155	S 3	0	0	0
1	B	96	Total 745	C 465	N 122	O 155	S 3	0	0	0

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	A	1	Total 5	O 4	S 1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	126	Total 127	O 127	0	1

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	113	Total 113	O 113	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: RIBONUCLEASE SA



- Molecule 1: RIBONUCLEASE SA



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	64.90Å 78.32Å 38.79Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	(Not available) – 1.80 27.56 – 1.83	Depositor EDS
% Data completeness (in resolution range)	(Not available) ((Not available)-1.80) 95.9 (27.56-1.83)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.00 (at 1.83Å)	Xtrriage
Refinement program	PROLSQ	Depositor
R, R_{free}	0.172 , (Not available) 0.158 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	18.3	Xtrriage
Anisotropy	0.359	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 57.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	1735	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.53% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.30	3/763 (0.4%)	2.56	49/1043 (4.7%)
1	B	1.28	2/763 (0.3%)	2.69	49/1043 (4.7%)
All	All	1.29	5/1526 (0.3%)	2.63	98/2086 (4.7%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3
1	B	0	4
All	All	0	7

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	35	VAL	C-N	8.75	1.54	1.34
1	B	74	GLU	CD-OE2	5.74	1.31	1.25
1	A	80	TYR	CE2-CZ	5.34	1.45	1.38
1	B	31	SER	CB-OG	-5.21	1.35	1.42
1	A	74	GLU	CD-OE1	5.20	1.31	1.25

All (98) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	63	ARG	NE-CZ-NH1	25.95	133.28	120.30
1	B	63	ARG	NE-CZ-NH2	-19.16	110.72	120.30
1	B	40	ARG	NE-CZ-NH1	16.57	128.59	120.30
1	A	65	ARG	NE-CZ-NH1	-16.18	112.21	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	65	ARG	NE-CZ-NH2	15.53	128.06	120.30
1	B	68	ARG	NE-CZ-NH2	-12.44	114.08	120.30
1	A	74	GLU	CB-CG-CD	11.91	146.35	114.20
1	A	30	TYR	CB-CG-CD1	-11.47	114.12	121.00
1	A	78	GLU	OE1-CD-OE2	11.31	136.88	123.30
1	A	30	TYR	CB-CG-CD2	11.30	127.78	121.00
1	A	1	ASP	CB-CG-OD1	11.09	128.28	118.30
1	A	49	TYR	CB-CG-CD2	-10.64	114.61	121.00
1	A	40	ARG	CD-NE-CZ	10.62	138.48	123.60
1	B	86	TYR	CB-CG-CD1	10.52	127.31	121.00
1	A	69	ARG	NE-CZ-NH2	10.39	125.49	120.30
1	B	72	CYS	CA-CB-SG	-10.02	95.97	114.00
1	B	52	TYR	CB-CG-CD1	-9.78	115.13	121.00
1	B	49	TYR	CB-CG-CD1	-9.63	115.22	121.00
1	A	74	GLU	OE1-CD-OE2	-9.58	111.80	123.30
1	A	63	ARG	NE-CZ-NH1	9.35	124.97	120.30
1	B	33	ASP	CB-CG-OD1	9.34	126.71	118.30
1	A	3	SER	O-C-N	-9.03	107.84	123.20
1	A	72	CYS	CA-CB-SG	-8.83	98.11	114.00
1	A	33	ASP	CB-CG-OD1	8.75	126.17	118.30
1	B	30	TYR	CB-CG-CD2	8.75	126.25	121.00
1	B	5	THR	O-C-N	8.71	136.63	122.70
1	B	40	ARG	CD-NE-CZ	8.58	135.62	123.60
1	B	51	TYR	CB-CG-CD2	-8.55	115.87	121.00
1	B	14	GLU	CG-CD-OE1	8.47	135.25	118.30
1	B	40	ARG	NE-CZ-NH2	-8.26	116.17	120.30
1	A	80	TYR	CB-CG-CD2	-7.98	116.21	121.00
1	B	30	TYR	CB-CG-CD1	-7.74	116.36	121.00
1	B	51	TYR	CB-CG-CD1	7.72	125.63	121.00
1	B	17	ASP	CB-CG-OD1	7.67	125.21	118.30
1	B	3	SER	N-CA-CB	7.65	121.97	110.50
1	A	49	TYR	CG-CD2-CE2	-7.62	115.21	121.30
1	B	74	GLU	CB-CG-CD	7.56	134.61	114.20
1	A	69	ARG	O-C-N	7.53	134.74	122.70
1	A	20	ASN	O-C-N	7.35	134.47	122.70
1	A	3	SER	CA-C-N	7.34	130.88	116.20
1	B	54	GLU	CA-CB-CG	-7.15	97.67	113.40
1	A	54	GLU	OE1-CD-OE2	7.14	131.86	123.30
1	A	28	PHE	CB-CG-CD1	-7.06	115.86	120.80
1	B	81	TYR	CB-CG-CD2	6.99	125.19	121.00
1	B	89	PHE	CG-CD1-CE1	-6.84	113.27	120.80
1	A	68	ARG	NE-CZ-NH2	-6.83	116.89	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	68	ARG	NE-CZ-NH1	6.80	123.70	120.30
1	A	49	TYR	CD1-CG-CD2	6.75	125.33	117.90
1	A	77	GLN	CB-CA-C	-6.56	97.27	110.40
1	B	28	PHE	CB-CG-CD2	-6.55	116.21	120.80
1	A	40	ARG	CA-CB-CG	6.35	127.36	113.40
1	A	84	ASP	CB-CG-OD1	6.28	123.95	118.30
1	B	63	ARG	CD-NE-CZ	6.20	132.28	123.60
1	A	71	ILE	CB-CG1-CD1	6.19	131.22	113.90
1	A	74	GLU	CG-CD-OE1	6.12	130.55	118.30
1	B	38	GLN	CA-CB-CG	-6.08	100.03	113.40
1	A	82	THR	O-C-N	6.06	133.50	123.20
1	A	89	PHE	O-C-N	6.01	132.32	122.70
1	B	89	PHE	CZ-CE2-CD2	-5.99	112.92	120.10
1	A	1	ASP	CB-CG-OD2	-5.98	112.92	118.30
1	B	54	GLU	CG-CD-OE2	-5.95	106.41	118.30
1	B	17	ASP	O-C-N	5.93	132.18	122.70
1	B	1	ASP	CB-CG-OD2	-5.92	112.97	118.30
1	A	73	GLY	N-CA-C	-5.83	98.53	113.10
1	B	74	GLU	CA-C-N	5.82	130.00	117.20
1	B	70	ILE	CA-CB-CG1	-5.80	99.98	111.00
1	B	3	SER	CB-CA-C	-5.77	99.13	110.10
1	A	25	ASP	CB-CG-OD2	-5.76	113.11	118.30
1	A	13	PRO	O-C-N	5.73	131.87	122.70
1	B	84	ASP	CB-CG-OD1	5.73	123.45	118.30
1	B	25	ASP	CA-CB-CG	-5.71	100.83	113.40
1	A	6	VAL	CA-CB-CG1	5.70	119.45	110.90
1	B	80	TYR	O-C-N	5.70	131.81	122.70
1	A	41	GLU	CG-CD-OE2	-5.56	107.18	118.30
1	B	30	TYR	O-C-N	5.51	131.52	122.70
1	B	14	GLU	CG-CD-OE2	-5.49	107.32	118.30
1	B	46	THR	CA-CB-OG1	-5.46	97.53	109.00
1	A	65	ARG	O-C-N	-5.45	113.94	123.20
1	A	85	HIS	C-N-CA	5.43	135.27	121.70
1	B	55	TYR	CG-CD1-CE1	-5.42	116.96	121.30
1	B	23	ALA	CB-CA-C	5.41	118.21	110.10
1	B	80	TYR	CG-CD1-CE1	-5.39	116.99	121.30
1	A	49	TYR	CG-CD1-CE1	-5.38	116.99	121.30
1	A	78	GLU	CG-CD-OE2	-5.37	107.56	118.30
1	A	14	GLU	CG-CD-OE1	5.26	128.82	118.30
1	A	80	TYR	CD1-CG-CD2	5.25	123.68	117.90
1	A	51	TYR	CD1-CE1-CZ	-5.17	115.14	119.80
1	B	53	HIS	C-N-CA	-5.17	108.77	121.70

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	77	GLN	N-CA-CB	5.16	119.89	110.60
1	B	59	THR	CA-CB-CG2	-5.16	105.17	112.40
1	B	2	VAL	O-C-N	5.14	130.93	122.70
1	A	86	TYR	CA-C-O	-5.09	109.40	120.10
1	B	77	GLN	CB-CA-C	-5.09	100.22	110.40
1	A	68	ARG	CD-NE-CZ	-5.08	116.49	123.60
1	B	79	ASP	CB-CG-OD1	5.05	122.84	118.30
1	B	2	VAL	CB-CA-C	5.04	120.97	111.40
1	A	51	TYR	CG-CD2-CE2	-5.04	117.27	121.30
1	A	69	ARG	NH1-CZ-NH2	-5.03	113.86	119.40

There are no chirality outliers.

All (7) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	40	ARG	Sidechain
1	A	65	ARG	Sidechain
1	A	68	ARG	Sidechain
1	B	40	ARG	Sidechain
1	B	63	ARG	Sidechain
1	B	65	ARG	Sidechain
1	B	68	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	745	0	693	11	0
1	B	745	0	693	10	0
2	A	5	0	0	0	0
3	A	127	0	0	6	0
3	B	113	0	0	3	0
All	All	1735	0	1386	21	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:53:HIS:HE1	3:A:121:HOH:O	1.67	0.76
1:B:54:GLU:OE2	3:B:177:HOH:O	2.09	0.71
1:A:68:ARG:NH1	3:A:204:HOH:O	2.16	0.59
1:A:54:GLU:HB3	1:A:71:ILE:HD12	1.87	0.57
1:A:53:HIS:HD2	3:A:189:HOH:O	1.89	0.56
1:A:74:GLU:HG2	3:A:120:HOH:O	2.09	0.52
1:A:36:VAL:HG11	1:A:49:TYR:CE1	2.44	0.52
1:B:75:ALA:HB3	1:B:78:GLU:HB2	1.95	0.49
1:A:74:GLU:CG	3:A:120:HOH:O	2.61	0.48
1:A:41:GLU:HG3	3:A:178:HOH:O	2.13	0.48
1:A:76:THR:HG22	1:A:77:GLN:HG2	1.97	0.47
1:B:40:ARG:HG2	3:B:159:HOH:O	2.16	0.46
1:B:40:ARG:HG3	1:B:41:GLU:OE2	2.16	0.46
1:B:46:THR:O	1:B:47:GLN:HG2	2.16	0.45
1:B:76:THR:O	1:B:77:GLN:HB2	2.16	0.45
1:A:21:LEU:HA	1:A:21:LEU:HD23	1.82	0.44
1:B:53:HIS:HD2	3:B:176:HOH:O	2.00	0.44
1:B:54:GLU:OE1	1:B:86:TYR:OH	2.21	0.43
1:B:46:THR:C	1:B:47:GLN:HG2	2.40	0.43
1:A:6:VAL:HG11	1:A:11:LEU:HD21	2.01	0.42
1:B:76:THR:O	1:B:77:GLN:CB	2.68	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	94/96 (98%)	93 (99%)	1 (1%)	0	100	100
1	B	94/96 (98%)	94 (100%)	0	0	100	100
All	All	188/192 (98%)	187 (100%)	1 (0%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	82/82 (100%)	80 (98%)	2 (2%)	49	36
1	B	82/82 (100%)	80 (98%)	2 (2%)	49	36
All	All	164/164 (100%)	160 (98%)	4 (2%)	49	36

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	13	PRO
1	A	91	LEU
1	B	41	GLU
1	B	47	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	53	HIS
1	B	47	GLN
1	B	53	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	SO4	A	97	-	4,4,4	0.64	0	6,6,6	1.17	1 (16%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	A	97	SO4	O4-S-O3	-2.57	98.08	109.06

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	96/96 (100%)	-0.70	0 100 100	9, 16, 30, 46	0
1	B	96/96 (100%)	-0.53	0 100 100	11, 17, 34, 47	0
All	All	192/192 (100%)	-0.62	0 100 100	9, 17, 32, 47	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	SO4	A	97	5/5	0.99	0.06	21,22,25,26	0

6.5 Other polymers [i](#)

There are no such residues in this entry.