

# wwPDB EM Validation Summary Report (i)

#### Nov 20, 2022 – 06:12 PM EST

PDB ID : 7SAX

EMDB ID : EMD-24958

Title: Structure of GldLM, the proton-powered motor that drives Type IX protein

secretion and gliding motility in Sphingobacterium wenxiniae

Authors: Hennell James, R.; Deme, J.C.; Lea, S.M.

Deposited on : 2021-09-23

Resolution : 3.00 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev43

MolProbity : 4.02b-467

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

MapQ : 1.9.9

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.31.3

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $ELECTRON\ MICROSCOPY$ 

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\#  ext{Entries})$	${ m EM~structures} \ (\#{ m Entries})$
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion <40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain						
1	A	263	70%	11%	19%				
1	В	263	71%	10%	20%				
2	С	212	· 25% · 72%						
2	D	212	22% • 74%						
2	Е	212	18% 5% 76%						
2	F	212	20% • 76%						
2	G	212	22% • 76%						



# 2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 5305 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called GldM.

$\mathbf{Mol}$	Chain	Residues		$\mathbf{At}$	oms			AltConf	Trace	
1	A	213		C		0	D	0	0	
			1627	1015	284	324	4			
1	B	211	Total	$\mathbf{C}$	N	O	S	0	0	
1	Ъ	211	1614	1007	281	322	4	0	U	

There are 78 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	225	GLU	-	expression tag	UNP A0A1I6R6I5
A	226	ASN	-	expression tag	UNP A0A1I6R6I5
A	227	LEU	_	expression tag	UNP A0A1I6R6I5
A	228	TYR	-	expression tag	UNP A0A1I6R6I5
A	229	PHE	-	expression tag	UNP A0A1I6R6I5
A	230	GLN	-	expression tag	UNP A0A1I6R6I5
A	231	GLY	-	expression tag	UNP A0A1I6R6I5
A	232	GLN	-	expression tag	UNP A0A1I6R6I5
A	233	PHE	-	expression tag	UNP A0A1I6R6I5
A	234	GLY	-	expression tag	UNP A0A1I6R6I5
A	235	SER	-	expression tag	UNP A0A1I6R6I5
A	236	TRP	-	expression tag	UNP A0A1I6R6I5
A	237	SER	-	expression tag	UNP A0A1I6R6I5
A	238	HIS	-	expression tag	UNP A0A1I6R6I5
A	239	PRO	-	expression tag	UNP A0A1I6R6I5
A	240	GLN	-	expression tag	UNP A0A1I6R6I5
A	241	PHE	-	expression tag	UNP A0A1I6R6I5
A	242	GLU	-	expression tag	UNP A0A1I6R6I5
A	243	LYS	-	expression tag	UNP A0A1I6R6I5
A	244	GLY	-	expression tag	UNP A0A1I6R6I5
A	245	GLY	-	expression tag	UNP A0A1I6R6I5
A	246	GLY	-	expression tag	UNP A0A1I6R6I5
A	247	SER	-	expression tag	UNP A0A1I6R6I5
A	248	GLY	-	expression tag	UNP A0A1I6R6I5
A	249	GLY	-	expression tag	UNP A0A1I6R6I5
A	250	GLY	-	expression tag	UNP A0A1I6R6I5

Continued on next page...



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	251	SER	-	expression tag	UNP A0A1I6R6I5
A	252	GLY	-	expression tag	UNP A0A1I6R6I5
A	253	GLY	-	expression tag	UNP A0A1I6R6I5
A	254	GLY	-	expression tag	UNP A0A1I6R6I5
A	255	SER	-	expression tag	UNP A0A1I6R6I5
A	256	TRP	-	expression tag	UNP A0A1I6R6I5
A	257	SER	-	expression tag	UNP A0A1I6R6I5
A	258	HIS	-	expression tag	UNP A0A1I6R6I5
A	259	PRO	_	expression tag	UNP A0A1I6R6I5
A	260	GLN	-	expression tag	UNP A0A1I6R6I5
A	261	PHE	-	expression tag	UNP A0A1I6R6I5
A	262	GLU	-	expression tag	UNP A0A1I6R6I5
A	263	LYS	-	expression tag	UNP A0A1I6R6I5
В	225	GLU	-	expression tag	UNP A0A1I6R6I5
В	226	ASN	-	expression tag	UNP A0A1I6R6I5
В	227	LEU	-	expression tag	UNP A0A1I6R6I5
В	228	TYR	-	expression tag	UNP A0A1I6R6I5
В	229	PHE	-	expression tag	UNP A0A1I6R6I5
В	230	GLN	_	expression tag	UNP A0A1I6R6I5
В	231	GLY	-	expression tag	UNP A0A1I6R6I5
В	232	GLN	-	expression tag	UNP A0A1I6R6I5
В	233	PHE	-	expression tag	UNP A0A1I6R6I5
В	234	GLY	-	expression tag	UNP A0A1I6R6I5
В	235	SER	-	expression tag	UNP A0A1I6R6I5
В	236	TRP	-	expression tag	UNP A0A1I6R6I5
В	237	SER	-	expression tag	UNP A0A1I6R6I5
В	238	HIS	-	expression tag	UNP A0A1I6R6I5
В	239	PRO	-	expression tag	UNP A0A1I6R6I5
В	240	GLN	-	expression tag	UNP A0A1I6R6I5
В	241	PHE	-	expression tag	UNP A0A1I6R6I5
В	242	GLU	-	expression tag	UNP A0A1I6R6I5
В	243	LYS	-	expression tag	UNP A0A1I6R6I5
В	244	GLY	-	expression tag	UNP A0A1I6R6I5
В	245	GLY	-	expression tag	UNP A0A1I6R6I5
В	246	GLY	-	expression tag	UNP A0A1I6R6I5
В	247	SER	-	expression tag	UNP A0A1I6R6I5
В	248	GLY	-	expression tag	UNP A0A1I6R6I5
В	249	GLY	-	expression tag	UNP A0A1I6R6I5
В	250	GLY	-	expression tag	UNP A0A1I6R6I5
В	251	SER	-	expression tag	UNP A0A1I6R6I5
В	252	GLY	-	expression tag	UNP A0A1I6R6I5
В	253	GLY	-	expression tag	UNP A0A1I6R6I5

Continued on next page...



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
В	254	GLY	-	expression tag	UNP A0A1I6R6I5
В	255	SER	-	expression tag	UNP A0A1I6R6I5
В	256	TRP	-	expression tag	UNP A0A1I6R6I5
В	257	SER	-	expression tag	UNP A0A1I6R6I5
В	258	HIS	-	expression tag	UNP A0A1I6R6I5
В	259	PRO	-	expression tag	UNP A0A1I6R6I5
В	260	GLN	-	expression tag	UNP A0A1I6R6I5
В	261	PHE	-	expression tag	UNP A0A1I6R6I5
В	262	GLU	-	expression tag	UNP A0A1I6R6I5
В	263	LYS	-	expression tag	UNP A0A1I6R6I5


#### $\bullet$ Molecule 2 is a protein called GldL.

Mol	Chain	Residues		Ato	ms		AltConf	Trace	
2	С	60	Total	С	N	О	S	0	0
	00	473	318	74	77	4	0	0	
2	D	55	Total	С	N	О	S	0	0
	ע	55	430	294	66	66	4	0	U
2	Е	50	Total	С	N	О	S	0	0
	<u> 1</u> 2	50	387	266	59	58	4		
2	F	50	Total	С	N	О	S	0	0
	Δ Γ	50	387	266	59	58	4		U
2	G	50	Total	С	N	О	S	0	0
	G	50	387	265	59	59	4	U	U




## 3 Residue-property plots (i)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.













# 4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	111727	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE	Depositor
	CORRECTION	
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{Å}^2)$	56	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K3 BIOQUANTUM (6k x 4k)	Depositor
Maximum map value	0.139	Depositor
Minimum map value	-0.079	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.005	Depositor
Recommended contour level	0.03	Depositor
Map size (Å)	212.992, 212.992, 212.992	wwPDB
Map dimensions	256, 256, 256	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.832, 0.832, 0.832	Depositor



# 5 Model quality (i)

#### 5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
WIOI		RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	A	0.34	0/1642	0.46	0/2214	
1	В	0.31	0/1629	0.45	0/2198	
2	С	0.37	0/485	0.47	0/656	
2	D	0.38	0/440	0.49	0/590	
2	Ε	0.36	0/396	0.49	0/532	
2	F	0.39	0/396	0.46	0/532	
2	G	0.38	0/396	0.46	0/533	
All	All	0.35	0/5384	0.46	0/7255	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

#### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1627	0	1674	22	0
1	В	1614	0	1658	18	0
2	С	473	0	483	7	0
2	D	430	0	454	6	0
2	Е	387	0	413	8	0
2	F	387	0	413	4	0
2	G	387	0	408	2	0
All	All	5305	0	5503	53	0



The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 53 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} \text{Interatomic} \\ \text{distance (Å)} \end{array}$	$\begin{array}{c} \text{Clash} \\ \text{overlap } (\text{\AA}) \end{array}$
1:A:210:HIS:HE1	1:A:214:LYS:HE2	1.54	0.71
1:B:9:ARG:NH1	2:C:47:GLU:OE2	2.17	0.71
1:A:51:GLN:NE2	1:A:55:ASP:OD1	2.24	0.71
1:A:210:HIS:CE1	1:A:214:LYS:HE2	2.27	0.70
1:B:138:ARG:NH2	1:B:164:ASP:OD1	2.30	0.65

There are no symmetry-related clashes.

#### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	$211/263\ (80\%)$	205 (97%)	6 (3%)	0	100	100
1	В	209/263~(80%)	200 (96%)	9 (4%)	0	100	100
2	$\mathbf{C}$	58/212 (27%)	57 (98%)	1 (2%)	0	100	100
2	D	53/212 (25%)	51 (96%)	2 (4%)	0	100	100
2	E	48/212 (23%)	47 (98%)	1 (2%)	0	100	100
2	F	48/212 (23%)	46 (96%)	2 (4%)	0	100	100
2	G	48/212 (23%)	47 (98%)	1 (2%)	0	100	100
All	All	675/1586 (43%)	653 (97%)	22 (3%)	0	100	100

There are no Ramachandran outliers to report.



#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	A	$175/212 \; (82\%)$	175 (100%)	0	100	100
1	В	$174/212 \ (82\%)$	173 (99%)	1 (1%)	86	95
2	С	48/170 (28%)	48 (100%)	0	100	100
2	D	43/170 (25%)	43 (100%)	0	100	100
2	E	39/170 (23%)	39 (100%)	0	100	100
2	F	39/170 (23%)	39 (100%)	0	100	100
2	G	39/170 (23%)	39 (100%)	0	100	100
All	All	557/1274 (44%)	556 (100%)	1 (0%)	93	98

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	В	27	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	210	HIS

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry (i)

There are no ligands in this entry.

### 5.7 Other polymers (i)

There are no such residues in this entry.

### 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.




## 6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-24958. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

### 6.1 Orthogonal projections (i)

#### 6.1.1 Primary map



#### 6.1.2 Raw map




The images above show the map projected in three orthogonal directions.




### 6.2 Central slices (i)


#### 6.2.1 Primary map








Y Index: 128




Z Index: 128


#### 6.2.2 Raw map



X Index: 128



Y Index: 128




Z Index: 128

The images above show central slices of the map in three orthogonal directions.




#### 6.3 Largest variance slices (i)


#### 6.3.1 Primary map








Y Index: 129




Z Index: 141


#### 6.3.2 Raw map



X Index: 127



Y Index: 129




Z Index: 141

The images above show the largest variance slices of the map in three orthogonal directions.




### 6.4 Orthogonal surface views (i)

#### 6.4.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.03. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

#### 6.4.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.




### 6.5 Mask visualisation (i)


This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure

#### $6.5.1 \quad \mathrm{emd} \, \underline{24958} \, \underline{\mathrm{msk}} \, \underline{1.\mathrm{map}} \, \, \underline{\mathrm{i}}$








# 7 Map analysis (i)

This section contains the results of statistical analysis of the map.


### 7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



#### 7.2 Volume estimate (i)




The volume at the recommended contour level is  $21 \text{ nm}^3$ ; this corresponds to an approximate mass of 19 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



### 7.3 Rotationally averaged power spectrum (i)




<sup>\*</sup>Reported resolution corresponds to spatial frequency of 0.333  $\rm \AA^{-1}$ 



## 8 Fourier-Shell correlation (i)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

#### 8.1 FSC (i)



\*Reported resolution corresponds to spatial frequency of 0.333  $\rm \mathring{A}^{-1}$ 



### 8.2 Resolution estimates (i)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
resolution estimate (A)	0.143	0.5	Half-bit
Reported by author	3.00	-	-
Author-provided FSC curve	3.04	3.60	3.17
Unmasked-calculated*	3.50	4.07	3.58


<sup>\*</sup>Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.50 differs from the reported value 3.0 by more than 10 %



# 9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-24958 and PDB model 7SAX. Per-residue inclusion information can be found in section 3 on page 6.


### 9.1 Map-model overlay (i)



The images above show the 3D surface view of the map at the recommended contour level 0.03 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.




### 9.2 Q-score mapped to coordinate model (i)



The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


#### 9.3 Atom inclusion mapped to coordinate model (i)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.03).



### 9.4 Atom inclusion (i)



At the recommended contour level, 78% of all backbone atoms, 69% of all non-hydrogen atoms, are inside the map.



# 9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (0.03) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.6870	0.5590
A	0.6930	0.5570
В	0.6252	0.5390
С	0.6709	0.5670
D	0.7103	0.5630
Е	0.7688	0.5930
F	0.7325	0.5820
G	0.7844	0.5810



