



wwPDB EM Validation Summary Report ⓘ

May 22, 2023 – 07:43 PM EDT

PDB ID : 8SAY
EMDB ID : EMD-40284
Title : CryoEM structure of DH270.3-CH848.10.17
Authors : Henderson, R.; Zhou, Y.; Stalls, V.; Bartesaghi, B.; Acharya, P.
Deposited on : 2023-04-02
Resolution : 3.40 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev50
Mogul : 1.8.5 (274361), CSD as541be (2020)
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.9
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.33


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	463	
1	F	463	
1	K	463	
2	B	132	
2	G	132	
2	L	132	
3	C	126	
3	H	126	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	M	126	95% 5%
4	D	110	88% 12% 8%
4	I	110	81% 19% 7%
4	N	110	90% 10% 7%
5	E	3	67% 33% 67%
5	P	3	67% 33% 67%
5	Q	3	67% 33% 33%
6	J	11	73% 27% 9%
6	R	11	82% 18% 9%
6	V	11	82% 18% 9%
7	O	5	60% 40% 60%
8	S	4	75% 25% 50%
8	W	4	75% 25% 50%
9	T	5	80% 60% 40%
10	U	2	50% 50% 50%
10	X	2	100% 50%

2 Entry composition i

There are 10 unique types of molecules in this entry. The entry contains 19902 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CH848.10.17 gp120.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	461	3607	2255	626	697	29	0	0
1	F	462	3615	2261	627	698	29	0	0
1	K	462	3615	2261	627	698	29	0	0

There are 30 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	31	ALA	-	expression tag	UNP A0A1W6IPB2
A	32	GLU	-	expression tag	UNP A0A1W6IPB2
A	33	ASN	-	expression tag	UNP A0A1W6IPB2
A	201	CYS	VAL	conflict	UNP A0A1W6IPB2
A	433	CYS	ALA	conflict	UNP A0A1W6IPB2
A	490	LYS	GLU	conflict	UNP A0A1W6IPB2
A	492	GLU	GLN	conflict	UNP A0A1W6IPB2
A	496	VAL	ILE	conflict	UNP A0A1W6IPB2
A	500	ARG	GLY	conflict	UNP A0A1W6IPB2
A	501	CYS	ALA	conflict	UNP A0A1W6IPB2
F	31	ALA	-	expression tag	UNP A0A1W6IPB2
F	32	GLU	-	expression tag	UNP A0A1W6IPB2
F	33	ASN	-	expression tag	UNP A0A1W6IPB2
F	201	CYS	VAL	conflict	UNP A0A1W6IPB2
F	433	CYS	ALA	conflict	UNP A0A1W6IPB2
F	490	LYS	GLU	conflict	UNP A0A1W6IPB2
F	492	GLU	GLN	conflict	UNP A0A1W6IPB2
F	496	VAL	ILE	conflict	UNP A0A1W6IPB2
F	500	ARG	GLY	conflict	UNP A0A1W6IPB2
F	501	CYS	ALA	conflict	UNP A0A1W6IPB2
K	31	ALA	-	expression tag	UNP A0A1W6IPB2
K	32	GLU	-	expression tag	UNP A0A1W6IPB2
K	33	ASN	-	expression tag	UNP A0A1W6IPB2
K	201	CYS	VAL	conflict	UNP A0A1W6IPB2

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
K	433	CYS	ALA	conflict	UNP A0A1W6IPB2
K	490	LYS	GLU	conflict	UNP A0A1W6IPB2
K	492	GLU	GLN	conflict	UNP A0A1W6IPB2
K	496	VAL	ILE	conflict	UNP A0A1W6IPB2
K	500	ARG	GLY	conflict	UNP A0A1W6IPB2
K	501	CYS	ALA	conflict	UNP A0A1W6IPB2

- Molecule 2 is a protein called CH848.10.17 gp41.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	B	122	Total	C	N	O	S	0	0
			967	609	168	184	6		
2	G	122	Total	C	N	O	S	0	0
			967	609	168	184	6		
2	L	122	Total	C	N	O	S	0	0
			967	609	168	184	6		

- Molecule 3 is a protein called DH270.3 variable heavy chain.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	C	126	Total	C	N	O	S	0	0
			991	631	169	186	5		
3	H	126	Total	C	N	O	S	0	0
			991	631	169	186	5		
3	M	126	Total	C	N	O	S	0	0
			991	631	169	186	5		

- Molecule 4 is a protein called DH270.3 variable light chain.


Mol	Chain	Residues	Atoms					AltConf	Trace
4	D	110	Total	C	N	O	S	0	0
			805	500	136	166	3		
4	I	110	Total	C	N	O	S	0	0
			805	500	136	166	3		
4	N	110	Total	C	N	O	S	0	0
			805	500	136	166	3		

- Molecule 5 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
5	E	3	39	22	2	15	0	0
5	P	3	39	22	2	15	0	0
5	Q	3	39	22	2	15	0	0

- Molecule 6 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
6	J	11	127	70	2	55	0	0
6	R	11	127	70	2	55	0	0
6	V	11	127	70	2	55	0	0

- Molecule 7 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
7	O	5	61	34	2	25	0	0


- Molecule 8 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.

e-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
8	S	4	50	28	2	20	0	0
8	W	4	50	28	2	20	0	0

- Molecule 9 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
9	T	5	61	34	2	25	0	0

- Molecule 10 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
10	U	2	28	16	2	10	0	0
10	X	2	28	16	2	10	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: CH848.10.17 gp120



- Molecule 1: CH848.10.17 gp120





- Molecule 1: CH848.10.17 gp120








- Molecule 2: CH848.10.17 gp41



- Molecule 2: CH848.10.17 gp41




- Molecule 2: CH848.10.17 gp41




- Molecule 3: DH270.3 variable heavy chain




- Molecule 3: DH270.3 variable heavy chain




- Molecule 3: DH270.3 variable heavy chain




- Molecule 4: DH270.3 variable light chain




- Molecule 4: DH270.3 variable light chain




- Molecule 4: DH270.3 variable light chain




- Molecule 5: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 5: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 5: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose




- Molecule 6: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-

(1-2)-alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose




- Molecule 6: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose





- Molecule 6: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose




- Molecule 7: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose





- Molecule 8: alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 8: alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 9: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 10: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 10: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	76274	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	60	Depositor
Minimum defocus (nm)	700	Depositor
Maximum defocus (nm)	4100	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	0.553	Depositor
Minimum map value	-0.346	Depositor
Average map value	0.002	Depositor
Map value standard deviation	0.024	Depositor
Recommended contour level	0.058	Depositor
Map size (Å)	204.672, 204.672, 204.672	wwPDB
Map dimensions	192, 192, 192	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.066, 1.066, 1.066	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MAN, NAG, BMA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.30	0/3683	0.61	0/5008
1	F	0.29	0/3692	0.60	0/5022
1	K	0.30	0/3692	0.61	0/5022
2	B	0.29	0/985	0.64	0/1337
2	G	0.30	0/985	0.66	1/1337 (0.1%)
2	L	0.30	0/985	0.62	0/1337
3	C	0.32	0/1018	0.62	0/1385
3	H	0.31	0/1018	0.61	0/1385
3	M	0.32	0/1018	0.64	0/1385
4	D	0.30	0/821	0.64	0/1116
4	I	0.29	0/821	0.62	0/1116
4	N	0.30	0/821	0.61	0/1116
All	All	0.30	0/19539	0.62	1/26566 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
2	B	0	1
4	D	0	1
4	I	0	1
4	N	0	1
All	All	0	5

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	G	581	LEU	CA-CB-CG	6.37	129.95	115.30

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	430	VAL	Peptide
2	B	569	THR	Peptide
4	D	309	ARG	Peptide
4	I	309	ARG	Peptide
4	N	309	ARG	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3607	0	3518	28	0
1	F	3615	0	3530	21	0
1	K	3615	0	3530	24	0
2	B	967	0	946	7	0
2	G	967	0	946	11	0
2	L	967	0	946	4	0
3	C	991	0	958	2	0
3	H	991	0	958	3	0
3	M	991	0	958	3	0
4	D	805	0	780	6	0
4	I	805	0	780	11	0
4	N	805	0	780	6	0
5	E	39	0	34	0	0
5	P	39	0	34	0	0
5	Q	39	0	34	1	0
6	J	127	0	106	0	0
6	R	127	0	106	0	0
6	V	127	0	106	0	0
7	O	61	0	52	0	0
8	S	50	0	43	0	0
8	W	50	0	43	0	0
9	T	61	0	52	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
10	U	28	0	25	0	0
10	X	28	0	25	0	0
All	All	19902	0	19290	111	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 111 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:37:THR:OG1	2:G:604:CYS:O	1.98	0.81
1:A:37:THR:OG1	2:B:604:CYS:O	1.99	0.80
1:A:195:ASN:ND2	1:A:201:CYS:SG	2.61	0.74
1:F:36:VAL:O	2:G:606:THR:OG1	2.04	0.74
1:A:256:SER:OG	1:A:259:LEU:O	2.06	0.73

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	457/463 (99%)	430 (94%)	26 (6%)	1 (0%)	47 78
1	F	460/463 (99%)	433 (94%)	26 (6%)	1 (0%)	47 78
1	K	460/463 (99%)	433 (94%)	26 (6%)	1 (0%)	47 78
2	B	120/132 (91%)	113 (94%)	7 (6%)	0	100 100
2	G	120/132 (91%)	114 (95%)	6 (5%)	0	100 100
2	L	120/132 (91%)	111 (92%)	8 (7%)	1 (1%)	19 51
3	C	124/126 (98%)	114 (92%)	9 (7%)	1 (1%)	19 51
3	H	124/126 (98%)	113 (91%)	10 (8%)	1 (1%)	19 51

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	M	124/126 (98%)	114 (92%)	9 (7%)	1 (1%)	19	51
4	D	108/110 (98%)	97 (90%)	11 (10%)	0	100	100
4	I	108/110 (98%)	98 (91%)	10 (9%)	0	100	100
4	N	108/110 (98%)	97 (90%)	11 (10%)	0	100	100
All	All	2433/2493 (98%)	2267 (93%)	159 (6%)	7 (0%)	44	72

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	123	THR
1	F	123	THR
1	K	123	THR
3	C	61	PRO
3	H	61	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	410/411 (100%)	410 (100%)	0	100	100
1	F	411/411 (100%)	411 (100%)	0	100	100
1	K	411/411 (100%)	411 (100%)	0	100	100
2	B	104/110 (94%)	104 (100%)	0	100	100
2	G	104/110 (94%)	104 (100%)	0	100	100
2	L	104/110 (94%)	104 (100%)	0	100	100
3	C	106/106 (100%)	106 (100%)	0	100	100
3	H	106/106 (100%)	106 (100%)	0	100	100
3	M	106/106 (100%)	106 (100%)	0	100	100
4	D	91/91 (100%)	91 (100%)	0	100	100
4	I	91/91 (100%)	91 (100%)	0	100	100
4	N	91/91 (100%)	91 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	2135/2154 (99%)	2135 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	F	103	GLN
1	K	103	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

64 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	NAG	E	1	1,5	14,14,15	0.95	1 (7%)	17,19,21	1.14	1 (5%)
5	NAG	E	2	5	14,14,15	0.21	0	17,19,21	0.48	0
5	BMA	E	3	5	11,11,12	0.82	0	15,15,17	1.06	1 (6%)
6	NAG	J	1	6,1	14,14,15	0.22	0	17,19,21	0.51	0
6	MAN	J	10	6	11,11,12	0.63	0	15,15,17	1.23	3 (20%)
6	MAN	J	11	6	11,11,12	0.75	0	15,15,17	0.90	1 (6%)
6	NAG	J	2	6	14,14,15	0.27	0	17,19,21	0.44	0
6	BMA	J	3	6	11,11,12	0.64	0	15,15,17	0.98	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	MAN	J	4	6	11,11,12	0.84	0	15,15,17	1.34	3 (20%)
6	MAN	J	5	6	11,11,12	1.09	1 (9%)	15,15,17	1.34	2 (13%)
6	MAN	J	6	6	11,11,12	0.70	0	15,15,17	1.14	2 (13%)
6	MAN	J	7	6	11,11,12	0.60	0	15,15,17	1.17	1 (6%)
6	MAN	J	8	6	11,11,12	0.81	1 (9%)	15,15,17	1.12	2 (13%)
6	MAN	J	9	6	11,11,12	0.75	1 (9%)	15,15,17	1.14	2 (13%)
7	NAG	O	1	7,1	14,14,15	0.37	0	17,19,21	0.51	0
7	NAG	O	2	7	14,14,15	0.28	0	17,19,21	0.49	0
7	BMA	O	3	7	11,11,12	0.55	0	15,15,17	1.00	0
7	MAN	O	4	7	11,11,12	0.70	0	15,15,17	1.30	2 (13%)
7	MAN	O	5	7	11,11,12	1.05	1 (9%)	15,15,17	1.19	2 (13%)
5	NAG	P	1	1,5	14,14,15	0.23	0	17,19,21	0.49	0
5	NAG	P	2	5	14,14,15	0.34	0	17,19,21	0.49	0
5	BMA	P	3	5	11,11,12	1.08	1 (9%)	15,15,17	1.29	3 (20%)
5	NAG	Q	1	1,5	14,14,15	1.13	1 (7%)	17,19,21	1.18	2 (11%)
5	NAG	Q	2	5	14,14,15	0.19	0	17,19,21	0.45	0
5	BMA	Q	3	5	11,11,12	0.79	0	15,15,17	0.95	1 (6%)
6	NAG	R	1	6,1	14,14,15	0.20	0	17,19,21	0.49	0
6	MAN	R	10	6	11,11,12	0.65	0	15,15,17	1.24	3 (20%)
6	MAN	R	11	6	11,11,12	0.74	0	15,15,17	0.89	1 (6%)
6	NAG	R	2	6	14,14,15	0.27	0	17,19,21	0.45	0
6	BMA	R	3	6	11,11,12	0.58	0	15,15,17	0.97	1 (6%)
6	MAN	R	4	6	11,11,12	0.95	0	15,15,17	1.35	3 (20%)
6	MAN	R	5	6	11,11,12	1.10	0	15,15,17	1.40	3 (20%)
6	MAN	R	6	6	11,11,12	0.69	0	15,15,17	1.08	2 (13%)
6	MAN	R	7	6	11,11,12	0.65	0	15,15,17	1.22	1 (6%)
6	MAN	R	8	6	11,11,12	0.76	0	15,15,17	1.05	2 (13%)
6	MAN	R	9	6	11,11,12	0.76	1 (9%)	15,15,17	1.15	2 (13%)
8	NAG	S	1	1,8	14,14,15	0.37	0	17,19,21	0.53	0
8	NAG	S	2	8	14,14,15	0.26	0	17,19,21	0.51	0
8	BMA	S	3	8	11,11,12	0.59	0	15,15,17	0.97	0
8	MAN	S	4	8	11,11,12	0.69	0	15,15,17	1.25	2 (13%)
9	NAG	T	1	9,1	14,14,15	0.22	0	17,19,21	0.46	0
9	NAG	T	2	9	14,14,15	0.32	0	17,19,21	0.51	0
9	BMA	T	3	9	11,11,12	1.00	0	15,15,17	1.14	2 (13%)
9	MAN	T	4	9	11,11,12	0.81	1 (9%)	15,15,17	1.30	2 (13%)
9	MAN	T	5	9	11,11,12	0.91	1 (9%)	15,15,17	1.28	2 (13%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
10	NAG	U	1	10,1	14,14,15	0.25	0	17,19,21	0.97	1 (5%)
10	NAG	U	2	10	14,14,15	0.47	0	17,19,21	0.62	0
6	NAG	V	1	6,1	14,14,15	0.19	0	17,19,21	0.49	0
6	MAN	V	10	6	11,11,12	0.67	0	15,15,17	1.23	3 (20%)
6	MAN	V	11	6	11,11,12	0.73	0	15,15,17	0.90	1 (6%)
6	NAG	V	2	6	14,14,15	0.22	0	17,19,21	0.46	0
6	BMA	V	3	6	11,11,12	0.53	0	15,15,17	1.03	2 (13%)
6	MAN	V	4	6	11,11,12	0.89	0	15,15,17	1.41	2 (13%)
6	MAN	V	5	6	11,11,12	1.11	1 (9%)	15,15,17	1.41	3 (20%)
6	MAN	V	6	6	11,11,12	0.65	0	15,15,17	1.21	2 (13%)
6	MAN	V	7	6	11,11,12	0.58	0	15,15,17	1.17	1 (6%)
6	MAN	V	8	6	11,11,12	0.76	0	15,15,17	1.08	2 (13%)
6	MAN	V	9	6	11,11,12	0.73	1 (9%)	15,15,17	1.16	2 (13%)
8	NAG	W	1	1,8	14,14,15	0.30	0	17,19,21	0.50	0
8	NAG	W	2	8	14,14,15	0.29	0	17,19,21	0.54	0
8	BMA	W	3	8	11,11,12	0.54	0	15,15,17	0.91	0
8	MAN	W	4	8	11,11,12	0.96	1 (9%)	15,15,17	1.35	3 (20%)
10	NAG	X	1	10,1	14,14,15	0.35	0	17,19,21	0.45	0
10	NAG	X	2	10	14,14,15	0.39	0	17,19,21	0.49	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	E	1	1,5	-	0/6/23/26	0/1/1/1
5	NAG	E	2	5	-	0/6/23/26	0/1/1/1
5	BMA	E	3	5	-	2/2/19/22	0/1/1/1
6	NAG	J	1	6,1	-	2/6/23/26	0/1/1/1
6	MAN	J	10	6	-	0/2/19/22	0/1/1/1
6	MAN	J	11	6	-	1/2/19/22	0/1/1/1
6	NAG	J	2	6	-	0/6/23/26	0/1/1/1
6	BMA	J	3	6	-	0/2/19/22	0/1/1/1
6	MAN	J	4	6	-	1/2/19/22	0/1/1/1
6	MAN	J	5	6	-	2/2/19/22	0/1/1/1
6	MAN	J	6	6	-	0/2/19/22	0/1/1/1
6	MAN	J	7	6	-	0/2/19/22	0/1/1/1
6	MAN	J	8	6	-	0/2/19/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	MAN	J	9	6	-	2/2/19/22	0/1/1/1
7	NAG	O	1	7,1	-	0/6/23/26	0/1/1/1
7	NAG	O	2	7	-	0/6/23/26	0/1/1/1
7	BMA	O	3	7	-	1/2/19/22	0/1/1/1
7	MAN	O	4	7	-	0/2/19/22	0/1/1/1
7	MAN	O	5	7	-	1/2/19/22	0/1/1/1
5	NAG	P	1	1,5	-	2/6/23/26	0/1/1/1
5	NAG	P	2	5	-	1/6/23/26	0/1/1/1
5	BMA	P	3	5	-	1/2/19/22	0/1/1/1
5	NAG	Q	1	1,5	-	2/6/23/26	0/1/1/1
5	NAG	Q	2	5	-	0/6/23/26	0/1/1/1
5	BMA	Q	3	5	-	1/2/19/22	0/1/1/1
6	NAG	R	1	6,1	-	1/6/23/26	0/1/1/1
6	MAN	R	10	6	-	0/2/19/22	0/1/1/1
6	MAN	R	11	6	-	1/2/19/22	0/1/1/1
6	NAG	R	2	6	-	0/6/23/26	0/1/1/1
6	BMA	R	3	6	-	0/2/19/22	0/1/1/1
6	MAN	R	4	6	-	2/2/19/22	0/1/1/1
6	MAN	R	5	6	-	2/2/19/22	0/1/1/1
6	MAN	R	6	6	-	0/2/19/22	0/1/1/1
6	MAN	R	7	6	-	0/2/19/22	0/1/1/1
6	MAN	R	8	6	-	0/2/19/22	0/1/1/1
6	MAN	R	9	6	-	2/2/19/22	0/1/1/1
8	NAG	S	1	1,8	-	0/6/23/26	0/1/1/1
8	NAG	S	2	8	-	0/6/23/26	0/1/1/1
8	BMA	S	3	8	-	1/2/19/22	0/1/1/1
8	MAN	S	4	8	-	0/2/19/22	0/1/1/1
9	NAG	T	1	9,1	-	2/6/23/26	0/1/1/1
9	NAG	T	2	9	-	1/6/23/26	0/1/1/1
9	BMA	T	3	9	-	1/2/19/22	0/1/1/1
9	MAN	T	4	9	-	0/2/19/22	0/1/1/1
9	MAN	T	5	9	-	2/2/19/22	0/1/1/1
10	NAG	U	1	10,1	-	0/6/23/26	0/1/1/1
10	NAG	U	2	10	-	0/6/23/26	0/1/1/1
6	NAG	V	1	6,1	-	0/6/23/26	0/1/1/1
6	MAN	V	10	6	-	0/2/19/22	0/1/1/1
6	MAN	V	11	6	-	1/2/19/22	0/1/1/1
6	NAG	V	2	6	-	0/6/23/26	0/1/1/1
6	BMA	V	3	6	-	0/2/19/22	0/1/1/1
6	MAN	V	4	6	-	2/2/19/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	MAN	V	5	6	-	2/2/19/22	0/1/1/1
6	MAN	V	6	6	-	0/2/19/22	0/1/1/1
6	MAN	V	7	6	-	0/2/19/22	0/1/1/1
6	MAN	V	8	6	-	0/2/19/22	0/1/1/1
6	MAN	V	9	6	-	2/2/19/22	0/1/1/1
8	NAG	W	1	1,8	-	0/6/23/26	0/1/1/1
8	NAG	W	2	8	-	0/6/23/26	0/1/1/1
8	BMA	W	3	8	-	2/2/19/22	0/1/1/1
8	MAN	W	4	8	-	0/2/19/22	0/1/1/1
10	NAG	X	1	10,1	-	2/6/23/26	0/1/1/1
10	NAG	X	2	10	-	1/6/23/26	0/1/1/1

The worst 5 of 13 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	Q	1	NAG	O5-C1	-3.84	1.37	1.43
5	E	1	NAG	O5-C1	-3.09	1.38	1.43
8	W	4	MAN	C1-C2	2.54	1.58	1.52
7	O	5	MAN	C1-C2	2.31	1.57	1.52
5	P	3	BMA	O5-C1	-2.31	1.40	1.43

The worst 5 of 76 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
9	T	5	MAN	C1-O5-C5	3.79	117.33	112.19
6	V	6	MAN	C1-O5-C5	3.54	116.98	112.19
6	R	5	MAN	O2-C2-C3	-3.49	103.15	110.14
6	R	7	MAN	C1-O5-C5	3.47	116.89	112.19
9	T	4	MAN	C1-O5-C5	3.45	116.87	112.19

There are no chirality outliers.

5 of 46 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
6	R	5	MAN	O5-C5-C6-O6
6	V	5	MAN	O5-C5-C6-O6
6	J	5	MAN	O5-C5-C6-O6
6	J	9	MAN	C4-C5-C6-O6
5	P	1	NAG	O5-C5-C6-O6



There are no ring outliers.


1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	Q	1	NAG	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.























5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-40284. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 96




Y Index: 96




Z Index: 96


6.2.2 Raw map



X Index: 96



Y Index: 96




Z Index: 96


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 85




Y Index: 90




Z Index: 113


6.3.2 Raw map



X Index: 85



Y Index: 90




Z Index: 108


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.058. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.6.1 emd_40284_msk_1.map [i](#)



X



Y




Z

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [\(i\)](#)



The volume at the recommended contour level is 178 nm^3 ; this corresponds to an approximate mass of 161 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum i




*Reported resolution corresponds to spatial frequency of 0.294 \AA^{-1}

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.294 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.40	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	3.55	3.86	3.59

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-40284 and PDB model 8SAY. Per-residue inclusion information can be found in section 3 on page 8.

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.058 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.058).



















































9.4 Atom inclusion [i](#)



At the recommended contour level, 84% of all backbone atoms, 76% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.058) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.7650	 0.4060
A	 0.7650	 0.4190
B	 0.7460	 0.3750
C	 0.8550	 0.4470
D	 0.7610	 0.3710
E	 0.3590	 0.2050
F	 0.7610	 0.4200
G	 0.7520	 0.3750
H	 0.8440	 0.4440
I	 0.7560	 0.3620
J	 0.7640	 0.3850
K	 0.7610	 0.4160
L	 0.7530	 0.3740
M	 0.8450	 0.4480
N	 0.7510	 0.3730
O	 0.3930	 0.2460
P	 0.4100	 0.2620
Q	 0.3330	 0.1730
R	 0.7720	 0.3940
S	 0.4400	 0.3070
T	 0.2790	 0.2080
U	 0.5360	 0.2180
V	 0.7720	 0.3920
W	 0.4600	 0.3550
X	 0.6070	 0.3690

