



wwPDB EM Validation Summary Report ⓘ

Nov 21, 2022 – 03:07 PM EST

PDB ID : 7SGR
EMDB ID : EMD-25116
Title : Structure of hemolysin A secretion system HlyB/D complex
Authors : Zhao, H.; Chen, J.
Deposited on : 2021-10-07
Resolution : 2.90 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev43
Mogul : 1.8.5 (274361), CSD as541be (2020)
MolProbity : 4.02b-467
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.9
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.31.3


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.







Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	707	
1	B	707	
1	E	707	
1	F	707	
1	I	707	
1	J	707	
2	C	356	
2	D	356	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	G	356	
2	H	356	
2	K	356	
2	L	356	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 34989 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Alpha-hemolysin translocation ATP-binding protein HlyB.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	J	570	Total 4505	C 2923	N 773	O 797	S 12	0	0
1	I	700	Total 5549	C 3595	N 953	O 989	S 12	0	0
1	A	700	Total 5543	C 3591	N 952	O 988	S 12	0	0
1	B	570	Total 4509	C 2926	N 774	O 797	S 12	0	0
1	E	700	Total 5535	C 3583	N 953	O 987	S 12	0	0
1	F	570	Total 4509	C 2926	N 774	O 797	S 12	0	0

- Molecule 2 is a protein called Membrane fusion protein (MFP) family protein, Hemolysin secretion protein D, chromosomal.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	H	69	Total 581	C 385	N 97	O 98	S 1	0	0
2	L	69	Total 581	C 385	N 97	O 98	S 1	0	0
2	D	69	Total 581	C 385	N 97	O 98	S 1	0	0
2	C	166	Total 1035	C 648	N 186	O 199	S 2	0	0
2	G	165	Total 1026	C 640	N 185	O 199	S 2	0	0
2	K	49	Total 396	C 261	N 65	O 69	S 1	0	0

- Molecule 3 is [(2 {R})-1-[2-azanylethoxy(oxidanyl)phosphoryl]oxy-3-hexadecanoyloxy-prop an-2-yl] ({Z})-octadec-9-enoate (three-letter code: 6OU) (formula: C₃₉H₇₆NO₈P).



Mol	Chain	Residues	Atoms				AltConf
3	J	1	Total	C			0
			82	82			
3	J	1	Total	C			0
			82	82			
3	J	1	Total	C			0
			82	82			
3	J	1	Total	C			0
			82	82			
3	J	1	Total	C			0
			82	82			
3	J	1	Total	C			0
			82	82			
3	I	1	Total	C	O	P	0
			131	122	8	1	
3	I	1	Total	C	O	P	0
			131	122	8	1	
3	I	1	Total	C	O	P	0
			131	122	8	1	
3	I	1	Total	C	O	P	0
			131	122	8	1	
3	I	1	Total	C	O	P	0
			131	122	8	1	
3	I	1	Total	C	O	P	0
			131	122	8	1	
3	I	1	Total	C	O	P	0
			131	122	8	1	

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				AltConf	
			Total	C	O	P		
3	I	1	131	122	8	1	0	
3	A	1	147	114	1	29	3	0
3	A	1	147	114	1	29	3	0
3	A	1	147	114	1	29	3	0
3	A	1	147	114	1	29	3	0
3	A	1	147	114	1	29	3	0
3	A	1	147	114	1	29	3	0
3	A	1	147	114	1	29	3	0
3	B	1	26	26				0
3	B	1	26	26				0
3	E	1	38	38				0
3	E	1	38	38				0
3	E	1	38	38				0
3	E	1	38	38				0
3	F	1	93	83	10			0
3	F	1	93	83	10			0
3	F	1	93	83	10			0
3	F	1	93	83	10			0
3	H	1	53	44	8	1		0
3	H	1	53	44	8	1		0
3	L	1	29	29				0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms				AltConf
3	L	1	Total	C			0
			29	29			
3	K	1	Total	C	O	P	0
			40	31	8	1	


3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Alpha-hemolysin translocation ATP-binding protein HlyB





• Molecule 1: Alpha-hemolysin translocation ATP-binding protein HlyB







• Molecule 1: Alpha-hemolysin translocation ATP-binding protein HlyB




• Molecule 1: Alpha-hemolysin translocation ATP-binding protein HlyB




• Molecule 1: Alpha-hemolysin translocation ATP-binding protein HlyB







● Molecule 1: Alpha-hemolysin translocation ATP-binding protein HlyB




● Molecule 2: Membrane fusion protein (MFP) family protein, Hemolysin secretion protein D, chromosomal






- Molecule 2: Membrane fusion protein (MFP) family protein, Hemolysin secretion protein D, chromosomal

Chain L: 15% 5% 81%




- Molecule 2: Membrane fusion protein (MFP) family protein, Hemolysin secretion protein D, chromosomal



Chain D: 15% 81%




- Molecule 2: Membrane fusion protein (MFP) family protein, Hemolysin secretion protein D, chromosomal



- Molecule 2: Membrane fusion protein (MFP) family protein, Hemolysin secretion protein D, chromosomal



- Molecule 2: Membrane fusion protein (MFP) family protein, Hemolysin secretion protein D, chromosomal



THR
LEU
SER
GLY
ARG
SER
LYS
GLU
PRO
LYS
ILE
LEU
ASN
GLU
ASN

THR
ARG
TYR
THR
ILE
LEU
SER
ARG
SER
ILE
LEU
ASN
LYS
LEU
PRO
GLU
VAL
LEU
LYS
TYR
PHE
GLN
SER
ASN
VAL
VAL
SER
SER
GLY
GLU
GLU
VAL
LEU
LEU
ARG
LYS
LEU
LEU
THR
THR
SER
ALA
LEU
GLY
ILE
LYS
ALA
GLY
GLU
GLN
ALA
ASP

LYS
LYS
ARG
ALA
GLU
ARG
LEU
THR
THR
ILE
LEU
ALA
LYS
ARG
ILE
ASN
ASN
ARG
TYR
GLU
ASN
LEU
ASN
LEU
SER
ALA
SER
ARG
VAL
GLU
GLU
LYS
SER
ARG
LEU
GLY
LEU
LEU
VAL
VAL
THR
ALA
LEU
VAL
VAL
ASN
ASN
LYS
ASP
ILE
GLY
SER
PHE
THR
ILE
ILE
ASN
VAL
GLY
ASN
GLN
ASN
ALA
ILE
ILE
LYS
VAL
VAL
ALA

PHE
PRO
TYR
THR
ARG
TYR
GLY
TYR
LEU
VAL
TYR
GLY
LYS
VAL
LYS
ASN
ASN
ILE
LEU
ASN
GLU
ASP
ALA
ASP
GLN
LYS
LEU
GLY
LEU
VAL
PHE
ASN
VAL
ILE
VAL
VAL
VAL
GLU
GLU
ASN
ASP
LEU
SER
THR
GLY
ASN
LYS
HIS
PHE
ILE
PRO
LEU
SER
SER
GLY
MET
ALA
VAL
VAL
THR
ALA
ILE

LYS
THR
GLY
MET
SER
SER
VAL
ILE
SER
SER
LEU
LEU
SER
PRO
LEU
GLU
SER
SER
VAL
THR
GLU
SER
LEU
HIS
GLU
ARG

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	136123	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	70	Depositor
Minimum defocus (nm)	1000	Depositor
Maximum defocus (nm)	3000	Depositor
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	4.434	Depositor
Minimum map value	-2.384	Depositor
Average map value	-0.001	Depositor
Map value standard deviation	0.104	Depositor
Recommended contour level	0.4	Depositor
Map size (Å)	410.88, 410.88, 410.88	wwPDB
Map dimensions	384, 384, 384	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.07, 1.07, 1.07	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 6OU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.30	0/5651	0.49	0/7663
1	B	0.25	0/4596	0.48	0/6232
1	E	0.26	0/5641	0.47	0/7648
1	F	0.27	0/4596	0.48	0/6232
1	I	0.25	0/5656	0.47	0/7669
1	J	0.25	0/4592	0.47	0/6228
2	C	0.25	0/1044	0.48	0/1433
2	D	0.24	0/595	0.53	0/807
2	G	0.24	0/1035	0.45	0/1420
2	H	0.24	0/595	0.51	0/807
2	K	0.23	0/404	0.47	0/550
2	L	0.25	0/595	0.53	0/807
All	All	0.26	0/35000	0.48	0/47496

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5543	0	5714	62	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	4509	0	4683	52	0
1	E	5535	0	5706	76	0
1	F	4509	0	4683	80	0
1	I	5549	0	5734	62	0
1	J	4505	0	4672	60	0
2	C	1035	0	815	9	0
2	D	581	0	609	12	0
2	G	1026	0	797	23	0
2	H	581	0	609	8	0
2	K	396	0	416	5	0
2	L	581	0	609	16	0
3	A	147	0	0	0	0
3	B	26	0	0	0	0
3	E	38	0	0	0	0
3	F	93	0	0	0	0
3	H	53	0	0	0	0
3	I	131	0	0	0	0
3	J	82	0	0	0	0
3	K	40	0	0	0	0
3	L	29	0	0	0	0
All	All	34989	0	35047	421	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 421 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:128:ARG:HH21	2:L:47:GLU:HG2	1.42	0.85
1:F:315:LEU:HD11	1:F:370:VAL:HG13	1.70	0.72
1:I:128:ARG:NH2	2:D:42:LEU:O	2.24	0.70
1:I:282:PHE:HB2	1:I:420:ILE:HD11	1.75	0.69
1:A:283:PHE:HZ	1:A:301:LEU:HD11	1.56	0.68

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	698/707 (99%)	664 (95%)	34 (5%)	0	100	100
1	B	568/707 (80%)	543 (96%)	25 (4%)	0	100	100
1	E	698/707 (99%)	674 (97%)	24 (3%)	0	100	100
1	F	568/707 (80%)	547 (96%)	21 (4%)	0	100	100
1	I	698/707 (99%)	663 (95%)	35 (5%)	0	100	100
1	J	568/707 (80%)	545 (96%)	23 (4%)	0	100	100
2	C	162/356 (46%)	147 (91%)	15 (9%)	0	100	100
2	D	67/356 (19%)	65 (97%)	2 (3%)	0	100	100
2	G	161/356 (45%)	141 (88%)	20 (12%)	0	100	100
2	H	67/356 (19%)	66 (98%)	1 (2%)	0	100	100
2	K	47/356 (13%)	46 (98%)	1 (2%)	0	100	100
2	L	67/356 (19%)	64 (96%)	3 (4%)	0	100	100
All	All	4369/6378 (68%)	4165 (95%)	204 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	603/615 (98%)	603 (100%)	0	100	100
1	B	494/615 (80%)	494 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	601/615 (98%)	601 (100%)	0	100	100
1	F	494/615 (80%)	493 (100%)	1 (0%)	93	98
1	I	605/615 (98%)	604 (100%)	1 (0%)	93	98
1	J	493/615 (80%)	493 (100%)	0	100	100
2	C	67/324 (21%)	67 (100%)	0	100	100
2	D	65/324 (20%)	65 (100%)	0	100	100
2	G	66/324 (20%)	66 (100%)	0	100	100
2	H	65/324 (20%)	65 (100%)	0	100	100
2	K	45/324 (14%)	45 (100%)	0	100	100
2	L	65/324 (20%)	65 (100%)	0	100	100
All	All	3663/5634 (65%)	3661 (100%)	2 (0%)	93	98

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	I	587	PHE
1	F	587	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	121	HIS
1	E	539	ASN
1	F	585	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry

37 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	6OU	A	803	-	32,32,48	1.45	6 (18%)	36,37,53	1.18	2 (5%)
3	6OU	F	803	-	31,31,48	1.32	3 (9%)	33,33,53	1.19	2 (6%)
3	6OU	F	804	-	11,11,48	0.70	0	10,10,53	0.81	0
3	6OU	I	802	-	9,9,48	0.74	0	8,8,53	0.79	0
3	6OU	J	803	-	11,11,48	0.68	0	10,10,53	0.82	0
3	6OU	B	801	-	13,13,48	0.80	0	12,12,53	0.74	0
3	6OU	F	801	-	31,31,48	1.34	4 (12%)	33,33,53	1.22	2 (6%)
3	6OU	E	803	-	7,7,48	0.72	0	6,6,53	0.72	0
3	6OU	H	402	-	11,11,48	0.70	0	10,10,53	0.81	0
3	6OU	I	806	-	10,10,48	0.67	0	9,9,53	0.81	0
3	6OU	J	801	-	16,16,48	1.43	2 (12%)	15,15,53	0.96	0
3	6OU	I	808	-	11,11,48	0.70	0	10,10,53	0.84	0
3	6OU	L	401	-	16,16,48	1.42	2 (12%)	15,15,53	1.03	1 (6%)
3	6OU	F	802	-	16,16,48	1.42	2 (12%)	15,15,53	0.96	1 (6%)
3	6OU	I	807	-	11,11,48	0.70	0	10,10,53	0.83	0
3	6OU	J	806	-	11,11,48	0.70	0	10,10,53	0.83	0
3	6OU	E	804	-	12,12,48	0.68	0	11,11,53	0.86	0
3	6OU	H	401	-	40,40,48	1.33	5 (12%)	44,45,53	1.13	2 (4%)
3	6OU	A	804	-	19,19,48	1.58	3 (15%)	21,21,53	1.37	2 (9%)
3	6OU	J	802	-	16,16,48	1.42	2 (12%)	15,15,53	0.98	1 (6%)
3	6OU	A	801	-	31,31,48	1.39	5 (16%)	34,36,53	1.15	2 (5%)
3	6OU	L	402	-	11,11,48	0.70	0	10,10,53	0.84	0
3	6OU	A	806	-	11,11,48	0.69	0	10,10,53	0.85	0
3	6OU	J	804	-	13,13,48	0.71	0	12,12,53	0.87	0
3	6OU	A	807	-	24,24,48	1.62	4 (16%)	28,29,53	1.26	2 (7%)
3	6OU	E	801	-	8,8,48	0.74	0	7,7,53	0.76	0
3	6OU	I	803	-	33,33,48	1.43	6 (18%)	37,38,53	1.17	2 (5%)
3	6OU	K	401	-	39,39,48	1.35	5 (12%)	43,44,53	1.09	2 (4%)
3	6OU	I	801	-	16,16,48	1.42	2 (12%)	15,15,53	0.97	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	6OU	I	804	-	10,10,48	0.67	0	9,9,53	0.84	0
3	6OU	A	805	-	7,7,48	0.64	0	6,6,53	0.73	0
3	6OU	I	809	-	12,12,48	0.70	0	11,11,53	0.83	0
3	6OU	B	802	-	11,11,48	0.69	0	10,10,53	0.83	0
3	6OU	J	805	-	9,9,48	0.74	0	8,8,53	0.80	0
3	6OU	I	805	-	10,10,48	0.67	0	9,9,53	0.81	0
3	6OU	E	802	-	7,7,48	0.71	0	6,6,53	0.74	0
3	6OU	A	802	-	16,16,48	1.43	2 (12%)	15,15,53	0.91	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	6OU	A	803	-	-	18/34/34/52	-
3	6OU	F	803	-	-	16/33/33/52	-
3	6OU	F	804	-	-	4/9/9/52	-
3	6OU	I	802	-	-	4/7/7/52	-
3	6OU	J	803	-	-	5/9/9/52	-
3	6OU	B	801	-	-	4/11/11/52	-
3	6OU	F	801	-	-	9/33/33/52	-
3	6OU	E	803	-	-	3/5/5/52	-
3	6OU	H	402	-	-	4/9/9/52	-
3	6OU	I	806	-	-	6/8/8/52	-
3	6OU	J	801	-	-	7/14/14/52	-
3	6OU	I	808	-	-	3/9/9/52	-
3	6OU	L	401	-	-	8/14/14/52	-
3	6OU	F	802	-	-	5/14/14/52	-
3	6OU	I	807	-	-	3/9/9/52	-
3	6OU	J	806	-	-	6/9/9/52	-
3	6OU	E	804	-	-	4/10/10/52	-
3	6OU	H	401	-	-	22/42/42/52	-
3	6OU	A	804	-	-	10/21/21/52	-
3	6OU	J	802	-	-	9/14/14/52	-
3	6OU	A	801	-	-	22/35/35/52	-
3	6OU	L	402	-	-	3/9/9/52	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	6OU	A	806	-	-	5/9/9/52	-
3	6OU	J	804	-	-	5/11/11/52	-
3	6OU	A	807	-	-	10/26/26/52	-
3	6OU	E	801	-	-	2/6/6/52	-
3	6OU	I	803	-	-	16/35/35/52	-
3	6OU	K	401	-	-	13/41/41/52	-
3	6OU	I	801	-	-	10/14/14/52	-
3	6OU	I	804	-	-	3/8/8/52	-
3	6OU	A	805	-	-	2/5/5/52	-
3	6OU	I	809	-	-	5/10/10/52	-
3	6OU	B	802	-	-	6/9/9/52	-
3	6OU	J	805	-	-	4/7/7/52	-
3	6OU	I	805	-	-	3/8/8/52	-
3	6OU	E	802	-	-	3/5/5/52	-
3	6OU	A	802	-	-	7/14/14/52	-

The worst 5 of 53 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	J	802	6OU	C41-C40	4.24	1.56	1.31
3	L	401	6OU	C41-C40	4.23	1.56	1.31
3	I	801	6OU	C41-C40	4.23	1.56	1.31
3	J	801	6OU	C41-C40	4.23	1.56	1.31
3	A	802	6OU	C41-C40	4.22	1.56	1.31

The worst 5 of 21 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	801	6OU	O30-C31-C33	4.05	120.23	111.50
3	A	807	6OU	O30-C31-C33	3.94	120.00	111.50
3	H	401	6OU	O30-C31-C33	3.94	119.98	111.50
3	A	801	6OU	O30-C31-C33	3.93	119.97	111.50
3	I	803	6OU	O30-C31-C33	3.91	119.94	111.50

There are no chirality outliers.


5 of 269 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	I	803	6OU	O18-C19-C20-O30
3	A	801	6OU	C27-O26-P23-O22
3	A	801	6OU	C27-O26-P23-O24
3	A	801	6OU	C27-O26-P23-O25
3	A	801	6OU	O26-C27-C28-N29


There are no ring outliers.


No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.
































5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-25116. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map




The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 177



Y Index: 193




Z Index: 201


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal surface views [i](#)


6.4.1 Primary map



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.4. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.5 Mask visualisation

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [\(i\)](#)



The volume at the recommended contour level is 291 nm³; this corresponds to an approximate mass of 262 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [i](#)



*Reported resolution corresponds to spatial frequency of 0.345\AA^{-1}


8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-25116 and PDB model 7SGR. Per-residue inclusion information can be found in section 3 on page 8.

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.4 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.4).



























9.4 Atom inclusion [i](#)



At the recommended contour level, 87% of all backbone atoms, 85% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.4) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.8476	 0.5180
A	 0.8547	 0.5260
B	 0.8276	 0.4990
C	 0.6364	 0.4040
D	 0.6973	 0.4320
E	 0.8935	 0.5530
F	 0.8788	 0.5330
G	 0.5799	 0.3200
H	 0.8414	 0.5200
I	 0.8781	 0.5420
J	 0.8740	 0.5280
K	 0.7190	 0.4620
L	 0.8468	 0.5010

