

wwPDB X-ray Structure Validation Summary Report (i)

Oct 15, 2023 – 05:30 AM EDT

PDB ID : 7SGX

Title : Crystal Structure of Turtle Cadherin-23 EC1-2

Authors: Nisler, C.R.; Sotomayor, M.

Deposited on : 2021-10-07

Resolution : 3.13 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.36

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 3.13 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	Similar resolution $(\# \text{Entries, resolution range}(\text{\AA}))$
R_{free}	130704	1626 (3.18-3.10)
Clashscore	141614	1735 (3.18-3.10)
Ramachandran outliers	138981	1677 (3.18-3.10)
Sidechain outliers	138945	1677 (3.18-3.10)
RSRZ outliers	127900	1588 (3.18-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	218	82%	12%	6%
1	В	218	6% 85%	9%	6%
1	С	218	5% 89%	6%	6%
1	D	218	83%	12%	5%



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 6517 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Cadherin 23.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
1	Λ	206	Total C N O	0	2	0
1	A	200	1602 1015 266 321	U	<i>L</i>	0
1	В	206	Total C N O	0	2	0
1	Б	200	1602 1015 266 321	U	2	0
1	С	206	Total C N O	0	2	0
1		200	1602 1015 266 321	U	2	0
1	D	208	Total C N O S	0	2	0
1	D	200	1618 1024 268 325 1	U	<i>L</i>	U

• Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	7	Total Ca 7 7	0	0
2	В	8	Total Ca 8 8	0	0
2	С	9	Total Ca 9 9	0	0
2	D	7	Total Ca 7 7	0	0

• Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Cl 1 1	0	0
3	D	1	Total Cl 1 1	0	0

• Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 6 3 3	0	0
4	D	1	Total C O 6 3 3	0	0


• Molecule 5 is water.

Mol	Chain	Residues	${f Atoms}$	ZeroOcc	AltConf
5	A	9	Total O 9 9	0	0
5	В	13	Total O 13 13	0	0
5	С	12	Total O 12 12	0	0
5	D	14	Total O 14 14	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants	94.98Å 94.98Å 358.36Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.06 - 3.13	Depositor
resolution (A)	49.01 - 3.13	EDS
% Data completeness	99.4 (49.06-3.13)	Depositor
(in resolution range)	99.5 (49.01-3.13)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.82 (at 3.12Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
P. P.	0.253 , 0.310	Depositor
R, R_{free}	0.251 , 0.305	DCC
R_{free} test set	1525 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	77.1	Xtriage
Anisotropy	0.669	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.31, 52.5	EDS
L-test for twinning ²	$ < L >=0.46, < L^2>=0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	6517	wwPDB-VP
Average B, all atoms (Å ²)	107.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 40.31 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.7900e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, GOL, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.87	1/1636~(0.1%)	0.78	0/2235
1	В	0.87	0/1636	0.76	0/2235
1	С	0.85	0/1636	0.77	0/2235
1	D	0.88	1/1652~(0.1%)	0.77	0/2256
All	All	0.87	$2/6560 \ (0.0\%)$	0.77	0/8961

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\operatorname{\AA})$	$\operatorname{Ideal}(ext{\AA})$
1	A	56	GLU	CD-OE2	6.09	1.32	1.25
1	D	145	SER	CA-CB	-5.30	1.45	1.52

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1602	0	1548	17	0
1	В	1602	0	1548	13	0
1	С	1602	0	1548	7	0
1	D	1618	0	1561	15	0
2	A	7	0	0	0	0

Continued on next page...



qe

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	В	8	0	0	0	0
2	С	9	0	0	0	0
2	D	7	0	0	0	0
3	A	1	0	0	0	0
3	D	1	0	0	0	0
4	A	6	0	8	0	0
4	D	6	0	8	0	0
5	A	9	0	0	0	0
5	В	13	0	0	2	0
5	С	12	0	0	0	0
5	D	14	0	0	0	0
All	All	6517	0	6221	46	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 46 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$\begin{array}{c} \text{Clash} \\ \text{overlap } (\text{\AA}) \end{array}$
1:B:130:ASN:ND2	5:B:401:HOH:O	2.32	0.62
1:C:107:PHE:CE1	1:C:131:ALA:HB2	2.36	0.61
1:D:107:PHE:CD2	1:D:195:ALA:HB2	2.37	0.60
1:B:110:GLN:N	1:B:110:GLN:OE1	2.34	0.59
1:B:63[A]:TRP:HH2	1:D:61:VAL:HG11	1.67	0.58

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	204/218 (94%)	187 (92%)	16 (8%)	1 (0%)	29 63	

Continued on next page...



Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	В	204/218 (94%)	184 (90%)	20 (10%)	0	100	100
1	С	204/218 (94%)	184 (90%)	20 (10%)	0	100	100
1	D	206/218 (94%)	189 (92%)	17 (8%)	0	100	100
All	All	818/872 (94%)	744 (91%)	73 (9%)	1 (0%)	51	82

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	135	ASP

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	180/192 (94%)	177 (98%)	3 (2%)	60 82
1	В	180/192 (94%)	176 (98%)	4 (2%)	52 77
1	С	180/192 (94%)	177 (98%)	3 (2%)	60 82
1	D	182/192 (95%)	177 (97%)	5 (3%)	44 72
All	All	722/768~(94%)	707 (98%)	15 (2%)	53 78

5 of 15 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	С	48	GLU
1	D	163	SER
1	С	133	ASP
1	D	202	THR
1	D	28	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:



Mol	Chain	Res	Type
1	С	181	ASN
1	D	86	GLN
1	D	196	ASN
1	В	185	GLN
1	A	103	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 35 ligands modelled in this entry, 33 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Ros Lin		Res Link		Bond lengths			В	ond ang	gles
IVIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2		
4	GOL	D	309	-	5,5,5	0.22	0	5,5,5	0.22	0		
4	GOL	A	309	-	5,5,5	0.29	0	5,5,5	0.18	0		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	D	309	-	-	3/4/4/4	-
4	GOL	A	309	-	-	3/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 6 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	D	309	GOL	C1-C2-C3-O3
4	D	309	GOL	O2-C2-C3-O3
4	A	309	GOL	O1-C1-C2-C3
4	A	309	GOL	C1-C2-C3-O3
4	A	309	GOL	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	#RSRZ>2		$OWAB(A^2)$	Q < 0.9
1	A	206/218 (94%)	0.22	7 (3%) 45	24	54, 94, 137, 157	0
1	В	206/218 (94%)	0.38	12 (5%) 23	10	74, 111, 166, 195	0
1	С	206/218 (94%)	0.33	11 (5%) 26	12	78, 110, 163, 193	0
1	D	208/218 (95%)	0.18	2 (0%) 82	70	59, 96, 142, 170	0
All	All	826/872 (94%)	0.28	32 (3%) 39	20	54, 102, 156, 195	0

The worst 5 of 32 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	С	5	TYR	4.5
1	В	6	PHE	3.3
1	D	79	GLU	3.0
1	В	49	ALA	2.9
1	В	68	LEU	2.9

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}(\mathring{\mathbf{A}}^2)$	Q<0.9
2	CA	A	306	1/1	0.89	0.17	94,94,94,94	0
2	CA	С	306	1/1	0.91	0.10	94,94,94,94	0
2	CA	В	308	1/1	0.93	0.30	94,94,94,94	0
2	CA	В	304	1/1	0.93	0.09	169,169,169,169	0
3	CL	D	308	1/1	0.93	0.10	69,69,69,69	0
2	CA	D	303	1/1	0.94	0.11	99,99,99,99	0
2	CA	В	307	1/1	0.94	0.24	87,87,87,87	0
2	CA	С	305	1/1	0.95	0.12	93,93,93,93	0
2	CA	В	301	1/1	0.95	0.09	106,106,106,106	0
2	CA	С	307	1/1	0.96	0.34	104,104,104,104	0
2	CA	С	308	1/1	0.96	0.28	85,85,85,85	0
2	CA	D	301	1/1	0.96	0.16	88,88,88,88	0
2	CA	A	305	1/1	0.96	0.18	82,82,82,82	0
2	CA	D	304	1/1	0.96	0.07	128,128,128,128	0
2	CA	D	305	1/1	0.96	0.18	95,95,95,95	0
2	CA	С	304	1/1	0.96	0.04	145,145,145,145	0
4	GOL	D	309	6/6	0.96	0.23	56,60,65,69	0
2	CA	D	302	1/1	0.97	0.16	$92,\!92,\!92,\!92$	0
2	CA	A	303	1/1	0.97	0.13	86,86,86,86	0
2	CA	A	304	1/1	0.97	0.09	106,106,106,106	0
2	CA	В	305	1/1	0.97	0.12	89,89,89,89	0
3	CL	A	308	1/1	0.97	0.26	52,52,52,52	0
2	CA	С	309	1/1	0.97	0.24	81,81,81,81	0
4	GOL	A	309	6/6	0.97	0.22	54,57,59,59	0
2	CA	В	306	1/1	0.97	0.10	97,97,97,97	0
2	CA	С	303	1/1	0.98	0.12	114,114,114,114	0
2	CA	В	303	1/1	0.98	0.14	96,96,96,96	0
2	CA	С	301	1/1	0.98	0.10	97,97,97,97	0
2	CA	D	307	1/1	0.98	0.16	85,85,85,85	0
2	CA	D	306	1/1	0.99	0.18	81,81,81,81	0
2	CA	A	307	1/1	0.99	0.22	75,75,75,75	0
2	CA	A	301	1/1	0.99	0.10	77,77,77,77	0
2	CA	С	302	1/1	0.99	0.12	93,93,93,93	0
2	CA	В	302	1/1	0.99	0.13	85,85,85,85	0
2	CA	A	302	1/1	0.99	0.15	82,82,82,82	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.














Electron density around CA B 308: $2 \mathrm{mF}_o\text{-DF}_c \text{ (at 0.7 rmsd) in gray} \\ \mathrm{mF}_o\text{-DF}_c \text{ (at 3 rmsd) in purple (negative)}$


and green (positive)




































Electron density around CA C 307:

 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









Electron density around CA C 308: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)








































































































6.5 Other polymers (i)

There are no such residues in this entry.

