

# wwPDB X-ray Structure Validation Summary Report (i)

#### Nov 7, 2023 – 07:07 PM EST

PDB ID : 1SM1

Title : COMPLEX OF THE LARGE RIBOSOMAL SUBUNIT FROM DEINOCOC-

CUS RADIODURANS WITH QUINUPRISTIN AND DALFOPRISTIN

Authors: Harms, J.M.; Schluenzen, F.; Fucini, P.; Bartels, H.; Yonath, A.

Deposited on : 2004-03-08

Resolution : 3.42 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : NOT EXECUTED

EDS : NOT EXECUTED

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$ 

The reported resolution of this entry is 3.42 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution		
Metric	$(\# \mathrm{Entries})$	$(\#  ext{Entries},  ext{resolution range}( ext{Å}))$		
Clashscore	141614	1572 (3.50-3.34)		
Ramachandran outliers	138981	1534 (3.50-3.34)		
Sidechain outliers	138945	1535 (3.50-3.34)		
RNA backbone	3102	1012 (3.88-2.96)		

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length		Quality of chain	
1	0	2880	29%	53%	13% •••
2	1	82		65%	35%
3	2	47		98%	
4	3	66		95%	5%
5	4	37		95%	5%
6	5	8	25%	50%	12% 12%

Continued on next page...



Continued from previous page...

Mol	Chain	Length	Quality of chain	
7	9	124	36% 54%	5% 5%
8	A	275	98%	
9	В	211	97%	
10	С	205	96%	
11	D	180	99%	
12	Е	212	83%	17%
13	F	146	36% 64%	
14	G	144	99%	
15	Н	174	82%	18%
16	I	134	99%	
17	J	156	90%	10%
18	K	142	87%	13%
19	L	116	97%	1370
20	M	114	97%	···
21	N	166		•
			75%	25%
22	0	118	97%	••
23	Р	100	100%	
24	Q	134	97%	•
25	R	95	98%	·
26	S	115	98%	•
27	Т	253	88%	12%
28	U	91	95%	5%
29	W	67	97%	•
30	X	55	100%	
31	Y	73	100%	

Continued on next page...



Continued from previous page...

Mol	Chain	Length	Quality of chain	
32	Z	60	95%	·

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
33	DOL	0	2882	X	-	-	-
6	DBB	5	3	-	-	X	-



# 2 Entry composition (i)

There are 33 unique types of molecules in this entry. The entry contains 65418 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a RNA chain called 23S RIBOSOMAL RNA.

Mol	Chain	Residues		-	Atoms	ZeroOcc	AltConf	Trace		
1	0	2766	Total 59359	C 26479	N 10949	O 19166	P 2765	0	0	0

• Molecule 2 is a protein called 50S RIBOSOMAL PROTEIN L33.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
2	1	53	Total C 53 53	0	0	53

• Molecule 3 is a protein called 50S RIBOSOMAL PROTEIN L34.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
3	2	46	Total C 46 46	0	0	46

• Molecule 4 is a protein called 50S RIBOSOMAL PROTEIN L35.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
4	3	63	Total C 63 63	0	0	63

• Molecule 5 is a protein called 50S RIBOSOMAL PROTEIN L36.

$\mathbf{Mol}$	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
5	4	35	Total C 35 35	0	0	35

• Molecule 6 is a protein called QUINUPRISTIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	5	Q	Total	С	N	О	S	0	0	0
0	9	9 0	73	53	9	10	1	U	0	



• Molecule 7 is a RNA chain called 5S RIBOSOMAL RNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
7	9	118	Total 2516	C 1124	N 464	O 811	P 117	0	0	0

• Molecule 8 is a protein called 50S RIBOSOMAL PROTEIN L2.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
8	A	270	Total C 270 270	0	0	270

• Molecule 9 is a protein called 50S RIBOSOMAL PROTEIN L3.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
9	В	205	Total C 205 205	0	0	205

• Molecule 10 is a protein called 50S RIBOSOMAL PROTEIN L4.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
10	С	197	Total C 197 197	0	0	197

• Molecule 11 is a protein called 50S RIBOSOMAL PROTEIN L5.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
11	D	178	Total C 178 178	0	0	178

• Molecule 12 is a protein called 50S RIBOSOMAL PROTEIN L6.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
12	E	177	Total C 177 177	0	0	177

• Molecule 13 is a protein called 50S RIBOSOMAL PROTEIN L9.

$\mathbf{N}$	/Iol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
	13	F	52	Total C 52 52	0	0	52

• Molecule 14 is a protein called 50S RIBOSOMAL PROTEIN L11.



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
14	G	143	Total C 143 143	0	0	143

 $\bullet$  Molecule 15 is a protein called 50S RIBOSOMAL PROTEIN L13.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	Trace
15	Н	143	Total 143	C 143	0	0	143

• Molecule 16 is a protein called 50S RIBOSOMAL PROTEIN L14.

Mo	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
16	I	132	Total ( 132 13	0	0	132

• Molecule 17 is a protein called 50S RIBOSOMAL PROTEIN L15.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	Trace
17	J	141	Total 141	C 141	0	0	141

• Molecule 18 is a protein called 50S RIBOSOMAL PROTEIN L16.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	Trace
18	К	124	Total 124	C 124	0	0	124

• Molecule 19 is a protein called 50S RIBOSOMAL PROTEIN L17.

N	Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	Trace
	19	L	114	Total 114	C 114	0	0	114

• Molecule 20 is a protein called 50S RIBOSOMAL PROTEIN L18.

Mol	Chain	Residues	Ato	ms	ZeroOcc	AltConf	Trace
20	M	111	Total 111	C 111	8	0	111

• Molecule 21 is a protein called 50S RIBOSOMAL PROTEIN L19.



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
21	N	125	Total C 125 125	0	0	125

• Molecule 22 is a protein called 50S RIBOSOMAL PROTEIN L20.

]	Mol	Chain	Residues	Aton	ns	ZeroOcc	AltConf	Trace
	22	О	117	Total 117	C 117	16	0	117

• Molecule 23 is a protein called 50S RIBOSOMAL PROTEIN L21.

$\mathbf{Mol}$	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
23	Р	100	Total C 100 100	0	0	100

• Molecule 24 is a protein called 50S RIBOSOMAL PROTEIN L22.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
24	Q	130	Total C 130 130	0	0	130

• Molecule 25 is a protein called 50S RIBOSOMAL PROTEIN L23.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
25	R	93	Total C 93 93	0	0	93

• Molecule 26 is a protein called 50S RIBOSOMAL PROTEIN L24.

Mol	Chain	Residues	Ator	ns	ZeroOcc	AltConf	Trace
26	S	113	Total 113	C 113	0	0	113

• Molecule 27 is a protein called GENERAL STRESS PROTEIN CTC.

Mo	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
27	Т	223	Total C 223 223	43	0	223

• Molecule 28 is a protein called 50S RIBOSOMAL PROTEIN L27.



$\mathbf{Mol}$	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
28	U	86	Total C 86 86	0	0	86

• Molecule 29 is a protein called 50S RIBOSOMAL PROTEIN L29.

Mo	l Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
29	W	65	Total C 65 65	0	0	65

• Molecule 30 is a protein called 50S RIBOSOMAL PROTEIN L30.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
30	X	55	Total C 55 55	4	0	55

• Molecule 31 is a protein called 50S RIBOSOMAL PROTEIN L31.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
31	Y	73	Total C 73 73	0	0	73

• Molecule 32 is a protein called 50S RIBOSOMAL PROTEIN L32.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
32	Z	58	Total C 58 58	0	0	58

• Molecule 33 is 5-(2-DIETHYLAMINO-ETHANESULFONYL)-21-HYDROXY-10-ISOPRO PYL-11,19-DIMETHYL-9,26-DIOXA-3,15,28-TRIAZA-TRICYCLO[23.2.1.00,255]OCTAC OSA-1(27),12,17,19,25(28)-PENTAENE-2,8,14,23-TETRAONE (three-letter code: DOL) (formula:  $C_{34}H_{50}N_4O_9S$ ).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf		
33	0	1	Total	С	N	О	S	0	0
33	U	1	48	34	4	9	1		




# 3 Residue-property plots (i)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


• Molecule 1: 23S RIBOSOMAL RNA






























Chain P:	100%	
There are no	outlier residues recorded for this chain.	
• Molecule 2	4: 50S RIBOSOMAL PROTEIN L22	
Chain Q:	97%	
MET THR ALA PRO ES KI34		
• Molecule 2	5: 50S RIBOSOMAL PROTEIN L23	
Chain R:	98%	
NET S2 Q94 ALA		
• Molecule 2	6: 50S RIBOSOMAL PROTEIN L24	
Chain S:	98%	
MET P2 1114 I ASP		
• Molecule 2	7: GENERAL STRESS PROTEIN CTC	
Chain T:	88%	12%
MET ALA HIS HIS GLY GLY THR ALA LYS	SER GLN GLY THR THR THR ASP ASP ASP ASP ASP ASP GLU GLU GLU GLU GLU GLU GLU GLU	
• Molecule 2	8: 50S RIBOSOMAL PROTEIN L27	
Chain U:	95%	5%
MET A2 E87 VAL ALA ALA ASP		
• Molecule 2	9: 50S RIBOSOMAL PROTEIN L29	
Chain W:	97%	
MET K2 Q66 GLN		
• Molecule 3	0: 50S RIBOSOMAL PROTEIN L30	
Chain X:	100%	
There are no	outlier residues recorded for this chain.	



• Molecule 31: 50S RIBOSOMAL PROTEIN L31

Chain Y: 100%

There are no outlier residues recorded for this chain.

• Molecule 32: 50S RIBOSOMAL PROTEIN L32

Chain Z: 95% ...





# 4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source	
Space group	I 2 2 2	Depositor	
Cell constants	168.50Å 406.00Å 693.00Å	Depositor	
a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.00° 90.00° 90.00°	Depositor	
Resolution (Å)	15.00 - 3.42	Depositor	
% Data completeness	(Not available) (15.00-3.42)	Depositor	
(in resolution range)	(100 available) (15.00-5.42)		
$R_{merge}$	(Not available)	Depositor	
$R_{sym}$	0.16	Depositor	
Refinement program	CNS	Depositor	
$R, R_{free}$	0.278 , 0.348	Depositor	
Estimated twinning fraction	No twinning to report.	Xtriage	
Total number of atoms	65418	wwPDB-VP	
Average B, all atoms (Å <sup>2</sup> )	80.0	wwPDB-VP	



## 5 Model quality (i)

#### 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: 004, MHU, MHW, DOL, MHV, MHT, DBB

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain		Bo	ond lengths	Bond angles		
MIOI	Chain	RMSZ	# Z  > 5	RMSZ	# Z >5	
1	0	0.70	$16/66467 \ (0.0\%)$	0.74	12/103673 (0.0%)	
6	5	0.85	0/13	0.67	0/15	
7	9	0.50	0/2813	0.65	0/4384	
All	All	0.70	$16/69293 \ (0.0\%)$	0.73	$12/108072 \ (0.0\%)$	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	0	0	146
6	5	1	1
7	9	0	1
All	All	1	148

The worst 5 of 16 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	Observed(A)	$\operatorname{Ideal}( ext{\AA})$
1	0	1962	С	N1-C2	-7.46	1.32	1.40
1	0	2255	G	C5-C6	-7.28	1.35	1.42
1	0	2789	U	N1-C2	6.94	1.44	1.38
1	0	868	U	N1-C2	6.93	1.44	1.38
1	0	806	A	C5-C6	6.88	1.47	1.41

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$\mathbf{Observed}(^{o})$	$\operatorname{Ideal}({}^{o})$
1	0	994	A	N9-C1'-C2'	-6.36	105.00	112.00
1	0	800	U	OP2-P-O3'	6.27	119.00	105.20

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
1	0	2056	С	N1-C1'-C2'	-6.04	105.36	112.00
1	0	1686	A	C5'-C4'-O4'	5.86	116.13	109.10
1	0	1938	U	C2'-C3'-O3'	5.75	122.90	113.70

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
6	5	8	MHT	C3

5 of 148 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	0	126	С	Sidechain
1	0	174	A	Sidechain
1	0	211	U	Sidechain
1	0	33	С	Sidechain
1	0	8	A	Sidechain

#### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	0	59359	0	29917	2138	0
2	1	53	0	0	0	0
3	2	46	0	0	0	0
4	3	63	0	0	0	0
5	4	35	0	0	0	0
6	5	73	0	64	6	0
7	9	2516	0	1286	66	0
8	A	270	0	0	1	0
9	В	205	0	0	1	0
10	С	197	0	0	0	0
11	D	178	0	0	0	0
12	Е	177	0	0	0	0
13	F	52	0	0	0	0
14	G	143	0	0	0	0
15	Н	143	0	0	0	0

Continued on next page...



1SM1

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
16	I	132	0	0	0	0
17	J	141	0	0	0	0
18	K	124	0	0	0	0
19	L	114	0	0	1	0
20	M	111	0	0	0	0
21	N	125	0	0	0	0
22	О	117	0	0	2	0
23	Р	100	0	0	0	0
24	Q	130	0	0	0	0
25	R	93	0	0	0	0
26	S	113	0	0	0	0
27	Τ	223	0	0	0	0
28	U	86	0	0	0	0
29	W	65	0	0	0	0
30	X	55	0	0	0	0
31	Y	73	0	0	0	0
32	Z	58	0	0	2	0
33	0	48	0	47	16	0
All	All	65418	0	31314	2213	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 23.

The worst 5 of 2213 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$egin{aligned} &  ext{Interatomic} \ &  ext{distance} \ &  ext{(Å)} \end{aligned}$	$egin{aligned}  ext{Clash} \  ext{overlap} & ( ext{Å}) \end{aligned}$
1:0:1463:A:H1'	1:0:1543:G:H22	1.05	1.14
1:0:128:C:H2'	1:0:129:A:H5"	1.19	1.10
1:0:1656:U:H2'	1:0:1657:A:H5"	1.34	1.10
1:0:940:G:H3'	1:0:941:U:H5"	1.23	1.09
1:0:2607:C:H3'	1:0:2608:A:H5'	1.10	1.08

There are no symmetry-related clashes.

### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	alysed Favoured Allowed		Outliers		
6	5	2/8 (25%)	1 (50%)	0	1 (50%)	0 0	

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
6	5	2	THR

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percei	ntiles
6	5	2/2 (100%)	2 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

#### 5.3.3 RNA (i)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	0	2757/2880 (95%)	433 (15%)	19 (0%)
7	9	117/124 (94%)	12 (10%)	0
All	All	2874/3004 (95%)	445 (15%)	19 (0%)

5 of 445 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	0	45	С
1	0	48	A
1	0	49	U
1	0	50	G
1	0	59	G



5 of 19 RNA pucker outliers are listed below:

Mol	Chain	Res	Type
1	0	2015	G
1	0	2377	U
1	0	2404	A
1	0	2261	G
1	0	1354	A

#### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

5 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

N / - 1	Mal True Chain D		Das	Bond lengths			Bond angles			
Mol	Type	Chain	Res	Link	Counts	RMSZ	$\mid \# Z  > 2$	Counts	RMSZ	# Z  > 2
6	004	5	7	6	9,10,11	1.69	2 (22%)	9,12,14	1.27	1 (11%)
6	MHW	5	1	6	9,9,10	0.76	0	10,11,13	1.56	1 (10%)
6	MHV	5	6	6	7,9,10	0.67	0	7,11,13	1.68	2 (28%)
6	MHU	5	5	6	14,15,16	1.14	1 (7%)	18,19,21	1.11	1 (5%)
6	DBB	5	3	6	4,5,6	0.58	0	1,5,7	0.06	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	004	5	7	6	-	2/4/6/8	0/1/1/1
6	MHW	5	1	6	-	2/2/2/4	0/1/1/1
6	MHV	5	6	6	-	0/1/12/14	0/1/1/1
6	MHU	5	5	6	-	2/9/12/14	0/1/1/1
6	DBB	5	3	6	-	1/3/4/6	-

All (3) bond length outliers are listed below:



Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	Observed(A)	$Ideal(\AA)$
6	5	7	004	CB-CA	3.12	1.55	1.52
6	5	7	004	CG2-CB	-2.70	1.34	1.39
6	5	5	MHU	CZ1-NZ	-2.59	1.39	1.45

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
6	5	1	MHW	O-C-CA	-4.20	120.24	124.22
6	5	6	MHV	CE-CD2-CG	3.22	117.29	111.89
6	5	5	MHU	O-C-CA	-2.83	117.37	124.78
6	5	7	004	CG2-CB-CA	2.31	124.38	120.65
6	5	6	MHV	CB-CA-N	-2.02	108.33	112.50

There are no chirality outliers.

5 of 7 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	5	1	MHW	O-C-CA-N
6	5	1	MHW	O-C-CA-CB
6	5	3	DBB	O-C-CA-CB
6	5	5	MHU	N-CA-CB-CG
6	5	5	MHU	C-CA-CB-CG

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	5	1	MHW	1	0
6	5	3	DBB	5	0

### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry (i)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The



Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

	Mol	Type	Chain Re	Res	Link	В	ond leng	$\overline{ ext{gths}}$	В	ond ang	gles
	Moi   Type	туре		nes	LIIIK	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
Ī	33	DOL	0	2882	-	43,50,50	4.58	11 (25%)	51,70,70	3.94	18 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
33	DOL	0	2882	-	2/2/14/20	20/58/77/77	0/2/3/3

The worst 5 of 11 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(A)	$\operatorname{Ideal}(\text{\AA})$
33	0	2882	DOL	O40-S39	18.61	1.77	1.44
33	0	2882	DOL	O41-S39	18.41	1.76	1.44
33	0	2882	DOL	C28-C29	-8.71	1.11	1.32
33	0	2882	DOL	C1-C37	4.83	1.62	1.52
33	0	2882	DOL	C8-C6	-4.51	1.42	1.50

The worst 5 of 18 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
33	0	2882	DOL	C4-N5-C1	-14.71	94.37	112.45
33	0	2882	DOL	O18-C17-C16	13.82	145.78	109.73
33	0	2882	DOL	C28-C26-N25	-9.38	97.30	114.97
33	0	2882	DOL	O40-S39-O41	-7.23	109.95	118.19
33	0	2882	DOL	O27-C26-C28	6.58	138.02	123.03

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
33	0	2882	DOL	C17
33	0	2882	DOL	C2

5 of 20 torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
33	0	2882	DOL	O7-C6-N5-C1
33	0	2882	DOL	C8-C6-N5-C1
33	0	2882	DOL	C1-C2-S39-O41
33	0	2882	DOL	C1-C2-S39-O40
33	0	2882	DOL	C1-C2-S39-C42


There are no ring outliers.

1 monomer is involved in 16 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
33	0	2882	DOL	16	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





### 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

#### 6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

#### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

#### 6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

### 6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

#### 6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

