

wwPDB EM Validation Summary Report (i)

Nov 26, 2022 – 12:22 PM EST

PDB ID	:	7STM
EMDB ID	:	EMD-25433
Title	:	Chitin Synthase 2 from Candida albicans bound to UDP-GlcNAc
Authors	:	Ren, Z.; Chhetri, A.; Lee, S.; Yokoyama, K.
Deposited on	:	2021-11-14
Resolution	:	3.02 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/EMValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


EMDB validation analysis	:	0.0.1.dev 43
Mogul	:	1.8.5 (274361), CSD as541be (2020)
MolProbity	:	4.02b-467
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ	:	1.9.9
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.31.2

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $ELECTRON\ MICROSCOPY$

The reported resolution of this entry is 3.02 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \textbf{Whole archive} \\ \textbf{(\#Entries)} \end{array}$	${f EM} {f structures} \ (\#{f Entries})$		
		(#Entries)		
Clashscore	158937	4297		
Ramachandran outliers	154571	4023		
Sidechain outliers	154315	3826		

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion < 40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain					
1	А	1039	74%	•	23%			
1	В	1039	74%	•	23%			



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 25450 atoms, of which 12510 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms					AltConf	Trace	
1	Δ	801	Total	С	Η	Ν	Ο	S	0	0
	Л		12321	4100	6050	1028	1115	28	0	0
1	Р	801	Total	С	Η	Ν	Ο	S	0	0
	D	801	12321	4100	6050	1028	1115	28	0	0

• Molecule 1 is a protein called Chitin synthase.

Chain	Residue	Modelled	Actual	Comment	Reference
А	1010	ALA	-	expression tag	UNP A0A1D8PTV3
А	1011	ASN	-	expression tag	UNP A0A1D8PTV3
А	1012	SER	-	expression tag	UNP A0A1D8PTV3
А	1013	LEU	-	expression tag	UNP A0A1D8PTV3
А	1014	GLU	-	expression tag	UNP A0A1D8PTV3
А	1015	VAL	-	expression tag	UNP A0A1D8PTV3
А	1016	LEU	-	expression tag	UNP A0A1D8PTV3
А	1017	PHE	-	expression tag	UNP A0A1D8PTV3
А	1018	GLN	-	expression tag	UNP A0A1D8PTV3
А	1019	GLY	-	expression tag	UNP A0A1D8PTV3
А	1020	PRO	-	expression tag	UNP A0A1D8PTV3
A	1021	ASP	-	expression tag	UNP A0A1D8PTV3
А	1022	TYR	-	expression tag	UNP A0A1D8PTV3
А	1023	LYS	-	expression tag	UNP A0A1D8PTV3
A	1024	ASP	-	expression tag	UNP A0A1D8PTV3
А	1025	ASP	-	expression tag	UNP A0A1D8PTV3
А	1026	ASP	-	expression tag	UNP A0A1D8PTV3
А	1027	ASP	-	expression tag	UNP A0A1D8PTV3
А	1028	LYS	-	expression tag	UNP A0A1D8PTV3
А	1029	ALA	-	expression tag	UNP A0A1D8PTV3
А	1030	HIS	-	expression tag	UNP A0A1D8PTV3
А	1031	HIS	-	expression tag	UNP A0A1D8PTV3
А	1032	HIS	-	expression tag	UNP A0A1D8PTV3
А	1033	HIS	-	expression tag	UNP A0A1D8PTV3
А	1034	HIS	-	expression tag	UNP A0A1D8PTV3
А	1035	HIS	_	expression tag	UNP A0A1D8PTV3

There are 60 discrepancies between the modelled and reference sequences:




Continued on next page...

	ed from pre	DC			
Chain	Residue	Modelled	Actual	Comment	Reference
A	1036	HIS	-	expression tag	UNP A0A1D8PTV3
A	1037	HIS	-	expression tag	UNP A0A1D8PTV3
A	1038	HIS	-	expression tag	UNP A0A1D8PTV3
A	1039	HIS	-	expression tag	UNP A0A1D8PTV3
В	1010	ALA	-	expression tag	UNP A0A1D8PTV3
В	1011	ASN	-	expression tag	UNP A0A1D8PTV3
В	1012	SER	-	expression tag	UNP A0A1D8PTV3
В	1013	LEU	-	expression tag	UNP A0A1D8PTV3
В	1014	GLU	-	expression tag	UNP A0A1D8PTV3
В	1015	VAL	-	expression tag	UNP A0A1D8PTV3
В	1016	LEU	-	expression tag	UNP A0A1D8PTV3
В	1017	PHE	-	expression tag	UNP A0A1D8PTV3
В	1018	GLN	-	expression tag	UNP A0A1D8PTV3
В	1019	GLY	-	expression tag	UNP A0A1D8PTV3
В	1020	PRO	-	expression tag	UNP A0A1D8PTV3
В	1021	ASP	-	expression tag	UNP A0A1D8PTV3
В	1022	TYR	-	expression tag	UNP A0A1D8PTV3
В	1023	LYS	-	expression tag	UNP A0A1D8PTV3
В	1024	ASP	-	expression tag	UNP A0A1D8PTV3
В	1025	ASP	-	expression tag	UNP A0A1D8PTV3
В	1026	ASP	-	expression tag	UNP A0A1D8PTV3
В	1027	ASP	-	expression tag	UNP A0A1D8PTV3
В	1028	LYS	-	expression tag	UNP A0A1D8PTV3
В	1029	ALA	-	expression tag	UNP A0A1D8PTV3
В	1030	HIS	-	expression tag	UNP A0A1D8PTV3
В	1031	HIS	-	expression tag	UNP A0A1D8PTV3
В	1032	HIS	-	expression tag	UNP A0A1D8PTV3
В	1033	HIS	-	expression tag	UNP A0A1D8PTV3
В	1034	HIS	-	expression tag	UNP A0A1D8PTV3
В	1035	HIS	-	expression tag	UNP A0A1D8PTV3
В	1036	HIS	-	expression tag	UNP A0A1D8PTV3
В	1037	HIS	-	expression tag	UNP A0A1D8PTV3
В	1038	HIS	-	expression tag	UNP A0A1D8PTV3
В	1039	HIS	-	expression tag	UNP A0A1D8PTV3

Continued from previous page...

• Molecule 2 is 1,2-Distearoyl-sn-glycerophosphoethanolamine (three-letter code: 3PE) (formula: C₄₁H₈₂NO₈P) (labeled as "Ligand of Interest" by depositor).






Mol	Chain	Residues		A	tom	5			AltConf
2	А	1	Total	С	Η	Ν	0	Р	0
	A	1	364	120	205	3	32	4	0
2	А	1	Total	С	Η	Ν	Ο	Р	0
	A	1	364	120	205	3	32	4	0
2	А	1	Total	С	Η	Ν	Ο	Р	0
	A	1	364	120	205	3	32	4	0
2	А	1	Total	С	Η	Ν	Ο	Р	0
	Л	1	364	120	205	3	32	4	0
2	В	1	Total	С	Η	Ν	Ο	Р	0
	D	1	364	120	205	3	32	4	0
2	В	1	Total	С	Η	Ν	Ο	Р	0
	D	1	364	120	205	3	32	4	0
2	В	1	Total	С	Η	Ν	Ο	Р	0
	D	1	364	120	205	3	32	4	0
2	В	1	Total	С	Η	Ν	Ο	Р	0
	D	L	364	120	205	3	32	4	U

• Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	AltConf
3	А	1	Total Mg 1 1	0
3	В	1	Total Mg 1 1	0

• Molecule 4 is URIDINE-DIPHOSPHATE-N-ACETYLGLUCOSAMINE (three-letter code: UD1) (formula: $C_{17}H_{27}N_3O_{17}P_2$) (labeled as "Ligand of Interest" by depositor).






Mol	Chain	Residues	Atoms					AltConf
4	٨	1	Total	С	Ν	0	Р	0
4	A	1	39	17	3	17	2	0
4	D	1	Total	С	Ν	0	Р	0
4	D	1	39	17	3	17	2	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Chitin synthase







R3 66 1.420 1.420 1.420 1.420 241 242 243 243 244 245 245 245 245 247 251 251 251 251 251 251 251 251 251 251 251 251 254





4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C2	Depositor
Number of particles used	287321	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^{-}/\text{\AA}^2)$	60	Depositor
Minimum defocus (nm)	1000	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	Not provided	
Image detector	GATAN K3 $(6k \ge 4k)$	Depositor
Maximum map value	0.058	Depositor
Minimum map value	-0.025	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.002	Depositor
Recommended contour level	0.0078	Depositor
Map size (Å)	233.28001, 233.28001, 233.28001	wwPDB
Map dimensions	216, 216, 216	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.08, 1.08, 1.08	Depositor



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: UD1, 3PE, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles			
	Unam	RMSZ	# Z > 5	RMSZ	# Z > 5		
1	А	0.31	0/6433	0.46	0/8739		
1	В	0.31	0/6433	0.46	0/8739		
All	All	0.31	0/12866	0.46	0/17478		

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	6271	6050	6075	16	0
1	В	6271	6050	6075	16	0
2	А	159	205	211	0	0
2	В	159	205	211	0	0
3	А	1	0	0	0	0
3	В	1	0	0	0	0
4	А	39	0	25	2	0
4	В	39	0	25	2	0
All	All	12940	12510	12622	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:1206:UD1:O1A	4:A:1206:UD1:H5'	1.89	0.73
4:B:1206:UD1:H5'	4:B:1206:UD1:O1A	1.89	0.72
1:A:474:SER:HG	1:A:541:PHE:HZ	1.58	0.50
1:A:465:ASP:OD1	1:A:466:CYS:N	2.47	0.48
1:B:465:ASP:OD1	1:B:466:CYS:N	2.47	0.48

The worst 5 of 32 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	797/1039~(77%)	778~(98%)	19 (2%)	0	100	100
1	В	797/1039~(77%)	778~(98%)	19~(2%)	0	100	100
All	All	1594/2078~(77%)	1556~(98%)	38~(2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	А	644/923~(70%)	640 (99%)	4 (1%)	86 95	

Continued on next page...



Continued from previous page...

Mol	Chain	Analysed	nalysed Rotameric Outliers		Percentiles		
1	В	644/923~(70%)	640~(99%)	4 (1%)	86 95		
All	All	1288/1846~(70%)	1280 (99%)	8 (1%)	86 95		

5 of 8 residues with a non-rotameric side chain are listed below:

Mol	Chain	Res	Type
1	В	989	CYS
1	В	703	PHE
1	В	645	ARG
1	А	989	CYS
1	В	699	TYR

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. There are no such side chains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 12 ligands modelled in this entry, 2 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mol	Turne	Chain	Res	Link	B	ond leng	gths	B	ond ang	les
	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2
4	UD1	А	1206	3	38,41,41	2.72	17 (44%)	57,62,62	1.40	8 (14%)
2	3PE	А	1204	-	36,36,50	1.00	3 (8%)	39,41,55	1.00	2 (5%)
2	3PE	В	1201	-	40,40,50	0.96	4 (10%)	43,45,55	0.98	1 (2%)
2	3PE	А	1203	-	41,41,50	0.95	3 (7%)	44,46,55	1.08	2 (4%)
4	UD1	В	1206	3	38,41,41	2.72	17 (44%)	57,62,62	1.39	8 (14%)
2	3PE	В	1204	-	36,36,50	1.00	3 (8%)	39,41,55	1.00	2(5%)
2	3PE	А	1202	-	38,38,50	0.98	3 (7%)	41,43,55	1.04	2 (4%)
2	3PE	В	1202	-	38,38,50	0.97	3 (7%)	41,43,55	1.04	2 (4%)
2	3PE	А	1201	-	40,40,50	0.96	4 (10%)	43,45,55	0.98	1 (2%)
2	3PE	В	1203	-	41,41,50	0.95	3 (7%)	44,46,55	1.08	2 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	UD1	А	1206	3	-	13/26/63/63	0/3/3/3
2	3PE	А	1204	-	-	19/40/40/54	-
2	3PE	В	1201	-	-	13/44/44/54	-
2	3PE	А	1203	-	-	16/45/45/54	-
4	UD1	В	1206	3	-	13/26/63/63	0/3/3/3
2	3PE	В	1204	-	-	19/40/40/54	-
2	3PE	А	1202	-	-	8/42/42/54	-
2	3PE	В	1202	-	-	8/42/42/54	-
2	3PE	А	1201	-	-	13/44/44/54	-
2	3PE	В	1203	-	_	16/45/45/54	_

The worst 5 of 60 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(Å)
4	А	1206	UD1	O4-C4	7.83	1.39	1.24
4	В	1206	UD1	O4-C4	7.82	1.39	1.24
4	А	1206	UD1	O5'-C1'	5.48	1.55	1.41
4	В	1206	UD1	O5'-C1'	5.48	1.55	1.41
4	А	1206	UD1	O3B-C3B	4.69	1.54	1.43

The worst 5 of 30 bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
2	В	1203	3PE	O21-C21-C22	4.00	120.12	111.50
2	А	1203	3PE	O21-C21-C22	3.99	120.09	111.50
4	В	1206	UD1	O3A-PB-O1'	3.48	109.51	102.48
4	А	1206	UD1	O3A-PB-O1'	3.48	109.50	102.48
4	А	1206	UD1	C4-N3-C2	-3.45	122.03	126.58

There are no chirality outliers.

5 of 138 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	А	1201	3PE	C11-O13-P-O12
2	А	1201	3PE	C11-O13-P-O14
2	А	1203	3PE	C1-O11-P-O12
2	А	1203	3PE	C1-O11-P-O13
2	А	1203	3PE	C1-O11-P-O14


There are no ring outliers.

2 monomers are involved in 4 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	А	1206	UD1	2	0
4	В	1206	UD1	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.
























5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.




6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-25433. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections (i)


6.1.1 Primary map



The images above show the map projected in three orthogonal directions.

6.2 Central slices (i)

6.2.1 Primary map



X Index: 108

Y Index: 108




Z Index: 108

The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices (i)

6.3.1 Primary map



X Index: 103


Y Index: 77

Z Index: 110

The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal surface views (i)

6.4.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.0078. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.



6.5 Mask visualisation (i)


This section was not generated. No masks/segmentation were deposited.



7 Map analysis (i)

This section contains the results of statistical analysis of the map.


7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



7.2 Volume estimate (i)




The volume at the recommended contour level is 131 $\rm nm^3;$ this corresponds to an approximate mass of 118 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



7.3 Rotationally averaged power spectrum (i)



*Reported resolution corresponds to spatial frequency of 0.331 ${\rm \AA^{-1}}$



8 Fourier-Shell correlation (i)


This section was not generated. No FSC curve or half-maps provided.



9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-25433 and PDB model 7STM. Per-residue inclusion information can be found in section 3 on page 7.


9.1 Map-model overlay (i)



The images above show the 3D surface view of the map at the recommended contour level 0.0078 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.




9.2 Q-score mapped to coordinate model (i)



The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


9.3 Atom inclusion mapped to coordinate model (i)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.0078).



9.4 Atom inclusion (i)



At the recommended contour level, 89% of all backbone atoms, 88% of all non-hydrogen atoms, are inside the map.



9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (0.0078) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.8806	0.5330
А	0.8867	0.5330
В	0.8856	0.5330



