



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 10, 2023 – 03:00 AM EDT

PDB ID : 7SU9
Title : KRAS-G12D specific TCR9a in complex with C*05-GADGVGKSL
Authors : Sim, M.J.W.; Sun, P.D.
Deposited on : 2021-11-16
Resolution : 1.99 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	D	206	 9% 82% 9% 10%
2	E	244	 % 95% . .
3	A	280	 15% 87% 10% .
4	B	100	 11% 92% 7% .
5	C	9	 89% 11%

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 6973 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TCR9a alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	D	186	1440	899	238	295	8	0	0	0

- Molecule 2 is a protein called TCR9a beta chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	E	241	1919	1206	340	368	5	0	0	0

- Molecule 3 is a protein called MHC class I antigen.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	A	273	2243	1398	415	424	6	0	0	0

- Molecule 4 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	B	100	837	533	141	159	4	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	MET	-	initiating methionine	UNP P61769


- Molecule 5 is a protein called KRAS-G12D-9mer with A18L substitution.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
5	C	9	56	33	10	13	0	0	0

- Molecule 6 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	D	1	Total Mg 1 1	0	0
6	E	1	Total Mg 1 1	0	0

- Molecule 7 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	E	1	Total C O 6 3 3	0	0
7	A	1	Total C O 6 3 3	0	0

- Molecule 8 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	E	1	Total O S 5 4 1	0	0
8	E	1	Total O S 5 4 1	0	0


- Molecule 9 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	D	120	Total O 120 120	0	0
9	E	189	Total O 189 189	0	0
9	A	113	Total O 113 113	0	0
9	B	28	Total O 28 28	0	0
9	C	4	Total O 4 4	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: TCR9a alpha chain




- Molecule 2: TCR9a beta chain




- Molecule 3: MHC class I antigen



- Molecule 4: Beta-2-microglobulin



- Molecule 5: KRAS-G12D-9mer with A18L substitution





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	72.72Å 74.18Å 107.39Å 90.00° 101.30° 90.00°	Depositor
Resolution (Å)	38.87 – 1.99 43.82 – 1.99	Depositor EDS
% Data completeness (in resolution range)	98.5 (38.87-1.99) 98.5 (43.82-1.99)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.75 (at 2.00Å)	Xtrriage
Refinement program	PHENIX (1.19.2_4158: ???)	Depositor
R, R_{free}	0.181 , 0.214 0.185 , 0.213	Depositor DCC
R_{free} test set	3648 reflections (4.83%)	wwPDB-VP
Wilson B-factor (Å ²)	28.4	Xtrriage
Anisotropy	0.531	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 43.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6973	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.14% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, GOL, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	D	0.41	0/1467	0.66	1/1992 (0.1%)
2	E	0.41	0/1972	0.62	0/2685
3	A	0.41	0/2305	0.65	0/3128
4	B	0.35	0/860	0.58	0/1162
5	C	0.45	0/55	0.57	0/70
All	All	0.40	0/6659	0.63	1/9037 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	144	ASP	CB-CG-OD1	5.35	123.11	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	1440	0	1380	15	0
2	E	1919	0	1809	10	0
3	A	2243	0	2106	19	0
4	B	837	0	803	7	0
5	C	56	0	58	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	D	1	0	0	0	0
6	E	1	0	0	0	0
7	A	6	0	8	0	0
7	E	6	0	8	1	0
8	E	10	0	0	0	0
9	A	113	0	0	2	0
9	B	28	0	0	0	0
9	C	4	0	0	0	0
9	D	120	0	0	1	0
9	E	189	0	0	1	0
All	All	6973	0	6172	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:83:ASN:HD21	4:B:90:PRO:HG3	1.46	0.80
2:E:174:CYS:SG	2:E:196:ARG:HD3	2.32	0.68
4:B:22:PHE:CE2	4:B:69:GLU:HG2	2.32	0.65
1:D:163:CYS:SG	2:E:174:CYS:HB2	2.37	0.64
3:A:137:ASP:HB2	9:A:401:HOH:O	1.98	0.64
1:D:162:LYS:O	9:D:401:HOH:O	2.15	0.63
3:A:201:LEU:HD12	3:A:249:VAL:HG21	1.81	0.62
1:D:40:GLN:HB3	1:D:50:ILE:HD11	1.82	0.61
3:A:138:LYS:HD2	3:A:138:LYS:H	1.67	0.60
1:D:129:ASP:HB2	1:D:135:LYS:HB2	1.84	0.59
1:D:185:ASP:O	1:D:185:ASP:OD1	2.22	0.58
1:D:163:CYS:SG	2:E:174:CYS:SG	3.04	0.55
2:E:35:VAL:HG13	2:E:94:CYS:SG	2.48	0.53
1:D:163:CYS:SG	2:E:174:CYS:CB	2.98	0.51
1:D:61:GLU:CD	1:D:61:GLU:H	2.15	0.50
3:A:117:ALA:HB2	4:B:60:TRP:CE2	2.47	0.49
3:A:197:HIS:O	3:A:197:HIS:CG	2.65	0.48
2:E:206:ASN:HD21	2:E:208:ARG:CZ	2.26	0.48
3:A:147:TRP:HZ2	5:C:9:LEU:HD23	1.78	0.48
4:B:83:ASN:ND2	4:B:90:PRO:HG3	2.23	0.48
4:B:22:PHE:CZ	4:B:69:GLU:HG2	2.48	0.48
1:D:5:ALA:HB2	1:D:30:ILE:HA	1.96	0.47
1:D:137:VAL:HG13	1:D:178:VAL:HG13	1.96	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:47:PRO:HB2	3:A:53:GLU:HG3	1.97	0.46
1:D:158:TYR:O	1:D:179:ALA:HA	2.17	0.45
1:D:165:LEU:HB3	2:E:174:CYS:HB3	1.98	0.45
1:D:163:CYS:HB3	2:E:174:CYS:HG	1.82	0.45
3:A:138:LYS:NZ	9:A:401:HOH:O	2.13	0.44
2:E:246:ALA:O	9:E:401:HOH:O	2.20	0.44
3:A:218:GLN:HG2	3:A:223:ASP:OD2	2.18	0.43
3:A:260:HIS:HA	3:A:270:LEU:O	2.18	0.43
3:A:255:GLN:H	3:A:255:GLN:CD	2.22	0.43
3:A:19:GLU:H	3:A:19:GLU:CD	2.22	0.42
3:A:219:ARG:HD3	3:A:256:ARG:NH2	2.34	0.42
7:E:301:GOL:H32	3:A:129:ASP:OD2	2.19	0.42
3:A:249:VAL:HG13	3:A:257:TYR:CE1	2.55	0.42
4:B:1:ILE:HA	4:B:1:ILE:HD13	1.80	0.41
1:D:38:TRP:CZ2	1:D:91:CYS:HB2	2.55	0.41
3:A:33:PHE:HB3	3:A:51:TRP:CH2	2.55	0.41
3:A:194:VAL:HG23	3:A:199:ALA:HA	2.02	0.41
3:A:77:ASN:CG	5:C:9:LEU:HG	2.40	0.41
3:A:119:ASP:HB3	4:B:0:MET:HG3	2.02	0.40
1:D:135:LYS:HB3	2:E:131:PHE:HE2	1.86	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	182/206 (88%)	175 (96%)	7 (4%)	0	100	100
2	E	239/244 (98%)	236 (99%)	3 (1%)	0	100	100
3	A	271/280 (97%)	267 (98%)	4 (2%)	0	100	100
4	B	98/100 (98%)	97 (99%)	1 (1%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
5	C	7/9 (78%)	7 (100%)	0	0	100	100
All	All	797/839 (95%)	782 (98%)	15 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	166/184 (90%)	166 (100%)	0	100	100
2	E	208/210 (99%)	205 (99%)	3 (1%)	67	72
3	A	229/235 (97%)	223 (97%)	6 (3%)	46	48
4	B	95/95 (100%)	93 (98%)	2 (2%)	53	57
5	C	5/5 (100%)	5 (100%)	0	100	100
All	All	703/729 (96%)	692 (98%)	11 (2%)	62	67

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	E	50	PHE
2	E	196	ARG
2	E	241	GLU
3	A	69	ARG
3	A	79	ARG
3	A	82	ARG
3	A	113	TYR
3	A	138	LYS
3	A	163	THR
4	B	1	ILE
4	B	70	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
2	E	47	GLN
4	B	83	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
8	SO4	E	302	-	4,4,4	0.13	0	6,6,6	0.09	0
7	GOL	E	301	-	5,5,5	0.85	0	5,5,5	1.43	1 (20%)
7	GOL	A	301	-	5,5,5	0.79	0	5,5,5	1.03	0
8	SO4	E	303	-	4,4,4	0.14	0	6,6,6	0.04	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	GOL	E	301	-	-	2/4/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	GOL	A	301	-	-	2/4/4/4	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	E	301	GOL	C3-C2-C1	-2.20	103.16	111.70

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	E	301	GOL	O1-C1-C2-O2
7	E	301	GOL	O1-C1-C2-C3
7	A	301	GOL	C1-C2-C3-O3
7	A	301	GOL	O2-C2-C3-O3

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	E	301	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	D	186/206 (90%)	0.68	18 (9%) 7 7	23, 30, 71, 81	0
2	E	241/244 (98%)	0.07	3 (1%) 79 78	20, 29, 45, 61	0
3	A	273/280 (97%)	0.82	42 (15%) 2 1	21, 37, 71, 77	0
4	B	100/100 (100%)	0.56	11 (11%) 5 5	25, 46, 66, 76	0
5	C	9/9 (100%)	0.60	0 100 100	23, 24, 28, 30	0
All	All	809/839 (96%)	0.53	74 (9%) 9 8	20, 33, 67, 81	0

All (74) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	A	197	HIS	10.1
3	A	17	ARG	8.3
1	D	185	ASP	8.2
3	A	108	ARG	7.3
1	D	184	SER	7.3
3	A	105	PRO	7.1
3	A	18	GLY	5.7
3	A	42	SER	4.9
1	D	152	SER	4.8
3	A	196	ASP	4.8
3	A	16	GLY	4.7
3	A	19	GLU	4.6
3	A	107	GLY	4.6
1	D	153	LYS	4.5
3	A	268	GLU	4.5
4	B	0	MET	4.4
3	A	41	ALA	4.3
1	D	154	ASP	4.1
3	A	253	GLU	4.1
3	A	181	ARG	3.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
4	B	48	LYS	3.9
3	A	250	PRO	3.9
3	A	195	SER	3.9
3	A	106	ASP	3.7
1	D	186	PHE	3.7
2	E	225	GLU	3.7
3	A	256	ARG	3.6
1	D	156	ASP	3.6
1	D	182	ASN	3.6
3	A	199	ALA	3.6
3	A	255	GLN	3.5
1	D	5	ALA	3.5
3	A	274	TRP	3.5
3	A	104	GLY	3.3
3	A	247	VAL	3.3
3	A	249	VAL	3.3
1	D	193	ASN	3.2
3	A	194	VAL	3.2
1	D	190	ASN	3.2
3	A	273	ARG	3.1
2	E	222	GLU	3.0
3	A	257	TYR	3.0
3	A	220	ASP	3.0
3	A	252	GLY	2.9
3	A	198	GLU	2.9
1	D	136	SER	2.8
3	A	177	LYS	2.8
3	A	89	GLU	2.7
4	B	1	ILE	2.6
3	A	267	PRO	2.6
4	B	74	GLU	2.6
1	D	131	LYS	2.6
3	A	2	SER	2.6
3	A	271	THR	2.5
4	B	46	ILE	2.5
4	B	75	LYS	2.5
4	B	77	GLU	2.5
3	A	99	TYR	2.4
1	D	183	LYS	2.4
3	A	270	LEU	2.4
3	A	219	ARG	2.4
1	D	128	ARG	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	A	248	VAL	2.3
3	A	227	ASP	2.3
3	A	272	LEU	2.3
4	B	47	GLU	2.2
1	D	103	LEU	2.2
1	D	189	ALA	2.2
4	B	56	PHE	2.2
2	E	223	ASN	2.2
1	D	188	CYS	2.2
3	A	193	PRO	2.1
4	B	69	GLU	2.1
4	B	44	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
7	GOL	A	301	6/6	0.67	0.21	31,37,43,47	0
7	GOL	E	301	6/6	0.86	0.19	33,36,44,48	0
6	MG	E	304	1/1	0.89	0.06	51,51,51,51	0
6	MG	D	301	1/1	0.91	0.10	52,52,52,52	0
8	SO4	E	303	5/5	0.96	0.23	59,72,87,91	0
8	SO4	E	302	5/5	0.97	0.15	38,60,69,77	0

6.5 Other polymers [i](#)

There are no such residues in this entry.