



# Full wwPDB X-ray Structure Validation Report ⓘ

May 8, 2024 – 04:04 PM EDT

PDB ID : 8SVQ  
Title : Crystal structure of pregnane X receptor ligand binding domain in complex with SJPYT-312  
Authors : Garcia-Maldonado, E.; Huber, A.D.; Nithianantham, S.; Miller, D.J.; Chen, T.  
Deposited on : 2023-05-17  
Resolution : 2.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36.2  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1235 (2.78-2.74)
Clashscore	141614	1277 (2.78-2.74)
Ramachandran outliers	138981	1257 (2.78-2.74)
Sidechain outliers	138945	1257 (2.78-2.74)
RSRZ outliers	127900	1207 (2.78-2.74)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	351	 5% 75% 20%
1	B	351	 6% 74% 7% 19%

## 2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 4556 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pregnane X receptor ligand binding domain fused to SRC-1 coactivator peptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	280	2224	1437	376	393	18	0	0	0
1	B	283	2246	1441	385	401	19	0	1	0

There are 26 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	123	HIS	-	expression tag	UNP O75469
A	124	HIS	-	expression tag	UNP O75469
A	125	HIS	-	expression tag	UNP O75469
A	126	HIS	-	expression tag	UNP O75469
A	127	HIS	-	expression tag	UNP O75469
A	128	HIS	-	expression tag	UNP O75469
A	129	GLY	-	expression tag	UNP O75469
A	435	SER	-	linker	UNP O75469
A	436	GLY	-	linker	UNP O75469
A	437	GLY	-	linker	UNP O75469
A	438	SER	-	linker	UNP O75469
A	439	GLY	-	linker	UNP O75469
A	440	GLY	-	linker	UNP O75469
B	123	HIS	-	expression tag	UNP O75469
B	124	HIS	-	expression tag	UNP O75469
B	125	HIS	-	expression tag	UNP O75469
B	126	HIS	-	expression tag	UNP O75469
B	127	HIS	-	expression tag	UNP O75469
B	128	HIS	-	expression tag	UNP O75469
B	129	GLY	-	expression tag	UNP O75469
B	435	SER	-	linker	UNP O75469
B	436	GLY	-	linker	UNP O75469
B	437	GLY	-	linker	UNP O75469
B	438	SER	-	linker	UNP O75469

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	439	GLY	-	linker	UNP O75469
B	440	GLY	-	linker	UNP O75469

- Molecule 2 is (1P)-N-(5-tert-butyl-2-[[[(3S)-hexan-3-yl]oxy]phenyl]-1-(2,5-dimethoxyphenyl)-5-methyl-1H-1,2,3-triazole-4-carboxamide (three-letter code: WSX) (formula: C<sub>28</sub>H<sub>38</sub>N<sub>4</sub>O<sub>4</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	36	28	4	4	0	0
2	B	1	36	28	4	4	0	0

- Molecule 3 is DIMETHYL SULFOXIDE (three-letter code: DMS) (formula: C<sub>2</sub>H<sub>6</sub>OS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	O	S		
3	B	1	4	2	1	1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	2	2	2	0	0
4	B	8	8	8	0	0


### 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Pregnane X receptor ligand binding domain fused to SRC-1 coactivator peptide



- Molecule 1: Pregnane X receptor ligand binding domain fused to SRC-1 coactivator peptide



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	84.33Å 87.90Å 104.53Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	67.28 – 2.75 67.28 – 2.75	Depositor EDS
% Data completeness (in resolution range)	100.0 (67.28-2.75) 100.0 (67.28-2.75)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.33 (at 2.73Å)	Xtrriage
Refinement program	PHENIX (1.20.1_4487)	Depositor
R, $R_{free}$	0.230 , 0.262 0.228 , 0.259	Depositor DCC
$R_{free}$ test set	1000 reflections (4.81%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	75.9	Xtrriage
Anisotropy	0.105	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 59.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.011 for k,h,-l	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	4556	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	79.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.56% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: WSX, DMS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.24	0/2272	0.42	0/3069
1	B	0.25	0/2294	0.44	0/3095
All	All	0.24	0/4566	0.43	0/6164

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2224	0	2179	9	0
1	B	2246	0	2205	15	0
2	A	36	0	0	0	0
2	B	36	0	0	0	0
3	B	4	0	6	0	0
4	A	2	0	0	0	0
4	B	8	0	0	0	0
All	All	4556	0	4390	23	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.



All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:412:LEU:HD21	1:B:429:PHE:HB3	1.78	0.65
1:B:216:ARG:NH2	1:B:302:GLY:O	2.29	0.64
1:A:216:ARG:NH2	1:A:302:GLY:O	2.31	0.64
1:A:237:PHE:HA	1:A:240:LEU:HD23	1.82	0.61
1:A:254:ILE:HD12	1:A:283:LEU:HB3	1.87	0.55
1:A:221:SER:HB3	1:B:228:PRO:HD3	1.89	0.54
1:B:254:ILE:HD12	1:B:283:LEU:HB3	1.92	0.51
1:A:279:ALA:HB1	1:A:283:LEU:HD13	1.93	0.51
1:B:319:LEU:HD12	1:B:325:LYS:HA	1.94	0.49
1:B:216:ARG:HG2	1:B:222:VAL:HG22	1.95	0.49
1:B:279:ALA:HB1	1:B:283:LEU:HD13	1.95	0.48
1:A:415:GLN:HG2	1:A:431:ILE:HD13	1.96	0.47
1:B:455:ARG:NH1	1:B:459:GLU:OE2	2.48	0.46
1:B:214:GLN:HB3	1:B:305:SER:HB2	1.98	0.46
1:B:411:LEU:HG	1:B:420:PHE:HZ	1.82	0.45
1:B:273:ILE:HG23	1:B:453:LEU:HD23	1.98	0.45
1:A:382:PRO:HD2	1:A:386:HIS:CE1	2.52	0.44
1:A:381:ARG:HB3	1:A:386:HIS:ND1	2.33	0.43
1:B:332:LYS:HE3	1:B:332:LYS:HB2	1.81	0.43
1:A:208:SER:HB2	1:A:209:LEU:HD12	2.01	0.43
1:B:349:PHE:O	1:B:366:GLN:HB2	2.21	0.41
1:B:324:LEU:HD23	1:B:324:LEU:HA	1.92	0.41
1:B:214:GLN:NE2	1:B:216:ARG:HH11	2.19	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [\(i\)](#)

### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	270/351 (77%)	264 (98%)	6 (2%)	0	<b>100</b> <b>100</b>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	274/351 (78%)	270 (98%)	4 (2%)	0	100	100
All	All	544/702 (78%)	534 (98%)	10 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	236/310 (76%)	236 (100%)	0	100	100
1	B	239/310 (77%)	238 (100%)	1 (0%)	91	93
All	All	475/620 (77%)	474 (100%)	1 (0%)	93	96

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	162	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	214	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	WSX	A	502	-	34,38,38	2.31	9 (26%)	49,54,54	1.48	6 (12%)
3	DMS	B	504	-	3,3,3	0.61	0	3,3,3	0.33	0
2	WSX	B	502	-	34,38,38	2.33	9 (26%)	49,54,54	1.51	8 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	WSX	A	502	-	-	9/28/31/31	0/3/3/3
2	WSX	B	502	-	-	11/28/31/31	0/3/3/3

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	502	WSX	C04-N03	6.98	1.55	1.41
2	A	502	WSX	C04-N03	6.88	1.54	1.41
2	B	502	WSX	O06-C05	5.45	1.46	1.37
2	A	502	WSX	O06-C05	5.29	1.46	1.37
2	A	502	WSX	C02-N03	4.93	1.48	1.35
2	B	502	WSX	C02-N03	4.89	1.48	1.35
2	B	502	WSX	O29-C28	4.07	1.43	1.37
2	A	502	WSX	O29-C28	4.02	1.43	1.37
2	B	502	WSX	C21-C02	3.45	1.55	1.50

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	502	WSX	O34-C33	3.34	1.44	1.37
2	B	502	WSX	O34-C33	3.26	1.44	1.37
2	B	502	WSX	C16-C15	3.10	1.60	1.53
2	A	502	WSX	C21-C02	3.09	1.55	1.50
2	A	502	WSX	C16-C15	3.01	1.60	1.53
2	A	502	WSX	C23-C22	2.63	1.55	1.49
2	B	502	WSX	C23-C22	2.59	1.55	1.49
2	A	502	WSX	C19-C16	-2.29	1.45	1.53
2	B	502	WSX	C19-C16	-2.27	1.45	1.53

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	502	WSX	O06-C05-C04	6.45	121.40	114.63
2	A	502	WSX	O06-C05-C04	5.98	120.90	114.63
2	B	502	WSX	C28-C27-N24	3.08	120.80	117.87
2	A	502	WSX	C28-C27-N24	2.81	120.55	117.87
2	A	502	WSX	C23-C22-C21	-2.80	126.34	129.71
2	A	502	WSX	O29-C28-C31	-2.54	120.01	124.37
2	B	502	WSX	C23-C22-C21	-2.54	126.66	129.71
2	B	502	WSX	C04-N03-C02	-2.43	120.28	126.93
2	A	502	WSX	C30-O29-C28	-2.27	114.10	117.53
2	B	502	WSX	C30-O29-C28	-2.27	114.10	117.53
2	B	502	WSX	C22-N24-N25	-2.22	108.87	112.40
2	B	502	WSX	O06-C05-C13	-2.22	118.65	123.87
2	B	502	WSX	O29-C28-C31	-2.18	120.63	124.37
2	A	502	WSX	C05-C04-N03	2.13	120.49	116.66

There are no chirality outliers.

All (20) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	502	WSX	O01-C02-C21-C22
2	A	502	WSX	C04-C05-O06-C07
2	B	502	WSX	O06-C07-C10-C11
2	B	502	WSX	O01-C02-C21-C22
2	B	502	WSX	C04-C05-O06-C07
2	B	502	WSX	C32-C33-O34-C35
2	B	502	WSX	C36-C33-O34-C35
2	A	502	WSX	O06-C07-C08-C09
2	B	502	WSX	O06-C07-C08-C09
2	A	502	WSX	C27-C28-O29-C30

*Continued on next page...*


*Continued from previous page...*


Mol	Chain	Res	Type	Atoms
2	A	502	WSX	C07-C10-C11-C12
2	B	502	WSX	C07-C10-C11-C12
2	A	502	WSX	C31-C28-O29-C30
2	A	502	WSX	C10-C07-C08-C09
2	B	502	WSX	C08-C07-C10-C11
2	B	502	WSX	C13-C05-O06-C07
2	B	502	WSX	C10-C07-C08-C09
2	A	502	WSX	C13-C05-O06-C07
2	B	502	WSX	C05-C04-N03-C02
2	A	502	WSX	C05-C04-N03-C02

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	280/351 (79%)	0.58	16 (5%) 23 28	52, 75, 124, 138	0
1	B	283/351 (80%)	0.62	22 (7%) 13 15	48, 74, 131, 203	0
All	All	563/702 (80%)	0.60	38 (6%) 17 21	48, 75, 125, 203	0

All (38) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	206	LEU	6.9
1	B	196	ALA	5.3
1	A	209	LEU	4.9
1	B	208	SER	4.8
1	B	197	ALA	4.5
1	A	199	TRP	4.5
1	B	411	LEU	4.1
1	A	446	LEU	3.8
1	B	202	VAL	3.7
1	B	213	LEU	3.5
1	B	319	LEU	3.5
1	B	205	ASP	3.4
1	A	197	ALA	3.3
1	A	206	LEU	3.2
1	B	209	LEU	3.1
1	A	447	THR	3.1
1	B	207	CYS	2.9
1	A	417	ILE	2.8
1	A	359	HIS	2.8
1	A	251	PHE	2.7
1	A	431	ILE	2.7
1	B	309	GLU	2.7
1	B	456	LEU	2.6
1	B	251	PHE	2.6

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	318	LEU	2.5
1	B	237	PHE	2.5
1	A	213	LEU	2.4
1	A	269	ILE	2.4
1	A	407	HIS	2.3
1	B	216	ARG	2.3
1	B	230	ASP	2.3
1	B	240	LEU	2.2
1	A	196	ALA	2.2
1	A	177	VAL	2.1
1	A	362	VAL	2.1
1	B	215	LEU	2.1
1	B	238	SER	2.0
1	B	199	TRP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	DMS	B	504	4/4	0.87	0.25	80,87,88,97	0
2	WSX	A	502	36/36	0.89	0.37	70,85,101,109	0
2	WSX	B	502	36/36	0.90	0.34	68,82,98,102	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around WSX A 502:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around WSX B 502:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.