

wwPDB X-ray Structure Validation Summary Report (i)

Aug 8, 2020 – 12:19 PM BST

PDB ID : 6TK5

Title: Femtosecond to millisecond structural changes in a light-driven sodium pump:

800fs+2ps structure of KR2 with extrapolated, light and dark datasets

Authors: Skopintsev, P.; Ehrenberg, D.; Weinert, T.; James, D.; Kar, R.; Johnson, P.;

Ozerov, D.; Furrer, A.; Martiel, I.; Dworkowski, F.; Nass, K.; Knopp, G.; Cirelli, C.; Gashi, D.; Mous, S.; Wranik, M.; Gruhl, T.; Kekilli, D.; Bruenle, S.; Deupi, X.; Schertler, G.F.X.; Benoit, R.; Panneels, V.; Nogly, P.; Schapiro,

I.; Milne, C.; Heberle, J.; Standfuss, J.

Deposited on : 2019-11-28

Resolution : 2.25 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

 $Mol Probity \quad : \quad 4.02b\text{--}467$

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.13.1

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.13.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \text{Whole archive} \\ (\#\text{Entries}) \end{array}$	$\begin{array}{c} {\rm Similar \; resolution} \\ (\#{\rm Entries, \; resolution \; range(\AA)}) \end{array}$
R_{free}	130704	1377 (2.26-2.26)
Clashscore	141614	1487 (2.26-2.26)
Ramachandran outliers	138981	1449 (2.26-2.26)
Sidechain outliers	138945	1450 (2.26-2.26)
RSRZ outliers	127900	1356 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain			
			3%			
1	A	290	78%	12%	•	9%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:



Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	LFA	A	306	-	-	-	X
3	LFA	A	315	-	-	-	X
3	LFA	A	316	-	-	-	X
3	LFA	A	321	-	-	-	X
3	LFA	A	322	-	-	-	X



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 2398 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Sodium pumping rhodopsin.


\mathbf{Mol}	Chain	Residues	Atoms			ZeroOcc	${f AltConf}$	Trace		
1	A	265	Total 2103	C 1405	N 320	O 369	S 9	0	4	0

There are 15 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	276	GLU	_	expression tag	UNP N0DKS8
A	277	ASN	-	expression tag	UNP N0DKS8
A	278	LEU	_	expression tag	UNP N0DKS8
A	279	TYR	-	expression tag	UNP N0DKS8
A	280	PHE	_	expression tag	UNP N0DKS8
A	281	GLN	_	expression tag	UNP N0DKS8
A	282	SER	-	expression tag	UNP N0DKS8
A	283	GLY	_	expression tag	UNP N0DKS8
A	284	SER	-	expression tag	UNP N0DKS8
A	285	HIS	-	expression tag	UNP N0DKS8
A	286	HIS	_	expression tag	UNP N0DKS8
A	287	HIS	-	expression tag	UNP N0DKS8
A	288	HIS	_	expression tag	UNP N0DKS8
A	289	HIS	=	expression tag	UNP N0DKS8
A	290	HIS	-	expression tag	UNP N0DKS8


• Molecule 2 is RETINAL (three-letter code: RET) (formula: C₂₀H₂₈O) (labeled as "Ligand of Interest" by author).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C 20 20	0	0

 \bullet Molecule 3 is EICOSANE (three-letter code: LFA) (formula: $\mathrm{C}_{20}\mathrm{H}_{42}).$



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C 16 16	0	0
3	A	1	Total C 8 8	0	0
3	A	1	Total C 6 6	0	0

Continued on next page...



Continued from previous page...

Mol		$egin{array}{c c} n & previous & pa \\ \hline \mathbf{Residues} \end{array}$	Atoms	ZeroOcc	AltConf
3	A	1	Total C 6 6	0	0
3	A	1	Total C 5 5	0	0
3	A	1	Total C 13 13	0	0
3	A	1	Total C 10 10	0	0
3	A	1	Total C 8 8	0	0
3	A	1	Total C 8 8	0	0
3	A	1	Total C 12 12	0	0
3	A	1	Total C 16 16	0	0
3	A	1	Total C 5 5	0	0
3	A	1	Total C 10 10	0	0
3	A	1	Total C 5 5	0	0
3	A	1	Total C 11 11	0	0
3	A	1	Total C 9 9	0	0
3	A	1	Total C 9 9	0	0
3	A	1	Total C 6 6	0	0
3	A	1	Total C 5 5	0	0
3	A	1	Total C 4 4	0	0
3	A	1	Total C 6 6	0	0
3	A	1	Total C 10 10	0	0
3	A	1	Total C	0	0
3	A	1	Total C 9 9	0	0

Continued on next page...



Continued from previous page...

Mol	Chain	Residues	${f Atoms}$	ZeroOcc	AltConf
3	A	1	Total C 10 10	0	0
3	A	1	Total C 12 12	0	0

• Molecule 4 is water.


Mol	Chain	Residues	${f Atoms}$	ZeroOcc	AltConf
4	A	53	Total O 53 53	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Sodium pumping rhodopsin





4 Data and refinement statistics (i)

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants	41.54Å 84.48Å 235.56Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	12.36 - 2.25	Depositor
resolution (A)	12.36 - 2.25	EDS
% Data completeness	89.4 (12.36-2.25)	Depositor
(in resolution range)	89.4 (12.36-2.25)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	5.28 (at 2.26Å)	Xtriage
Refinement program	PHENIX 1.16_3549	Depositor
R, R_{free}	0.272 , 0.328	Depositor
It, It free	0.272 , 0.328	DCC
R_{free} test set	1168 reflections (6.48%)	wwPDB-VP
Wilson B-factor (Å ²)	31.7	Xtriage
Anisotropy	0.432	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.39 , 118.9	EDS
L-test for twinning ²	$ < L >=0.51, < L^2>=0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	2398	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 6.54% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $< L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: LFA, RET

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
		RMSZ	# Z >5	RMSZ	# Z > 5	
1	A	0.43	0/2173	0.53	0/2959	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2103	0	2078	25	0
2	A	20	0	27	5	0
3	A	222	0	403	12	0
4	A	53	0	0	3	0
All	All	2398	0	2508	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 35 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	$egin{aligned} ext{Interatomic} \ ext{distance} \ (ext{Å}) \end{aligned}$	$egin{aligned} ext{Clash} \ ext{overlap } (ext{Å}) \end{aligned}$
1:A:114:LEU:HD22	3:A:307:LFA:H31	1.67	0.76
1:A:8:ALA:HB3	1:A:12:ASN:HD22	1.51	0.76
1:A:189:ILE:HD12	1:A:208:TRP:HB2	1.80	0.62
3:A:302:LFA:H141	3:A:316:LFA:H111	1.80	0.61
3:A:312:LFA:H61	3:A:323:LFA:H61	1.86	0.58

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	\mathbf{ntiles}
1	A	$267/290 \ (92\%)$	257 (96%)	10 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	A	$222/250 \ (89\%)$	210 (95%)	12 (5%)	22 22		

5 of 12 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	157	GLN
1	A	161	VAL

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type
1	A	224	MET
1	A	142	PHE
1	A	189	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	26	GLN
1	A	61	ASN
1	A	180	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

27 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	ol Type Chain Res L		Link	Bo	Bond lengths			Bond angles		
WIOI	туре	Chain	res	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	$\mid \# Z > 2 \mid$
3	LFA	A	314	-	9,9,19	0.40	0	8,8,18	0.66	0
3	LFA	A	319	-	5,5,19	0.32	0	4,4,18	0.54	0
3	LFA	A	313	-	4,4,19	0.33	0	3,3,18	0.57	0



Mal	T	Chain	Dog	T : 1-	Bo	Bond lengths			ond ang	les
Mol	Type	Chain	Res	Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	LFA	A	327	_	11,11,19	0.34	0	10,10,18	0.78	0
3	LFA	A	326	-	9,9,19	0.37	0	8,8,18	0.61	0
2	RET	A	301	1	20,20,21	0.81	1 (5%)	27,27,28	0.79	1 (3%)
3	LFA	A	316	-	10,10,19	0.35	0	9,9,18	0.70	0
3	LFA	A	312	-	15,15,19	0.36	0	14,14,18	0.72	0
3	LFA	A	325	-	8,8,19	0.35	0	7,7,18	0.70	0
3	LFA	A	315	-	4,4,19	0.39	0	3,3,18	0.43	0
3	LFA	A	321	_	3,3,19	0.43	0	2,2,18	0.70	0
3	LFA	A	302	_	15,15,19	0.34	0	14,14,18	0.80	0
3	LFA	A	310	_	7,7,19	0.36	0	6,6,18	0.65	0
3	LFA	A	323	_	9,9,19	0.36	0	8,8,18	0.59	0
3	LFA	A	308	-	9,9,19	0.31	0	8,8,18	0.79	0
3	LFA	A	304	_	5,5,19	0.35	0	4,4,18	0.52	0
3	LFA	A	320	-	4,4,19	0.34	0	3,3,18	0.45	0
3	LFA	A	309	_	7,7,19	0.44	0	6,6,18	0.42	0
3	LFA	A	307	_	12,12,19	0.34	0	11,11,18	0.72	0
3	LFA	A	305	-	5,5,19	0.36	0	4,4,18	0.44	0
3	LFA	A	306	_	4,4,19	0.33	0	3,3,18	0.54	0
3	LFA	A	311	-	11,11,19	0.39	0	10,10,18	0.70	0
3	LFA	A	322	-	5,5,19	0.35	0	4,4,18	0.52	0
3	LFA	A	303	-	7,7,19	0.30	0	6,6,18	0.70	0
3	LFA	A	318	-	8,8,19	0.42	0	7,7,18	0.51	0
3	LFA	A	324	-	2,2,19	0.33	0	0,1,18	0.00	-
3	LFA	A	317	-	8,8,19	0.33	0	7,7,18	0.74	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	LFA	A	314	_	-	1/7/7/17	-
3	LFA	A	319	-	-	1/3/3/17	-
3	LFA	A	313	_	_	1/2/2/17	_
3	LFA	A	327	-	-	4/9/9/17	-
3	LFA	A	326	-	-	5/7/7/17	-
2	RET	A	301	1	-	0/13/30/31	0/1/1/1
3	LFA	A	316	_	-	5/8/8/17	-
3	LFA	A	312	_	-	6/13/13/17	-
3	LFA	A	325	_	-	4/6/6/17	-
3	LFA	A	315	_	_	0/2/2/17	-

Continued on next page...



Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	LFA	A	321	-	-	1/1/1/17	-
3	LFA	A	302	-	-	6/13/13/17	-
3	LFA	A	310	-	-	2/5/5/17	-
3	LFA	A	323	-	-	3/7/7/17	-
3	LFA	A	308	-	-	5/7/7/17	-
3	LFA	A	304	-	-	0/3/3/17	-
3	LFA	A	320	-	-	0/2/2/17	-
3	LFA	A	309	-	-	1/5/5/17	-
3	LFA	A	307	-	-	5/10/10/17	-
3	LFA	A	305	-	-	1/3/3/17	-
3	LFA	A	306	-	-	2/2/2/17	-
3	LFA	A	311	-	-	3/9/9/17	-
3	LFA	A	322	-	-	1/3/3/17	-
3	LFA	A	303	-		2/5/5/17	
3	LFA	A	318	-		3/6/6/17	
3	LFA	A	317	-	-	3/6/6/17	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(\mathbf{\mathring{A}})$	$\operatorname{Ideal}(ext{\AA})$
2	A	301	RET	C14-C13	2.85	1.36	1.33

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^o)$
2	A	301	RET	C8-C7-C6	2.51	134.25	127.20

There are no chirality outliers.

5 of 65 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	321	LFA	C3-C4-C5-C6
3	A	318	LFA	C3-C4-C5-C6
3	A	323	LFA	C3-C4-C5-C6
3	A	326	LFA	C6-C7-C8-C9
3	A	310	LFA	C4-C5-C6-C7


There are no ring outliers.

9 monomers are involved in 17 short contacts:





Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	326	LFA	1	0
2	A	301	RET	5	0
3	A	316	LFA	3	0
3	A	312	LFA	5	0
3	A	302	LFA	4	0
3	A	310	LFA	1	0
3	A	323	LFA	2	0
3	A	307	LFA	1	0
3	A	318	LFA	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.






























































5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\#\mathrm{RSRZ}{>}2$		$\mathrm{OWAB}(\mathrm{\AA}^2)$	Q < 0.9	
1	A	$265/290 \ (91\%)$	-0.04	9 (3%)	45	47	22, 40, 65, 89	0

The worst 5 of 9 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	272	ASN	5.8
1	A	158	PHE	4.6
1	A	88	PHE	2.8
1	A	231	ASP	2.6
1	A	271	LYS	2.6

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	${f Res}$	Atoms	RSCC	RSR	${f B\text{-factors}}({f A}^2)$	Q<0.9
3	LFA	A	311	12/20	0.30	0.31	59,79,85,89	0
3	LFA	A	306	5/20	0.32	0.47	74,85,103,106	0

Continued on next page...



Continued from previous page...


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}({ ext{\AA}}^2)$	Q<0.9
3	LFA	A	315	5/20	0.42	0.69	71,74,81,89	0
3	LFA	A	309	8/20	0.47	0.36	72,78,84,89	0
3	LFA	A	321	4/20	0.54	0.55	60,68,74,74	0
3	LFA	A	318	9/20	0.54	0.29	59,71,78,78	0
3	LFA	A	326	10/20	0.55	0.27	58,70,74,75	0
3	LFA	A	314	10/20	0.57	0.26	52,61,77,85	0
3	LFA	A	324	3/20	0.57	0.23	60,60,83,86	0
3	LFA	A	322	6/20	0.62	0.43	76,78,83,87	0
3	LFA	A	317	9/20	0.62	0.34	64,78,90,95	0
3	LFA	A	312	16/20	0.66	0.19	60,69,79,80	0
3	LFA	A	302	16/20	0.67	0.22	53,72,74,80	0
3	LFA	A	316	11/20	0.67	0.43	64,80,96,98	0
3	LFA	A	308	10/20	0.71	0.26	$46,\!59,\!65,\!79$	0
3	LFA	A	325	9/20	0.72	0.33	59,62,68,72	0
3	LFA	A	307	13/20	0.72	0.23	54,60,69,73	0
3	LFA	A	319	6/20	0.73	0.28	53,59,63,66	0
3	LFA	A	323	10/20	0.75	0.30	67,78,82,91	0
3	LFA	A	304	6/20	0.77	0.30	61,69,72,74	0
3	LFA	A	320	5/20	0.77	0.16	57,59,62,68	0
3	LFA	A	310	8/20	0.78	0.25	46,50,56,59	0
3	LFA	A	305	6/20	0.78	0.34	54,64,74,75	0
3	LFA	A	327	12/20	0.79	0.30	39,57,60,61	0
3	LFA	A	313	5/20	0.83	0.25	54,55,68,71	0
3	LFA	A	303	8/20	0.86	0.20	40,45,49,53	8
2	RET	A	301	20/21	0.94	0.12	21,35,44,45	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around LFA A 311:


 $2 \mathrm{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









Electron density around LFA A 306:












Electron density around LFA A 315:


 $2 \text{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\text{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around LFA A 309:
















Electron density around LFA A 318:


 $2 \mathrm{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around LFA A 326:

















Electron density around LFA A 317: 2mF_o-DF_c (at 0.7 rmsd) in gray mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)


Electron density around LFA A 312: 2mF_o-DF_c (at 0.7 rmsd) in gray mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)




Electron density around LFA A 302:


 $2 \mathrm{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









Electron density around LFA A 316:












Electron density around LFA A 308:


 $2 \mathrm{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









Electron density around LFA A 325:












Electron density around LFA A 307:


 $2 \mathrm{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









Electron density around LFA A 319:












Electron density around LFA A 323:


 $2 \mathrm{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









Electron density around LFA A 310:












Electron density around LFA A 305:


 $2 \mathrm{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around LFA A 327:
















Electron density around LFA A 303:


 $2 \mathrm{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









Electron density around RET A 301:









6.5 Other polymers (i)

There are no such residues in this entry.

