



Full wwPDB X-ray Structure Validation Report

Dec 9, 2020 – 12:07 am GMT

PDB ID : 6TLH
Title : Crystal structure of mouse KANK3 ankyrin repeats
Authors : Khan, R.; Singh, A.K.; Goult, B.T.
Deposited on : 2019-12-02
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.15.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.15.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	256	 2% 89% 6%
1	B	256	 6% 85% 7% 7%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3950 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called KN motif and ankyrin repeat domain-containing protein 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	241	1777	1087	323	356	11	0	0	0
1	B	239	1770	1082	324	353	11	0	1	0

There are 12 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	518	GLY	-	expression tag	UNP Q9Z1P7
A	519	ILE	-	expression tag	UNP Q9Z1P7
A	520	ASP	-	expression tag	UNP Q9Z1P7
A	521	PRO	-	expression tag	UNP Q9Z1P7
A	522	PHE	-	expression tag	UNP Q9Z1P7
A	523	THR	-	expression tag	UNP Q9Z1P7
B	518	GLY	-	expression tag	UNP Q9Z1P7
B	519	ILE	-	expression tag	UNP Q9Z1P7
B	520	ASP	-	expression tag	UNP Q9Z1P7
B	521	PRO	-	expression tag	UNP Q9Z1P7
B	522	PHE	-	expression tag	UNP Q9Z1P7
B	523	THR	-	expression tag	UNP Q9Z1P7

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is FORMIC ACID (three-letter code: FMT) (formula: CH₂O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			3	1	2		


- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	191	Total	O	0	0
			191	191		
4	B	179	Total	O	0	0
			179	179		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: KN motif and ankyrin repeat domain-containing protein 3

Chain A:  2% 89% 6%



- Molecule 1: KN motif and ankyrin repeat domain-containing protein 3

Chain B:  6% 85% 7% 7%



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	149.44Å 45.78Å 69.60Å 90.00° 95.06° 90.00°	Depositor
Resolution (Å)	53.17 – 1.80 53.12 – 1.80	Depositor EDS
% Data completeness (in resolution range)	97.1 (53.17-1.80) 97.1 (53.12-1.80)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.73 (at 1.80Å)	Xtrriage
Refinement program	REFMAC 5.8.0222	Depositor
R, R_{free}	0.199 , 0.236 0.209 , 0.241	Depositor DCC
R_{free} test set	2143 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	22.7	Xtrriage
Anisotropy	0.793	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 42.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.53$, $\langle L^2 \rangle = 0.37$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3950	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 40.31 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.7886e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, FMT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.67	0/1795	0.76	1/2436 (0.0%)
1	B	0.66	0/1789	0.76	0/2428
All	All	0.66	0/3584	0.76	1/4864 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	6
1	B	0	10
All	All	0	16

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	547	ASN	CB-CA-C	5.62	121.63	110.40

There are no chirality outliers.

All (16) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	534	ARG	Sidechain
1	A	536	ARG	Sidechain
1	A	550	ARG	Sidechain
1	A	561	ARG	Sidechain
1	A	569	ARG	Sidechain

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
1	A	725	ARG	Sidechain
1	B	536	ARG	Sidechain
1	B	548	ARG	Sidechain
1	B	561	ARG	Sidechain
1	B	569	ARG	Sidechain
1	B	575	ARG	Sidechain
1	B	725	ARG	Sidechain
1	B	766	LEU	Mainchain
1	B	767[A]	HIS	Mainchain
1	B	767[B]	HIS	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1777	0	1755	10	0
1	B	1770	0	1744	9	0
2	A	6	0	8	1	0
2	B	24	0	32	3	0
3	A	3	0	1	0	0
4	A	191	0	0	4	0
4	B	179	0	0	2	0
All	All	3950	0	3540	20	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:532:ASN:CG	1:B:533:PRO:HD2	2.13	0.68
1:B:562:LEU:O	1:B:562:LEU:HD12	1.95	0.66
1:A:562:LEU:O	1:A:562:LEU:HD12	1.99	0.63
1:A:761:GLU:HG3	4:A:1015:HOH:O	2.06	0.55
1:A:550:ARG:HG3	1:A:550:ARG:NH2	2.25	0.52
1:B:767[B]:HIS:CE1	4:B:906:HOH:O	2.62	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:722:GLU:OE2	2:B:804:GOL:H12	2.09	0.51
1:A:550:ARG:HH21	1:A:550:ARG:HG3	1.76	0.49
1:B:719:CYS:HA	2:B:804:GOL:H11	1.94	0.48
1:B:568:PHE:HB3	2:B:801:GOL:H31	1.95	0.48
1:A:639:ARG:CZ	1:A:639:ARG:HB3	2.45	0.47
1:A:619:ASN:ND2	1:A:622:ILE:HD12	2.32	0.45
1:B:649:ALA:HA	1:B:695:MET:HG2	2.00	0.42
1:A:579:GLU:HB2	4:A:1047:HOH:O	2.18	0.42
1:A:534:ARG:NH2	1:A:534:ARG:HG2	2.34	0.42
1:B:620:LEU:HD13	1:B:658:GLU:HB3	2.02	0.42
1:B:589:LYS:HE2	4:B:992:HOH:O	2.21	0.41
1:A:744:ASP:OD1	1:A:744:ASP:C	2.58	0.41
1:A:569:ARG:HD3	4:A:1072:HOH:O	2.20	0.41
2:A:801:GOL:H2	4:A:1021:HOH:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	239/256 (93%)	237 (99%)	2 (1%)	0	100	100
1	B	238/256 (93%)	235 (99%)	3 (1%)	0	100	100
All	All	477/512 (93%)	472 (99%)	5 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	186/198 (94%)	182 (98%)	4 (2%)	52	39
1	B	185/198 (93%)	179 (97%)	6 (3%)	39	25
All	All	371/396 (94%)	361 (97%)	10 (3%)	49	31

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	534	ARG
1	A	550	ARG
1	A	579	GLU
1	A	725	ARG
1	B	536	ARG
1	B	655	GLN
1	B	725	ARG
1	B	767[A]	HIS
1	B	767[B]	HIS
1	B	769	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	565	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GOL	B	803	-	5,5,5	0.50	0	5,5,5	0.62	0
2	GOL	B	804	-	5,5,5	0.39	0	5,5,5	0.48	0
2	GOL	A	801	-	5,5,5	0.73	0	5,5,5	1.46	0
2	GOL	B	802	-	5,5,5	0.66	0	5,5,5	0.70	0
2	GOL	B	801	-	5,5,5	0.59	0	5,5,5	0.90	0
3	FMT	A	802	-	0,2,2	0.00	-	0,1,1	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	B	803	-	-	0/4/4/4	-
2	GOL	B	804	-	-	4/4/4/4	-
2	GOL	B	801	-	-	3/4/4/4	-
2	GOL	A	801	-	-	4/4/4/4	-
2	GOL	B	802	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (13) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	804	GOL	O1-C1-C2-C3
2	B	804	GOL	C1-C2-C3-O3
2	A	801	GOL	C1-C2-C3-O3
2	B	801	GOL	O1-C1-C2-C3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	B	801	GOL	O1-C1-C2-O2
2	B	802	GOL	O1-C1-C2-C3
2	B	804	GOL	O1-C1-C2-O2
2	B	804	GOL	O2-C2-C3-O3
2	A	801	GOL	O1-C1-C2-O2
2	A	801	GOL	O2-C2-C3-O3
2	B	801	GOL	O2-C2-C3-O3
2	B	802	GOL	O1-C1-C2-O2
2	A	801	GOL	O1-C1-C2-C3

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	804	GOL	2	0
2	A	801	GOL	1	0
2	B	801	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	241/256 (94%)	-0.11	6 (2%) 57 52	19, 27, 56, 89	0
1	B	239/256 (93%)	0.13	15 (6%) 20 15	17, 28, 68, 117	0
All	All	480/512 (93%)	0.01	21 (4%) 34 28	17, 28, 65, 117	0

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	552	VAL	8.2
1	B	655	GLN	8.1
1	A	654	GLY	5.9
1	B	654	GLY	5.5
1	B	559	ALA	5.0
1	B	550	ARG	4.8
1	B	551	GLY	4.4
1	A	655	GLN	4.2
1	B	556	ASP	4.1
1	A	533	PRO	3.9
1	B	767[A]	HIS	3.2
1	B	534	ARG	3.1
1	B	532	ASN	2.9
1	B	558	ASN	2.7
1	A	530	GLU	2.7
1	B	557	GLY	2.7
1	A	550	ARG	2.4
1	A	770	LEU	2.3
1	B	531	LEU	2.3
1	B	769	HIS	2.2
1	B	554	SER	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	GOL	B	804	6/6	0.81	0.21	59,63,67,67	0
2	GOL	B	802	6/6	0.88	0.18	32,49,50,51	0
2	GOL	B	803	6/6	0.90	0.16	31,46,53,66	0
2	GOL	A	801	6/6	0.94	0.12	23,29,31,32	0
3	FMT	A	802	3/3	0.94	0.10	28,28,38,52	0
2	GOL	B	801	6/6	0.95	0.08	25,33,37,38	0

6.5 Other polymers [i](#)

There are no such residues in this entry.