

wwPDB X-ray Structure Validation Summary Report (i)

Jun 24, 2024 – 10:59 AM EDT

PDB ID : 6TQJ

Title: Crystal structure of the c14 ring of the F1FO ATP synthase from spinach

chloroplast

Authors: Kovalev, K.; Gushchin, I.; Vlasov, A.; Round, E.; Polovinkin, V.; Gordeliy, V.

Deposited on : 2019-12-16

Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 2022.3.0, CSD as 543be (2022)

Xtriage (Phenix) : 1.20.1

EDS : 2.37.1

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.37.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries,\ resolution\ range(\mathring{A})}) \end{array}$
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	Λ	81	2%	
1	A	91	95%	• •
1	В	81	93%	6% •
1	С	81	91%	7% •
1	D	81	91%	5% •
1	Е	81	90%	9% •

Continued on next page...



Continued from previous page...

Mol	Chain	Length	Quality of chain	
1	F	81	89%	10% •
1	G	81	91%	6% •
1	Н	81	91%	7% •
1	I	81	91%	5% •
1	J	81	91%	7% •
1	K	81	95%	
1	L	81	94%	5% •
1	M	81	91%	6% •
1	N	81	94%	6%



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 8459 atoms, of which 0 are hydrogens and 0 are deuteriums.


In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called ATP synthase subunit c, chloroplastic.

Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf	Trace
1	A	81	Total	С	N	О	S	0	1	0
1	A	01	560	365	92	102	1	U	1	
1	В	81	Total	С	N	О	S	0	2	0
1	D	01	568	369	94	104	1	U	2	
1	С	81	Total	С	N	O	S	0	2	0
1		01	569	369	94	105	1	U	2	
1	D	81	Total	С	N	O	S	0	2	0
1	D	01	569	369	94	105	1	U	2	
1	Е	81	Total	С	N	O	\mathbf{S}	0	2	0
1	ш	01	568	369	94	104	1	U	2	0
1	F	81	Total	С	N	O	\mathbf{S}	0	2	0
1	I.	01	568	369	94	104	1	U	<i>Z</i>	
1	G	81	Total	С	N	O	S	0	2	0
1	G	01	568	369	94	104	1			
1	Н	81	Total	С	N	О	S	0	2	0
1	11	01	568	369	94	104	1	0	Δ	0
1	I	81	Total	С	N	О	S	0	2	0
1	1	01	569	369	94	105	1	U	2	
1	J	81	Total	С	N	О	S	0	1	0
1		01	561	365	92	103	1	U	1	
1	K	81	Total	С	N	O	S	0	2	0
1	11	01	566	368	93	104	1	U	2	
1	L	81	Total	С	N	O	S	0	2	0
1	П	01	568	369	94	104	1	U	2	
1	1 M	81	Total	С	N	О	S	0	2	0
	1V1	01	568	369	94	104	1	U		
1	N	81	Total	С	N	О	S	0	2	0
1	11	01	568	369	94	104	1	U	<u> </u>	U

• Molecule 2 is EICOSANE (three-letter code: LFA) (formula: $C_{20}H_{42}$).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C 14 14	0	0
2	В	1	Total C 14 14	0	0
2	С	1	Total C 7 7	0	0
2	С	1	Total C 7 7	0	0
2	С	1	Total C 14 14	0	0
2	D	1	Total C 14 14	0	0
2	Е	1	Total C 14 14	0	0
2	F	1	Total C 14 14	0	0
2	G	1	Total C 14 14	0	0
2	G	1	Total C 7 7	0	0
2	G	1	Total C 14 14	0	0
2	Н	1	Total C 10 10	0	0
2	Н	1	Total C 16 16	0	0
2	Н	1	Total C 8 8	0	0
			C_{α}	ntinued on r	


Continued on next page...



 $Continued\ from\ previous\ page...$

Mol		Residues		ZeroOcc	AltConf
2	Н	1	Total C 14 14	0	0
2	I	1	Total C 16 16	0	0
2	I	1	Total C 14 14	0	0
2	J	1	Total C 14 14	0	0
2	L	1	Total C 14 14	0	0
2	L	1	Total C 14 14	0	0
2	M	1	Total C 8 8	0	0
2	M	1	Total C 10 10	0	0
2	M	1	Total C 14 14	0	0
2	N	1	Total C 8 8	0	0
2	N	1	Total C 8 8	0	0
2	N	1	Total C 14 14	0	0

• Molecule 3 is (2R)-2,3-dihydroxypropyl (9Z)-octadec-9-enoate (three-letter code: OLC) (formula: $C_{21}H_{40}O_4$).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	Н	1	Total C O 13 9 4	0	0

• Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	13	Total O 13 13	0	0
4	В	13	Total O 13 13	0	0
4	С	13	Total O 13 13	0	0
4	D	13	Total O 13 13	0	0
4	Е	15	Total O 15 15	0	0
4	F	14	Total O 14 14	0	0
4	G	13	Total O 13 13	0	0
4	Н	17	Total O 17 17	0	0
4	I	16	Total O 16 16	0	0
4	J	10	Total O 10 10	0	0
4	K	13	Total O 13 13	0	0
4	L	14	Total O 14 14	0	0
4	M	15	Total O 15 15	0	0
4	N	14	Total O 14 14	0	0



3 Residue-property plots (i)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: ATP synthase subunit c, chloroplastic



• Molecule 1: ATP synthase subunit c, chloroplastic







• Molecule 1: ATP synthase subunit c, chloroplastic

Chain M:

91%

6% •

Molecule 1: ATP synthase subunit c, chloroplastic

Chain N:

94%

6%





4 Data and refinement statistics (i)

Property	Value	Source
Space group	I 1 2 1	Depositor
Cell constants	93.14Å 96.34Å 158.68Å	Depositor
a, b, c, α , β , γ	90.00° 106.72° 90.00°	Depositor
Resolution (Å)	20.00 - 2.30	Depositor
rtesolution (A)	48.17 - 2.30	EDS
% Data completeness	96.3 (20.00-2.30)	Depositor
(in resolution range)	96.5 (48.17-2.30)	EDS
R_{merge}	0.19	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.51 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.8.0257	Depositor
R, R_{free}	0.204 , 0.237	Depositor
It, It free	0.216 , 0.248	DCC
R_{free} test set	2851 reflections (4.94%)	wwPDB-VP
Wilson B-factor (\mathring{A}^2)	29.1	Xtriage
Anisotropy	0.320	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.34, 46.0	EDS
L-test for twinning ²	$ < L >=0.53, < L^2>=0.37$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8459	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 9.04% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: OLC, LFA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles	
MIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.66	0/570	0.64	0/777
1	В	0.66	0/578	0.65	0/789
1	С	0.66	0/579	0.65	0/789
1	D	0.66	0/579	0.65	0/789
1	Е	0.66	0/578	0.66	0/789
1	F	0.66	0/578	0.64	0/789
1	G	0.66	0/578	0.65	0/789
1	Н	0.66	0/578	0.64	0/789
1	I	0.66	0/579	0.64	0/789
1	J	0.65	0/571	0.65	0/777
1	K	0.66	0/579	0.65	0/789
1	L	0.67	0/578	0.65	0/789
1	M	0.67	0/578	0.66	0/789
1	N	0.67	0/578	0.64	0/789
All	All	0.66	0/8081	0.65	0/11022

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	560	0	594	3	0
1	В	568	0	600	8	0
1	С	569	0	600	8	0
1	D	569	0	600	11	0
1	Е	568	0	600	11	0
1	F	568	0	600	13	0
1	G	568	0	600	9	0
1	Н	568	0	600	10	0
1	I	569	0	600	11	0
1	J	561	0	594	4	0
1	K	566	0	600	4	0
1	L	568	0	600	5	0
1	M	568	0	600	8	0
1	N	568	0	600	5	0
2	A	14	0	27	0	0
2	В	14	0	27	1	0
2	С	28	0	47	2	0
2	D	14	0	27	1	0
2	Е	14	0	27	1	0
2	F	14	0	27	1	0
2	G	35	0	64	1	0
2	Н	48	0	92	5	0
2	I	30	0	58	3	0
2	J	14	0	27	2	0
2	L	28	0	54	2	0
2	M	32	0	61	2	0
2	N	30	0	57	2	0
3	Н	13	0	15	0	0
4	A	13	0	0	1	0
4	В	13	0	0	1	0
4	С	13	0	0	1	0
4	D	13	0	0	1	0
4	Е	15	0	0	2	0
4	F	14	0	0	1	0
4	G	13	0	0	0	0
4	Н	17	0	0	2	0
4	I	16	0	0	1	0
4	J	10	0	0	2	0
4	K	13	0	0	0	0
4	L	14	0	0	0	0
4	M	15	0	0	1	0
4	N	14	0	0	1	0
All	All	8459	0	8998	109	0



The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 109 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:D:55:LEU:HD21	1:E:54:LEU:HD21	1.14	1.05
1:D:55:LEU:HD21	1:E:54:LEU:CD2	2.05	0.83
1:L:2[B]:ASN:N	1:L:3:PRO:HD2	2.00	0.77
1:B:2[B]:ASN:N	1:B:3:PRO:HD2	2.01	0.76
1:F:2[B]:ASN:N	1:F:3:PRO:HD2	2.01	0.76

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	$80/81\ (99\%)$	79 (99%)	1 (1%)	0	100	100
1	В	81/81 (100%)	80 (99%)	1 (1%)	0	100	100
1	С	81/81 (100%)	80 (99%)	1 (1%)	0	100	100
1	D	81/81 (100%)	80 (99%)	1 (1%)	0	100	100
1	E	81/81 (100%)	80 (99%)	1 (1%)	0	100	100
1	F	81/81 (100%)	80 (99%)	1 (1%)	0	100	100
1	G	81/81 (100%)	80 (99%)	1 (1%)	0	100	100
1	Н	81/81 (100%)	80 (99%)	1 (1%)	0	100	100
1	I	81/81 (100%)	80 (99%)	1 (1%)	0	100	100
1	J	80/81 (99%)	79 (99%)	1 (1%)	0	100	100
1	K	81/81 (100%)	80 (99%)	1 (1%)	0	100	100
1	L	81/81 (100%)	80 (99%)	1 (1%)	0	100	100

Continued on next page...



Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles		
1	M	81/81 (100%)	80 (99%)	1 (1%)	0	100	100	
1	N	81/81 (100%)	80 (99%)	1 (1%)	0	100	100	
All	All	1132/1134 (100%)	1118 (99%)	14 (1%)	0	100	100	

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	A	52/53~(98%)	50 (96%)	2 (4%)	33 47		
1	В	53/53 (100%)	49 (92%)	4 (8%)	13 17		
1	С	53/53 (100%)	49 (92%)	4 (8%)	13 17		
1	D	53/53 (100%)	49 (92%)	4 (8%)	13 17		
1	E	53/53 (100%)	49 (92%)	4 (8%)	13 17		
1	F	53/53 (100%)	51 (96%)	2 (4%)	33 47		
1	G	53/53 (100%)	49 (92%)	4 (8%)	13 17		
1	Н	53/53 (100%)	49 (92%)	4 (8%)	13 17		
1	I	53/53 (100%)	49 (92%)	4 (8%)	13 17		
1	J	52/53 (98%)	49 (94%)	3 (6%)	20 27		
1	K	53/53 (100%)	50 (94%)	3 (6%)	20 28		
1	L	53/53 (100%)	49 (92%)	4 (8%)	13 17		
1	M	53/53 (100%)	48 (91%)	5 (9%)	8 10		
1	N	53/53 (100%)	51 (96%)	2 (4%)	33 47		
All	All	740/742 (100%)	691 (93%)	49 (7%)	21 22		

5 of 49 residues with a non-rotameric sidechain are listed below:

\mathbf{Mol}	Chain	Res	Type
1	I	2[A]	ASN

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type
1	K	2[A]	ASN
1	I	2[B]	ASN
1	J	15	LEU
1	K	15	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	I	42	GLN
1	M	42	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

27 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Pog	Link	Bond lengths			Bond angles		
IVIOI	туре	Chain	rtes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	LFA	Н	103	-	7,7,19	0.11	0	6,6,18	0.11	0
2	LFA	F	101	-	13,13,19	0.08	0	12,12,18	0.07	0
2	LFA	Н	104	-	13,13,19	0.07	0	12,12,18	0.07	0



Mal	Trens	Chain	Dag	Link	Во	nd leng	ths	В	ond ang	gles
Mol	Type	Chain	Res	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	LFA	С	102	-	6,6,19	0.10	0	5,5,18	0.12	0
2	LFA	С	101	-	6,6,19	0.09	0	5,5,18	0.17	0
2	LFA	I	102	-	13,13,19	0.08	0	12,12,18	0.08	0
2	LFA	N	102	-	7,7,19	0.09	0	6,6,18	0.10	0
3	OLC	Н	105	-	12,12,24	1.30	1 (8%)	13,13,25	1.16	2 (15%)
2	LFA	I	101	-	15,15,19	0.09	0	14,14,18	0.05	0
2	LFA	M	101	-	7,7,19	0.09	0	6,6,18	0.10	0
2	LFA	Н	101	-	9,9,19	0.09	0	8,8,18	0.08	0
2	LFA	A	101	-	13,13,19	0.08	0	12,12,18	0.07	0
2	LFA	G	101	-	13,13,19	0.09	0	12,12,18	0.09	0
2	LFA	N	101	-	7,7,19	0.10	0	6,6,18	0.12	0
2	LFA	L	102	-	13,13,19	0.08	0	12,12,18	0.07	0
2	LFA	M	103	-	13,13,19	0.09	0	12,12,18	0.06	0
2	LFA	G	103	-	13,13,19	0.07	0	12,12,18	0.07	0
2	LFA	В	101	-	13,13,19	0.09	0	12,12,18	0.08	0
2	LFA	D	101	-	13,13,19	0.08	0	12,12,18	0.07	0
2	LFA	N	103	-	13,13,19	0.09	0	12,12,18	0.07	0
2	LFA	J	101	-	13,13,19	0.08	0	12,12,18	0.06	0
2	LFA	G	102	-	6,6,19	0.08	0	5,5,18	0.12	0
2	LFA	Н	102	-	15,15,19	0.09	0	14,14,18	0.06	0
2	LFA	С	103	-	13,13,19	0.07	0	12,12,18	0.07	0
2	LFA	Е	101	-	13,13,19	0.07	0	12,12,18	0.07	0
2	LFA	M	102	-	9,9,19	0.10	0	8,8,18	0.09	0
2	LFA	L	101	-	13,13,19	0.08	0	12,12,18	0.06	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	LFA	Н	103	-	-	1/5/5/17	-
2	LFA	F	101	-	-	7/11/11/17	-
2	LFA	Н	104	-	-	4/11/11/17	-
2	LFA	С	102	-	-	0/4/4/17	-
2	LFA	С	101	-	-	2/4/4/17	-
2	LFA	I	102	-	-	8/11/11/17	-
2	LFA	N	102	-	-	1/5/5/17	-
3	OLC	Н	105	-	-	6/12/12/24	-
2	LFA	I	101	-	-	5/13/13/17	-
2	LFA	M	101	-	-	2/5/5/17	-

Continued on next page...



Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	LFA	Н	101	-	-	3/7/7/17	-
2	LFA	A	101	-	-	6/11/11/17	-
2	LFA	G	101	-	-	9/11/11/17	-
2	LFA	N	101	-	-	2/5/5/17	-
2	LFA	L	102	-	-	6/11/11/17	-
2	LFA	M	103	-	-	7/11/11/17	-
2	LFA	G	103	-	-	6/11/11/17	-
2	LFA	В	101	-	-	5/11/11/17	-
2	LFA	D	101	-	-	9/11/11/17	-
2	LFA	N	103	-	-	7/11/11/17	-
2	LFA	J	101	-	-	5/11/11/17	-
2	LFA	G	102	-	-	3/4/4/17	-
2	LFA	Н	102	-	-	6/13/13/17	-
2	LFA	С	103	-		8/11/11/17	-
2	LFA	Е	101	-	-	8/11/11/17	-
2	LFA	M	102	_		3/7/7/17	-
2	LFA	L	101	_	_	5/11/11/17	_

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\text{\AA})$	$Ideal(\AA)$
3	Н	105	OLC	O20-C1	4.29	1.45	1.33

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$Ideal(^{o})$
3	Н	105	OLC	O20-C1-C2	3.05	121.14	111.83
3	Н	105	OLC	O20-C1-O19	-2.23	118.04	123.63

There are no chirality outliers.

5 of 134 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	Н	105	OLC	O20-C21-C22-C24
3	Н	105	OLC	C2-C1-O20-C21
3	Н	105	OLC	O20-C21-C22-O23
2	В	101	LFA	C10-C11-C12-C13
3	Н	105	OLC	O19-C1-O20-C21




There are no ring outliers.

14 monomers are involved in 14 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	F	101	LFA	1	0
2	Н	104	LFA	4	0
2	I	102	LFA	3	0
2	L	102	LFA	1	0
2	M	103	LFA	2	0
2	G	103	LFA	1	0
2	В	101	LFA	1	0
2	D	101	LFA	1	0
2	N	103	LFA	2	0
2	J	101	LFA	2	0
2	Н	102	LFA	1	0
2	С	103	LFA	2	0
2	Е	101	LFA	1	0
2	L	101	LFA	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





Ligand LFA F 101								
	^							
Bond lengths	Bond angles							
~~~~								
Torsions	Rings							
Ligand Li	FA H 104							
~~~~	~~~~							
Bond lengths	Bond angles							
~~~~								
Torsions	Rings							
Ligand L	FA C 101							
/ / / / / / / / / /	✓							
Bond lengths	Bond angles							
Torsions	Rings							
Ligand LFA I 102								
^	^							
Bond lengths	Bond angles							
~~~~								
Torsions	Rings							
Ligand L	FA N 102							
Bond lengths	Bond angles							
Torsions	Rings							







Ligand LFA A 101								
Bond lengths	Bond angles							
Dond lengths	Don't angles							
Torsions	Rings							
Ligand LFA G 101								
<b>~~~</b>	<b>~~~</b>							
Bond lengths	Bond angles							
~~~~								
Torsions	Rings							
Ligand Li	FA N 101							
/								
Bond lengths	Bond angles							
Torsions	Rings							
Ligand L	FA L 102							
D 11 41								
Bond lengths	Bond angles							
Torsions	Rings							
Ligand Li	FA M 103							
Bond lengths	Bond angles							
Dond lengths	Dong angles							
Torsions	Rings							
Ligand Li	Ligand LFA G 103							
Bond lengths	Bond angles							
Torsions	Rings							



Ligand LFA B 101							
	^						
Bond lengths	Bond angles						
~~~~							
Torsions	Rings						
Ligand LFA D 101							
^	^						
Bond lengths	Bond angles						
~~~~							
Torsions	Rings						
Ligand L	FA N 103						
~~~~	~~~~						
Bond lengths	Bond angles						
	D.						
Torsions	Rings						
Ligand L	FA J 101						
~~~~	<b>~~~~</b>						
Bond lengths	Bond angles						
Torsions	Rings						
Ligand L	FA G 102						
Bond lengths	Bond angles						
Torsions	Rings						
	-						
Ligand L	FA H 102						
<b>\\\\\</b>	<b>~~~~</b> \						
Bond lengths	Bond angles						
~~~~							
1							
Torsions	Rings						
Ligand L	FA C 103						
	~~~~						
Bond lengths	Bond angles						
/ / ////							
Torsions	Rings						



Ligand LFA E 101							
^	//////						
Bond lengths	Bond angles						
~~~~							
Torsions	Rings						
Ligand I	FA M 102						
~~~	~~~						
Bond lengths	Bond angles						
Torsions	Rings						
Ligand 1	LFA L 101						
	<b>~~~~</b>						
Bond lengths	Bond angles						
<b>^</b>							
Torsions	Rings						

### 5.7 Other polymers (i)

There are no such residues in this entry.

### 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



### 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed <rsr< th=""><th>#RSRZ</th><th>&gt;2</th><th>$\mathbf{OWAB}(\mathbf{\mathring{A}}^2)$</th><th>Q &lt; 0.9</th></rsr<>		#RSRZ	>2	$\mathbf{OWAB}(\mathbf{\mathring{A}}^2)$	Q < 0.9
1	A	81/81~(100%)	-0.14	2 (2%) 57	64	19, 28, 47, 74	0
1	В	81/81 (100%)	-0.14	3 (3%) 41	48	18, 28, 48, 79	0
1	С	81/81 (100%)	-0.09	2 (2%) 57	64	19, 28, 48, 86	0
1	D	81/81 (100%)	-0.17	2 (2%) 57	64	20, 28, 46, 78	0
1	Е	81/81 (100%)	-0.17	2 (2%) 57	64	19, 26, 45, 67	0
1	F	81/81 (100%)	-0.16	2 (2%) 57	64	18, 27, 40, 81	0
1	G	81/81 (100%)	-0.18	2 (2%) 57	64	15, 28, 48, 75	0
1	Н	81/81 (100%)	-0.14	2 (2%) 57	64	19, 27, 44, 71	0
1	I	81/81 (100%)	-0.17	2 (2%) 57	64	18, 26, 46, 79	0
1	J	81/81 (100%)	-0.18	3 (3%) 41	48	20, 28, 42, 73	0
1	K	81/81 (100%)	0.00	4 (4%) 29	36	20, 29, 46, 71	0
1	L	81/81 (100%)	-0.18	2 (2%) 57	64	19, 29, 50, 90	0
1	M	81/81 (100%)	-0.23	2 (2%) 57	64	18, 28, 45, 83	0
1	N	81/81 (100%)	-0.10	1 (1%) 79	83	18, 26, 45, 76	0
All	All	1134/1134 (100%)	-0.15	31 (2%) 54	62	15, 28, 51, 90	0

The worst 5 of 31 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	K	1	MET	6.8
1	С	1	MET	6.7
1	D	1	MET	6.7
1	M	1	MET	5.9
1	J	1	MET	5.5



### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.


### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}({ ext{\AA}}^2)$	Q < 0.9
2	LFA	Ε	101	14/20	0.49	0.34	76,86,91,97	0
2	LFA	Н	104	14/20	0.53	0.36	72,91,109,109	0
2	LFA	В	101	14/20	0.56	0.26	57,77,87,87	0
2	LFA	A	101	14/20	0.60	0.28	63,78,84,87	0
2	LFA	L	102	14/20	0.60	0.26	52,78,87,89	0
2	LFA	G	103	14/20	0.63	0.27	63,85,104,105	0
2	LFA	J	101	14/20	0.65	0.35	81,99,110,110	0
2	LFA	С	103	14/20	0.67	0.24	61,77,82,83	0
2	LFA	I	102	14/20	0.67	0.23	54,74,86,87	0
2	LFA	N	103	14/20	0.67	0.24	72,81,87,89	0
2	LFA	L	101	14/20	0.68	0.24	71,74,92,93	0
2	LFA	D	101	14/20	0.71	0.26	64,72,83,85	0
2	LFA	Н	103	8/20	0.71	0.29	47,60,72,74	0
2	LFA	F	101	14/20	0.72	0.28	71,92,101,102	0
2	LFA	M	103	14/20	0.73	0.24	64,81,88,88	0
3	OLC	Н	105	13/25	0.75	0.24	47,58,63,63	0
2	LFA	Н	101	10/20	0.77	0.19	45,54,59,60	0
2	LFA	G	101	14/20	0.77	0.20	45,60,67,67	0
2	LFA	С	102	7/20	0.79	0.36	63,66,74,74	0
2	LFA	M	102	10/20	0.80	0.29	42,51,60,63	0
2	LFA	M	101	8/20	0.81	0.24	51,63,69,69	0
2	LFA	С	101	7/20	0.84	0.16	41,45,49,50	0
2	LFA	I	101	16/20	0.85	0.19	39,46,56,57	0
2	LFA	Н	102	16/20	0.85	0.21	36,53,74,78	0
2	LFA	N	102	8/20	0.90	0.22	39,45,60,66	0
2	LFA	N	101	8/20	0.92	0.28	39,41,45,46	0
2	LFA	G	102	7/20	0.94	0.15	38,48,56,60	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







### Electron density around LFA H 104:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 0.7 rmsd) in gray  $\mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)










### Electron density around LFA B 101:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 0.7 rmsd) in gray  $\mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)












### Electron density around LFA A 101:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 0.7 rmsd) in gray  $\mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)










### Electron density around LFA L 102:

 $2 \mathrm{mF}_o\text{-DF}_c$  (at 0.7 rmsd) in gray  $\mathrm{mF}_o\text{-DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)












# Electron density around LFA G 103: 2mF_o-DF_c (at 0.7 rmsd) in gray mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)

## Electron density around LFA J 101:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$  (at 0.7 rmsd) in gray  ${\rm mF}_o\text{-}{\rm DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)










### Electron density around LFA C 103:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 0.7 rmsd) in gray  $\mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)










### Electron density around LFA I 102:


 $2 \mathrm{mF}_o\text{-DF}_c$  (at 0.7 rmsd) in gray  $\mathrm{mF}_o\text{-DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)














 $2 {\rm mF}_o\text{-}{\rm DF}_c$  (at 0.7 rmsd) in gray  ${\rm mF}_o\text{-}{\rm DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)













# Electron density around LFA D 101:












### Electron density around LFA H 103:

 $2 {
m mF}_o {
m -DF}_c$  (at 0.7 rmsd) in gray  ${
m mF}_o {
m -DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)












### Electron density around LFA F 101:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 0.7 rmsd) in gray  $\mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)










### Electron density around LFA M 103:


 $2 {\rm mF}_o\text{-}{\rm DF}_c$  (at 0.7 rmsd) in gray  ${\rm mF}_o\text{-}{\rm DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)



















### Electron density around LFA M 102:

 $2 {
m mF}_o {
m -DF}_c$  (at 0.7 rmsd) in gray  ${
m mF}_o {
m -DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)












### Electron density around LFA M 101:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 0.7 rmsd) in gray  $\mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)










### Electron density around LFA C 101:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 0.7 rmsd) in gray  $\mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)











# Electron density around LFA I 101: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${ m mF}_o{ m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive) Electron density around LFA H 102: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $mF_o$ -DF_c (at 3 rmsd) in purple (negative) and green (positive)




### Electron density around LFA N 102:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 0.7 rmsd) in gray  $\mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)










### Electron density around LFA N 101:


 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 0.7 rmsd) in gray  $\mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)











### 6.5 Other polymers (i)

There are no such residues in this entry.

