

# wwPDB X-ray Structure Validation Summary Report (i)

Aug 2, 2022 – 01:33 pm BST

PDB ID : 6TXN

Title : Crystal structure of thermotoga maritima Ferritin in apo form Authors : Wilk, P.; Grudnik, P.; Kumar, M.; Heddle, J.; Chakraborti, S.

Deposited on : 2020-01-14

Resolution : 2.01 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul: 1.8.4, CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.29

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0267

CCP4 : 7.1.010 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.29

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$ 

The reported resolution of this entry is 2.01 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries,\ resolution\ range(\mathring{\rm A})}) \end{array}$
$R_{free}$	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	164	90%	9% •
1	В	164	87%	12% •
1	С	164	88%	11% •
1	D	164	90%	10%
1	E	164	94%	6%



 $Continued\ from\ previous\ page...$ 

Mol	Chain	Length	Quality of chain	
1	F	164	8%	11%
1	G	164	92%	7% •
1	Н	164	6% 88%	12%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	SO4	С	203	-	-	-	X
3	SO4	С	204	-	-	-	X



## 2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 22696 atoms, of which 10836 are hydrogens and 0 are deuteriums.


In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Ferritin.

Mol	Chain	Residues			Atom	ıs			ZeroOcc	AltConf	Trace
1	A	164	Total	С	Н	N	О	S	0	1	0
1	Λ	104	2702	879	1329	224	264	6	U	1	
1	В	164	Total	С	Н	N	О	S	0	4	0
1	D	104	2728	887	1341	226	268	6	U	4	
1	С	164	Total	С	Н	N	О	S	0	1	0
1		104	2693	878	1320	223	266	6	U	1	U
1	D	164	Total	С	Н	N	O	S	0	3	0
1	D	104	2724	886	1343	225	264	6	U		U
1	E	164	Total	С	Н	N	O	S	0	2	0
1	ш	104	2712	881	1334	226	265	6	U		
1	F	164	Total	С	Н	N	О	S	0	1	0
1	I.	104	2663	868	1304	222	263	6	U	1	0
1	G	164	Total	С	Η	N	О	S	0	0	0
1	G	104	2668	871	1305	222	264	6	U	<u> </u>	
1	Н	164	Total	С	Н	N	О	S	0	4	0
1	11	104	2757	893	1362	231	265	6	U	1	

• Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>) (labeled as "Ligand of Interest" by depositor).






Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C H O 9 3 3 3	0	0
2	В	1	Total C H O 9 3 3 3	0	0
2	В	1	Total C H O 9 3 3 3	0	0
2	С	1	Total C H O 9 3 3 3	0	0
2	С	1	Total C H O 9 3 3 3	0	0
2	D	1	Total C H O 9 3 3 3	0	0
2	D	1	Total C H O 9 3 3 3	0	0
2	Е	1	Total C H O 9 3 3 3	0	0
2	G	1	Total C H O 9 3 3 3	0	0
2	Н	1	Total C H O 9 3 3 3	0	0

 $\bullet$  Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S) (labeled as "Ligand of Interest" by depositor).






Mol	Chain	Residues	Ato	oms		ZeroOcc	AltConf
3	A	1	Total	О	S	0	0
3	A	1	5	4	1	0	U
3	В	1	Total	О	S	0	0
	Ъ	1	5	4	1	U	U
3	В	1	Total	Ο	S	0	0
	D	1	5	4	1	0	U
3	В	1	Total	Ο	S	0	0
	D	1	5	4	1	0	Ů
3	С	1	Total	Ο	S	0	0
		1	5	4	1	O	Ŭ
3	С	1	Total	Ο	S	0	0
		1	5	4	1	Ü	Ü
3	D	1	Total	O	S	0	0
			5	4	1	Ŭ	Ŭ
3	Е	1	Total	O	S	0	0
	_	_	5	4	1		<u> </u>
3	Е	1	Total	O	S	0	0
			5	4	1		_
3	F	1	Total	O	S	0	0
			5	4	1		-
3	F	1	Total	O	S	0	0
			5	4	1		
3	Н	1	Total	O	S	0	0
			5	4	1		_

 $\bullet$  Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Mg 1 1	0	0
4	В	1	Total Mg 1 1	0	0
4	С	1	Total Mg 1 1	0	0
4	D	1	Total Mg 1 1	0	0
4	E	1	Total Mg 1 1	0	0
4	F	1	Total Mg 1 1	0	0
4	G	1	Total Mg 1 1	0	0
4	Н	1	Total Mg 1 1	0	0

 $\bullet$  Molecule 5 is EICOSANE (three-letter code: LFA) (formula:  $C_{20}H_{42})$  (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	В	1	Total C H 62 20 42	0	0
5	D	1	Total C H 62 20 42	0	0
5	Е	1	Total C H 62 20 42	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	
5	G	1	Total 62	C 20	H 42	0	0

#### • Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	115	Total O 115 115	0	0
6	В	105	Total O 105 105	0	0
6	С	76	Total O 76 76	0	0
6	D	81	Total O 81 81	0	0
6	Е	91	Total O 91 91	0	0
6	F	66	Total O 66 66	0	0
6	G	63	Total O 63 63	0	0
6	Н	46	Total O 46 46	0	0




## 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.









## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants	176.02Å 176.02Å 356.27Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $120.00^{\circ}$	Depositor
Resolution (Å)	48.28 - 2.01	Depositor
resolution (A)	48.28 - 2.01	EDS
% Data completeness	99.8 (48.28-2.01)	Depositor
(in resolution range)	93.2 (48.28-2.01)	EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	0.73 (at 2.01Å)	Xtriage
Refinement program	PHENIX 1.14_3260	Depositor
P. P.	0.193 , 0.221	Depositor
$R, R_{free}$	0.195 , $0.223$	DCC
$R_{free}$ test set	2000 reflections (1.43%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	40.9	Xtriage
Anisotropy	0.114	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	(Not available), (Not available)	EDS
L-test for twinning <sup>2</sup>	$  <  L  > = 0.48, < L^2 > = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	22696	wwPDB-VP
Average B, all atoms $(Å^2)$	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 2.68% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $<L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

### 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MG, LFA, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Во	Bond lengths		ond angles
IVIOI	Chain	RMSZ	# Z  > 5	RMSZ	# Z >5
1	A	0.79	1/1405 (0.1%)	0.82	2/1887 (0.1%)
1	В	0.78	0/1438	0.81	4/1933~(0.2%)
1	С	0.73	0/1405	0.77	3/1888 (0.2%)
1	D	0.72	0/1419	0.76	6/1905~(0.3%)
1	Е	0.72	0/1413	0.73	0/1898
1	F	0.77	0/1390	0.79	2/1869~(0.1%)
1	G	0.73	$2/1392 \ (0.1\%)$	0.66	0/1872
1	Н	0.60	0/1436	0.64	0/1927
All	All	0.73	3/11298 (0.0%)	0.75	17/15179 (0.1%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$\operatorname{Observed}(\operatorname{\AA})$	$\operatorname{Ideal}(\text{\AA})$
1	G	52	GLU	CD-OE1	-9.11	1.15	1.25
1	G	51	GLU	CD-OE2	-6.56	1.18	1.25
1	A	65	GLU	CD-OE1	-5.72	1.19	1.25

The worst 5 of 17 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$\mathbf{Observed}(^{o})$	$\operatorname{Ideal}({}^{o})$
1	A	69	ARG	NE-CZ-NH1	10.40	125.50	120.30
1	С	69	ARG	NE-CZ-NH2	-9.91	115.35	120.30
1	С	69	ARG	NE-CZ-NH1	8.25	124.42	120.30
1	A	69	ARG	NE-CZ-NH2	-7.89	116.36	120.30
1	С	18	ARG	NE-CZ-NH2	-6.93	116.84	120.30

There are no chirality outliers.

There are no planarity outliers.



#### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1373	1329	1332	11	0
1	В	1387	1341	1334	14	2
1	С	1373	1320	1325	13	0
1	D	1381	1343	1347	15	0
1	Ε	1378	1334	1337	9	0
1	F	1359	1304	1308	12	0
1	G	1363	1305	1308	12	0
1	Н	1395	1362	1366	16	0
2	A	6	3	8	0	0
2	В	12	6	15	0	0
2	С	12	6	16	2	0
2	D	12	6	16	1	0
2	Ε	6	3	7	0	0
2	G	6	3	8	1	0
2	Н	6	3	8	0	0
3	A	5	0	0	0	0
3	В	15	0	0	2	0
3	С	10	0	0	2	0
3	D	5	0	0	1	0
3	Ε	10	0	0	0	0
3	F	10	0	0	0	0
3	Н	5	0	0	0	0
4	A	1	0	0	0	0
4	В	1	0	0	0	0
4	С	1	0	0	0	0
4	D	1	0	0	0	0
4	Ε	1	0	0	0	0
4	F	1	0	0	0	0
4	G	1	0	0	0	0
4	Н	1	0	0	0	0
5	В	20	42	42	0	0
5	D	20	42	42	1	0
5	Е	20	42	42	0	0
5	G	20	42	42	4	0
6	A	115	0	0	4	0
6	В	105	0	0	7	0
6	С	76	0	0	2	0



Continued	trom	mmoninonic	maaa
COHABABACA		DIEUIUU	DUIUE
0 0 1000100000			

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	D	81	0	0	6	0
6	Ε	91	0	0	3	0
6	F	66	0	0	3	0
6	G	63	0	0	2	0
6	Н	46	0	0	8	0
All	All	11860	10836	10903	100	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 100 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
3:B:201:SO4:O1	6:B:301:HOH:O	1.53	1.27
1:C:50:GLN:NE2	6:C:301:HOH:O	1.91	1.01
1:H:18[B]:ARG:NH2	6:H:303:HOH:O	2.03	0.91
1:F:18:ARG:NH2	6:F:301:HOH:O	2.04	0.91
1:H:97:GLU:OE1	6:H:302:HOH:O	1.94	0.85

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:B:108:GLU:OE1	1:B:114:LYS:NZ[2_665]	2.01	0.19
1:B:108:GLU:OE1	1:B:114:LYS:HZ2[2_665]	1.45	0.15

### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	163/164 (99%)	163 (100%)	0	0	100 100



Continued from previous page.					
	C'	ontinued	from	previous	page

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	В	166/164 (101%)	165 (99%)	1 (1%)	0	100	100
1	С	163/164~(99%)	163 (100%)	0	0	100	100
1	D	165/164~(101%)	163 (99%)	2 (1%)	0	100	100
1	E	$164/164 \; (100\%)$	163 (99%)	1 (1%)	0	100	100
1	F	163/164~(99%)	161 (99%)	2 (1%)	0	100	100
1	G	162/164~(99%)	159 (98%)	3 (2%)	0	100	100
1	Н	166/164 (101%)	164 (99%)	2 (1%)	0	100	100
All	All	1312/1312 (100%)	1301 (99%)	11 (1%)	0	100	100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	145/144 (101%)	144 (99%)	1 (1%)	84 88
1	В	148/144 (103%)	147~(99%)	1 (1%)	84 88
1	С	145/144 (101%)	142 (98%)	3 (2%)	53 57
1	D	146/144 (101%)	145 (99%)	1 (1%)	84 88
1	E	146/144 (101%)	146 (100%)	0	100 100
1	F	142/144~(99%)	141 (99%)	1 (1%)	84 88
1	G	143/144 (99%)	143 (100%)	0	100 100
1	Н	148/144 (103%)	147 (99%)	1 (1%)	84 88
All	All	1163/1152 (101%)	1155 (99%)	8 (1%)	84 88

5 of 8 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type	
1	Н	28	SER	
1	F	147	ASN	



Continued from previous page...

Mol	Chain	Res	Type
1	С	58	LYS
1	С	52	GLU
1	D	9	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	ol Chain Res		Type	
1	F	81	ASN	
1	F	147	ASN	
1	G	55	HIS	

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry (i)

Of 34 ligands modelled in this entry, 8 are monoatomic - leaving 26 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bo	ond leng	ths	В	ond ang	cles
MIOI	туре	Chain	nes	LIIIK	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	GOL	С	202	-	5,5,5	1.15	1 (20%)	5,5,5	1.26	1 (20%)
5	LFA	G	202	-	19,19,19	0.36	0	18,18,18	0.45	0



Mol	Trino	Chain	Dag	Link	Во	nd leng	ths	В	ond ang	eles
IVIOI	Type	Chain	Res	Link	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	GOL	A	201	-	5,5,5	1.14	0	5,5,5	0.82	0
2	GOL	Е	202	-	5,5,5	1.19	1 (20%)	5,5,5	0.87	0
5	LFA	В	205	-	19,19,19	0.25	0	18,18,18	0.41	0
3	SO4	Н	201	-	4,4,4	0.11	0	6,6,6	0.18	0
3	SO4	D	204	-	4,4,4	0.14	0	6,6,6	0.18	0
2	GOL	В	202	-	5,5,5	1.45	1 (20%)	5,5,5	1.06	0
3	SO4	F	202	-	4,4,4	0.17	0	6,6,6	0.05	0
3	SO4	В	201	-	4,4,4	0.15	0	6,6,6	0.05	0
2	GOL	G	201	-	5,5,5	1.31	0	5,5,5	1.04	0
3	SO4	A	202	-	4,4,4	0.24	0	6,6,6	0.27	0
5	LFA	D	203	-	19,19,19	0.24	0	18,18,18	0.50	0
2	GOL	Н	202	-	5,5,5	0.69	0	5,5,5	1.03	0
2	GOL	D	201	-	5,5,5	0.83	0	5,5,5	1.05	0
3	SO4	С	204	-	4,4,4	0.14	0	6,6,6	0.23	0
2	GOL	С	201	-	5,5,5	0.79	0	5,5,5	1.02	0
3	SO4	В	206	-	4,4,4	0.19	0	6,6,6	0.16	0
3	SO4	В	203	-	4,4,4	0.14	0	6,6,6	0.05	0
2	GOL	В	204	-	5,5,5	0.74	0	5,5,5	0.88	0
5	LFA	Е	203	-	19,19,19	0.19	0	18,18,18	0.27	0
3	SO4	Е	201	-	4,4,4	0.14	0	6,6,6	0.05	0
3	SO4	F	201	-	4,4,4	0.14	0	6,6,6	0.04	0
2	GOL	D	202	-	5,5,5	0.73	0	5,5,5	1.28	0
3	SO4	Е	204	-	4,4,4	0.14	0	6,6,6	0.08	0
3	SO4	С	203	-	4,4,4	0.68	0	6,6,6	0.13	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	В	204	-	-	4/4/4/4	-
2	GOL	В	202	-	-	4/4/4/4	-
2	GOL	С	202	-	-	2/4/4/4	-
5	LFA	G	202	-	-	8/17/17/17	-
2	GOL	G	201	-	-	3/4/4/4	-
5	LFA	Е	203	-	-	7/17/17/17	-
5	LFA	D	203	-	-	5/17/17/17	-
2	GOL	Н	202	-	-	2/4/4/4	-
2	GOL	D	201	-	-	0/4/4/4	-



Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	201	-	-	2/4/4/4	-
2	GOL	E	202	-	-	2/4/4/4	-
5	LFA	В	205	-	-	3/17/17/17	-
2	GOL	С	201	-	-	4/4/4/4	-
2	GOL	D	202	-	-	4/4/4/4	-

#### All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$\operatorname{Observed}(\text{\AA})$	Ideal(A)
2	В	202	GOL	O2-C2	-2.43	1.36	1.43
2	Е	202	GOL	O2-C2	-2.22	1.36	1.43
2	С	202	GOL	C1-C2	2.20	1.60	1.51

#### All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
2	С	202	GOL	C3-C2-C1	-2.04	103.77	111.70

There are no chirality outliers.

5 of 50 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	В	204	GOL	O1-C1-C2-C3
2	С	202	GOL	O1-C1-C2-C3
2	D	202	GOL	O1-C1-C2-O2
2	D	202	GOL	O1-C1-C2-C3
2	D	202	GOL	C1-C2-C3-O3

There are no ring outliers.

10 monomers are involved in 14 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	С	202	GOL	2	0
5	G	202	LFA	4	0
3	D	204	SO4	1	0
3	В	201	SO4	1	0
2	G	201	GOL	1	0
5	D	203	LFA	1	0
3	С	204	SO4	1	0
3	В	203	SO4	1	0




Continued from previous page...


Mo	ol	Chain	Res	Type	Clashes	Symm-Clashes
2		D	202	GOL	1	0
3		С	203	SO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











































































































### 5.7 Other polymers (i)

There are no such residues in this entry.

### 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



### 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle { m RSRZ} \rangle$	$\# \mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q<0.9
1	A	164/164 (100%)	0.26	1 (0%) 89 88	33, 43, 64, 102	0
1	В	164/164 (100%)	0.01	4 (2%) 59 57	33, 43, 66, 88	0
1	С	164/164 (100%)	0.04	5 (3%) 50 49	36, 47, 69, 91	0
1	D	164/164 (100%)	-0.01	3 (1%) 68 66	37, 47, 66, 95	0
1	Е	164/164 (100%)	0.10	2 (1%) 79 78	35, 47, 65, 99	0
1	F	164/164 (100%)	0.40	13 (7%) 12 11	34, 54, 85, 101	0
1	G	164/164 (100%)	0.11	4 (2%) 59 57	41, 52, 79, 103	0
1	Н	164/164 (100%)	0.21	10 (6%) 21 20	40, 55, 87, 105	0
All	All	1312/1312 (100%)	0.14	42 (3%) 47 46	33, 49, 75, 105	0

The worst 5 of 42 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1	MET	7.4
1	Н	154	PHE	5.3
1	F	154	PHE	4.6
1	F	159	TYR	4.5
1	Н	151	SER	4.3

### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.



### 6.4 Ligands (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}(\mathring{\mathbf{A}}^2)$	Q<0.9
4	MG	Ε	205	1/1	0.38	0.12	63,63,63,63	0
3	SO4	С	203	5/5	0.59	0.59	118,120,153,153	0
3	SO4	F	201	5/5	0.66	0.29	82,107,124,167	0
2	GOL	A	201	6/6	0.69	0.17	67,81,97,97	0
3	SO4	В	201	5/5	0.71	0.33	109,117,178,199	0
2	GOL	С	201	6/6	0.72	0.16	70,80,96,96	0
3	SO4	D	204	5/5	0.75	0.25	77,99,115,121	0
2	GOL	D	202	6/6	0.76	0.19	67,83,87,90	0
2	GOL	В	204	6/6	0.76	0.28	66,80,96,96	0
2	GOL	D	201	6/6	0.76	0.32	72,86,96,98	0
3	SO4	С	204	5/5	0.77	0.43	98,98,102,117	0
2	GOL	Н	202	6/6	0.78	0.28	81,91,117,117	0
4	MG	G	203	1/1	0.79	0.06	58,58,58,58	0
5	LFA	G	202	20/20	0.79	0.30	46,69,95,105	0
3	SO4	В	206	5/5	0.80	0.41	76,87,95,110	0
2	GOL	С	202	6/6	0.82	0.24	57,74,79,89	0
4	MG	F	203	1/1	0.83	0.16	57,57,57,57	0
2	GOL	G	201	6/6	0.83	0.18	54,71,89,89	0
3	SO4	Ε	204	5/5	0.83	0.23	107,111,144,214	0
3	SO4	Ε	201	5/5	0.84	0.25	69,89,127,158	0
4	MG	Н	203	1/1	0.88	0.26	30,30,30,30	0
4	MG	С	205	1/1	0.88	0.09	60,60,60,60	0
3	SO4	A	202	5/5	0.89	0.22	73,85,100,116	0
3	SO4	F	202	5/5	0.89	0.34	90,95,108,111	0
5	LFA	D	203	20/20	0.89	0.18	45,63,88,89	0
3	SO4	Н	201	5/5	0.89	0.16	82,85,98,121	0
4	MG	D	205	1/1	0.90	0.08	58,58,58,58	0
5	LFA	В	205	20/20	0.90	0.17	40,61,87,98	0
5	LFA	E	203	20/20	0.91	0.18	41,65,85,90	0
2	GOL	Ε	202	6/6	0.92	0.19	63,69,85,87	0
2	GOL	В	202	6/6	0.94	0.11	52,62,63,65	0
4	MG	A	203	1/1	0.94	0.08	56,56,56,56	0
3	SO4	В	203	5/5	0.96	0.39	80,83,91,92	0
4	MG	В	207	1/1	0.97	0.09	47,47,47,47	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers



as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.












### Electron density around SO4 F 201:


 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 0.7 rmsd) in gray  $\mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)













## Electron density around SO4 B 201: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_{o}\text{-}\mathrm{DF}_{c}$ (at 3 rmsd) in purple (negative) and green (positive)




# Electron density around GOL C 201: 2mF<sub>o</sub>-DF<sub>c</sub> (at 0.7 rmsd) in gray mF<sub>o</sub>-DF<sub>c</sub> (at 3 rmsd) in purple (negative) and green (positive)



## Electron density around SO4 D 204: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${ m mF}_o{ m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









## Electron density around GOL B 204: 2mF<sub>o</sub>-DF<sub>c</sub> (at 0.7 rmsd) in gray mF<sub>o</sub>-DF<sub>c</sub> (at 3 rmsd) in purple (negative) and green (positive)









### Electron density around SO4 C 204:












### 




### Electron density around MG G 203:












## Electron density around SO4 B 206: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${ m mF}_o{ m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)








## Electron density around MG F 203: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${ m mF}_o{ m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)








## Electron density around SO4 E 204: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${ m mF}_o{ m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)




## Electron density around SO4 E 201: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_{o}\text{-}\mathrm{DF}_{c}$ (at 3 rmsd) in purple (negative) and green (positive)










### Electron density around MG C 205:









### Electron density around SO4 A 202:











## Electron density around SO4 F 202: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









### Electron density around SO4 H 201:











### 












## Electron density around GOL E 202: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${ m mF}_o{ m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









## Electron density around MG A 203: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${ m mF}_o{ m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









### 6.5 Other polymers (i)

There are no such residues in this entry.

