



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 5, 2024 – 10:07 AM EDT

PDB ID : 8TXL
Title : Nan Regulatory Protein full length mutant R148A
Authors : Wood, D.M.; Horne, C.R.; Dobson, R.C.J.
Deposited on : 2023-08-23
Resolution : 1.82 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.002 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.38.3


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.82 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	9242 (1.84-1.80)
Clashscore	180529	1080 (1.82-1.82)
Ramachandran outliers	177936	1073 (1.82-1.82)
Sidechain outliers	177891	1073 (1.82-1.82)
RSRZ outliers	164620	9241 (1.84-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	283	 10% 81% 16% ..
1	B	283	 4% 86% 11% .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	ACT	A	303	-	-	X	-

2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 9199 atoms, of which 4454 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Phosphosugar-binding transcriptional regulator.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	275	4354	1396	2150	368	430	10	0	0	0
1	B	276	4484	1433	2222	382	437	10	0	5	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	148	ALA	ARG	engineered mutation	UNP A0A4M6CQT5
B	148	ALA	ARG	engineered mutation	UNP A0A4M6CQT5

- Molecule 2 is PENTAETHYLENE GLYCOL (three-letter code: 1PE) (formula: C₁₀H₂₂O₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
2	A	1	38	10	22	6	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
2	A	1	38	10	22	6	0	0

- Molecule 3 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
3	A	1	7	2	3	2	0	0
3	B	1	7	2	3	2	0	0

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
			Total	C	H			O
4	A	1	10	2	6	2	0	0
4	B	1	10	2	6	2	0	0

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	
			Total	O			S
5	A	1	5	4	1	0	0
5	A	1	5	4	1	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 6 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Cl	0	0
			1	1		
6	B	2	Total	Cl	0	0
			2	2		

- Molecule 7 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	O	P	0	0
			5	4	1		
7	B	1	Total	O	P	0	0
			5	4	1		
7	B	1	Total	O	P	0	0
			5	4	1		

- Molecule 8 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
8	B	1	Total	C	H	O	0	0
			11	3	5	3		
8	B	1	Total	C	H	O	0	0
			14	3	8	3		
8	B	1	Total	C	H	O	0	0
			13	3	7	3		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	84	Total	O	0	0
			84	84		
9	B	96	Total	O	0	0
			96	96		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Phosphosugar-binding transcriptional regulator



- Molecule 1: Phosphosugar-binding transcriptional regulator



4 Data and refinement statistics i

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	62.76Å 104.73Å 205.66Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.34 – 1.82 42.34 – 1.82	Depositor EDS
% Data completeness (in resolution range)	99.9 (42.34-1.82) 99.9 (42.34-1.82)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.33 (at 1.82Å)	Xtrriage
Refinement program	REFMAC 8.0.013, PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.210 , 0.252 0.214 , 0.258	Depositor DCC
R_{free} test set	3062 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	39.6	Xtrriage
Anisotropy	0.525	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 51.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.015 for 1/2*h-1/2*k,-3/2*h-1/2*k,-l 0.029 for 1/2*h+1/2*k,3/2*h-1/2*k,-l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9199	wwPDB-VP
Average B, all atoms (Å ²)	56.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.28% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, CL, PO4, EDO, 1PE, ACT, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.54	0/2245	0.70	1/3032 (0.0%)
1	B	0.56	0/2313	0.70	0/3121
All	All	0.55	0/4558	0.70	1/6153 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	156	LEU	CA-CB-CG	-5.14	103.49	115.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	56	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2204	2150	2150	33	1
1	B	2262	2222	2220	23	1
2	A	32	44	44	5	0
3	A	4	3	3	2	0
3	B	4	3	3	1	0
4	A	4	6	6	0	0
4	B	4	6	6	0	0
5	A	15	0	0	0	0
6	A	1	0	0	0	0
6	B	2	0	0	0	0
7	A	5	0	0	0	0
7	B	10	0	0	1	0
8	B	18	20	24	3	0
9	A	84	0	0	3	1
9	B	96	0	0	1	0
All	All	4745	4454	4456	55	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:6:ILE:HD12	1:A:7:ALA:N	1.88	0.88
1:A:169:MET:HG2	1:A:199:ALA:HB2	1.72	0.71
1:B:51:GLN:H	8:B:303:GOL:H11	1.52	0.71
1:B:262:PHE:O	1:B:265[B]:TYR:HD1	1.77	0.67
1:A:169:MET:HE3	1:A:175:VAL:HG21	1.79	0.63
1:A:6:ILE:HD12	1:A:7:ALA:H	1.66	0.59
1:B:142:MET:HE1	1:B:243:ILE:HG22	1.84	0.58
1:A:269:LYS:HZ1	2:A:302:1PE:H241	1.69	0.56
1:A:41:GLN:N	1:A:41:GLN:OE1	2.39	0.56
1:A:166:THR:HA	1:A:169:MET:HE1	1.90	0.54
1:A:79:ASN:ND2	9:A:403:HOH:O	2.41	0.54
1:A:34:ILE:HD12	1:A:34:ILE:H	1.74	0.53
1:A:269:LYS:HZ1	2:A:302:1PE:H252	1.74	0.52
1:A:182:GLY:HA3	1:A:207:VAL:HG22	1.92	0.52
1:B:141:GLU:HA	1:B:144:LEU:HD12	1.93	0.51
1:B:127:VAL:O	1:B:153:CYS:HA	2.10	0.51
1:A:120:LEU:HD22	1:A:200:LYS:HD3	1.93	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:207:VAL:HG12	1:B:207:VAL:O	2.09	0.51
1:A:252:GLU:HG3	1:B:90:LEU:HD21	1.91	0.51
1:A:127:VAL:O	1:A:153:CYS:HA	2.10	0.50
1:A:185:PRO:HD2	9:A:413:HOH:O	2.12	0.50
1:A:138:VAL:HG13	1:A:243:ILE:HD11	1.93	0.50
1:B:52:ALA:HB3	8:B:304:GOL:H31	1.92	0.50
1:B:240:LEU:HA	1:B:243:ILE:HG12	1.94	0.49
1:A:266:TRP:CE3	3:A:303:ACT:H3	2.47	0.49
1:B:262:PHE:O	1:B:265[B]:TYR:CD1	2.61	0.49
1:A:265:TYR:HB3	2:A:302:1PE:H242	1.95	0.49
1:B:131:GLY:O	1:B:157:THR:HA	2.14	0.48
1:B:99:SER:O	1:B:103:GLU:HG3	2.13	0.48
1:B:140:ARG:NH2	9:B:403:HOH:O	2.49	0.46
1:B:232[A]:ARG:HH11	1:B:232[A]:ARG:HG3	1.79	0.46
1:A:143:LYS:HZ3	2:A:301:1PE:H241	1.80	0.46
1:B:252:GLU:OE2	1:B:255:ARG:NH1	2.49	0.46
1:B:206:SER:OG	3:B:301:ACT:H3	2.16	0.45
1:A:34:ILE:HD12	1:A:34:ILE:N	2.31	0.45
1:A:143:LYS:NZ	1:A:154:GLU:OE1	2.39	0.45
1:A:40:SER:O	1:A:44:THR:HB	2.17	0.44
1:A:108:LEU:HD22	1:A:220:LEU:HD23	1.99	0.44
1:A:130:PHE:HE1	1:A:156:LEU:HD13	1.83	0.44
1:B:240:LEU:HD23	1:B:243:ILE:HD11	2.00	0.44
1:B:114:LEU:HB2	1:B:249:TYR:CE2	2.52	0.44
1:A:270:LYS:NZ	3:A:303:ACT:H1	2.33	0.44
1:A:105:THR:O	1:A:109:ILE:HG13	2.18	0.44
1:B:51:GLN:HB2	8:B:303:GOL:C1	2.47	0.43
1:B:90:LEU:HD12	1:B:90:LEU:H	1.83	0.43
1:A:269:LYS:NZ	2:A:302:1PE:H241	2.33	0.43
1:A:86:LYS:NZ	9:A:411:HOH:O	2.52	0.43
1:B:102:ARG:NH2	1:B:103:GLU:HG2	2.32	0.43
1:B:184:THR:OG1	7:B:309:PO4:O2	2.37	0.42
1:B:142:MET:CE	1:B:243:ILE:HG22	2.48	0.42
1:A:166:THR:HA	1:A:169:MET:CE	2.50	0.42
1:A:180:LEU:HD21	1:A:224:HIS:HE1	1.85	0.42
1:A:66:TYR:CZ	1:A:70:ILE:HG21	2.55	0.41
1:A:34:ILE:H	1:A:34:ILE:CD1	2.32	0.41
1:A:6:ILE:O	1:A:9:VAL:HG13	2.21	0.41

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
9:A:458:HOH:O	9:A:458:HOH:O[3_554]	1.99	0.21
1:A:59:LYS:HZ1	1:B:107:ASP:OD2[4_565]	1.59	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	273/283 (96%)	268 (98%)	5 (2%)	0	100	100
1	B	279/283 (99%)	275 (99%)	4 (1%)	0	100	100
All	All	552/566 (98%)	543 (98%)	9 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	241/252 (96%)	235 (98%)	6 (2%)	42	25
1	B	248/252 (98%)	243 (98%)	5 (2%)	50	35
All	All	489/504 (97%)	478 (98%)	11 (2%)	45	31

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	44	THR
1	A	90	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	100	ASN
1	A	134	SER
1	A	160	ASP
1	A	211	ASP
1	B	46	LYS
1	B	159	GLN
1	B	160	ASP
1	B	237	LEU
1	B	264	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	224	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 18 ligands modelled in this entry, 3 are monoatomic - leaving 15 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	SO4	A	306	-	4,4,4	0.35	0	6,6,6	0.42	0
8	GOL	B	304	-	5,5,5	0.72	0	5,5,5	1.03	0
3	ACT	B	301	-	3,3,3	0.46	0	3,3,3	1.59	1 (33%)
7	PO4	B	308	-	4,4,4	1.39	0	6,6,6	1.21	0
2	1PE	A	301	-	15,15,15	0.13	0	14,14,14	0.18	0
3	ACT	A	303	-	3,3,3	1.06	0	3,3,3	1.49	1 (33%)
4	EDO	A	304	-	3,3,3	0.52	0	2,2,2	0.25	0
5	SO4	A	307	-	4,4,4	0.26	0	6,6,6	0.36	0
2	1PE	A	302	-	15,15,15	0.18	0	14,14,14	0.28	0
7	PO4	A	309	-	4,4,4	0.95	0	6,6,6	0.75	0
8	GOL	B	303	-	5,5,5	0.52	0	5,5,5	0.88	0
7	PO4	B	309	-	4,4,4	1.01	0	6,6,6	0.43	0
4	EDO	B	302	-	3,3,3	0.45	0	2,2,2	0.35	0
8	GOL	B	305	-	5,5,5	1.05	0	5,5,5	1.01	0
5	SO4	A	305	-	4,4,4	0.30	0	6,6,6	0.81	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	GOL	B	304	-	-	2/4/4/4	-
2	1PE	A	301	-	-	11/13/13/13	-
4	EDO	A	304	-	-	1/1/1/1	-
2	1PE	A	302	-	-	7/13/13/13	-
8	GOL	B	303	-	-	2/4/4/4	-
4	EDO	B	302	-	-	1/1/1/1	-
8	GOL	B	305	-	-	2/4/4/4	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	303	ACT	OXT-C-O	2.09	129.79	122.03
3	B	301	ACT	OXT-C-O	2.06	129.67	122.03

There are no chirality outliers.

All (26) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
8	B	305	GOL	C1-C2-C3-O3
2	A	302	1PE	OH6-C15-C25-OH5
2	A	301	1PE	OH5-C14-C24-OH4
2	A	301	1PE	OH2-C12-C22-OH3
2	A	301	1PE	OH7-C16-C26-OH6
2	A	302	1PE	OH2-C12-C22-OH3
2	A	301	1PE	OH4-C13-C23-OH3
8	B	303	GOL	C1-C2-C3-O3
8	B	304	GOL	O1-C1-C2-C3
8	B	305	GOL	O2-C2-C3-O3
2	A	302	1PE	OH7-C16-C26-OH6
2	A	301	1PE	C24-C14-OH5-C25
4	B	302	EDO	O1-C1-C2-O2
8	B	303	GOL	O2-C2-C3-O3
2	A	301	1PE	C15-C25-OH5-C14
2	A	301	1PE	C14-C24-OH4-C13
2	A	301	1PE	C25-C15-OH6-C26
2	A	302	1PE	C13-C23-OH3-C22
4	A	304	EDO	O1-C1-C2-O2
2	A	302	1PE	C23-C13-OH4-C24
2	A	301	1PE	C13-C23-OH3-C22
2	A	302	1PE	OH4-C13-C23-OH3
2	A	301	1PE	OH6-C15-C25-OH5
2	A	301	1PE	C23-C13-OH4-C24
2	A	302	1PE	OH5-C14-C24-OH4
8	B	304	GOL	O1-C1-C2-O2

There are no ring outliers.

7 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	B	304	GOL	1	0
3	B	301	ACT	1	0
2	A	301	1PE	1	0
3	A	303	ACT	2	0
2	A	302	1PE	4	0
8	B	303	GOL	2	0
7	B	309	PO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	275/283 (97%)	0.58	28 (10%) 13 12	36, 54, 85, 144	0
1	B	276/283 (97%)	0.41	12 (4%) 40 39	25, 52, 80, 119	5 (1%)
All	All	551/566 (97%)	0.50	40 (7%) 22 20	25, 53, 81, 144	5 (0%)

All (40) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	265[A]	TYR	5.6
1	A	71	PHE	4.8
1	B	271	LEU	4.4
1	B	208	PRO	3.9
1	B	3	LYS	3.6
1	A	34	ILE	3.5
1	A	73	TYR	3.5
1	A	278	LYS	3.4
1	A	6	ILE	3.3
1	B	274	TYR	3.2
1	A	38	LEU	2.8
1	A	211	ASP	2.7
1	A	212	SER	2.7
1	A	75	HIS	2.6
1	A	47	LEU	2.6
1	A	80	GLN	2.6
1	A	277	GLN	2.6
1	A	274	TYR	2.5
1	A	70	ILE	2.5
1	A	249	TYR	2.4
1	A	210	LYS	2.4
1	B	278	LYS	2.4
1	B	207	VAL	2.4
1	A	49	ILE	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	266	TRP	2.4
1	A	229	TYR	2.3
1	A	31	ALA	2.3
1	A	44	THR	2.3
1	A	7	ALA	2.2
1	B	4	PRO	2.2
1	A	4	PRO	2.2
1	A	63	PHE	2.2
1	A	276	ARG	2.2
1	A	36	ASP	2.1
1	A	32	GLU	2.1
1	B	75	HIS	2.1
1	A	168	ILE	2.1
1	A	27	TYR	2.0
1	B	77	ALA	2.0
1	B	90	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
8	GOL	B	304	6/6	0.72	0.16	58,72,82,98	0
5	SO4	A	307	5/5	0.78	0.10	70,72,96,104	0
8	GOL	B	303	6/6	0.80	0.13	60,74,89,93	0
7	PO4	B	309	5/5	0.80	0.13	73,82,92,93	0
8	GOL	B	305	6/6	0.81	0.22	60,82,99,99	0
4	EDO	B	302	4/4	0.83	0.14	61,73,89,89	0
4	EDO	A	304	4/4	0.86	0.14	54,68,81,85	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	1PE	A	301	16/16	0.88	0.13	48,74,103,108	0
7	PO4	A	309	5/5	0.88	0.10	54,65,73,88	0
3	ACT	A	303	4/4	0.90	0.14	71,78,86,86	0
5	SO4	A	306	5/5	0.90	0.10	41,60,63,69	0
3	ACT	B	301	4/4	0.91	0.11	46,53,64,64	0
7	PO4	B	308	5/5	0.91	0.10	52,59,61,64	0
2	1PE	A	302	16/16	0.92	0.11	36,60,87,93	0
6	CL	B	306	1/1	0.92	0.12	69,69,69,69	0
5	SO4	A	305	5/5	0.97	0.07	34,36,39,39	0
6	CL	B	307	1/1	0.97	0.12	40,40,40,40	0
6	CL	A	308	1/1	0.98	0.11	46,46,46,46	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.