



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 29, 2020 – 11:06 pm BST

PDB ID : 3U61  
Title : Structure of T4 Bacteriophage Clamp Loader Bound To Closed Clamp, DNA and ATP Analog and ADP  
Authors : Kelch, B.A.; Makino, D.L.; O'Donnell, M.; Kuriyan, J.  
Deposited on : 2011-10-11  
Resolution : 3.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1133 (3.20-3.20)
Clashscore	141614	1253 (3.20-3.20)
Ramachandran outliers	138981	1234 (3.20-3.20)
Sidechain outliers	138945	1233 (3.20-3.20)
RSRZ outliers	127900	1095 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	324	
1	C	324	
1	D	324	
1	E	324	
2	A	199	
3	F	228	

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	G	228	<p>%</p> <p>61% 36%</p>
3	H	228	<p>66% 31%</p>
4	I	20	<p>20% 20% 60%</p>
5	J	10	<p>50% 30% 20%</p>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	ADP	B	700	-	-	X	-

## 2 Entry composition i

There are 8 unique types of molecules in this entry. The entry contains 16748 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase accessory protein 44.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	B	305	Total 2407	C 1525	N 412	O 454	S 16	0	0	0
1	C	319	Total 2503	C 1583	N 430	O 473	S 17	0	0	0
1	D	320	Total 2514	C 1590	N 432	O 475	S 17	0	0	0
1	E	294	Total 2292	C 1451	N 394	O 432	S 15	0	0	0

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-4	GLY	-	EXPRESSION TAG	UNP P04526
B	-3	PRO	-	EXPRESSION TAG	UNP P04526
B	-2	GLY	-	EXPRESSION TAG	UNP P04526
B	-1	GLY	-	EXPRESSION TAG	UNP P04526
B	0	SER	-	EXPRESSION TAG	UNP P04526
C	-4	GLY	-	EXPRESSION TAG	UNP P04526
C	-3	PRO	-	EXPRESSION TAG	UNP P04526
C	-2	GLY	-	EXPRESSION TAG	UNP P04526
C	-1	GLY	-	EXPRESSION TAG	UNP P04526
C	0	SER	-	EXPRESSION TAG	UNP P04526
D	-4	GLY	-	EXPRESSION TAG	UNP P04526
D	-3	PRO	-	EXPRESSION TAG	UNP P04526
D	-2	GLY	-	EXPRESSION TAG	UNP P04526
D	-1	GLY	-	EXPRESSION TAG	UNP P04526
D	0	SER	-	EXPRESSION TAG	UNP P04526
E	-4	GLY	-	EXPRESSION TAG	UNP P04526
E	-3	PRO	-	EXPRESSION TAG	UNP P04526
E	-2	GLY	-	EXPRESSION TAG	UNP P04526
E	-1	GLY	-	EXPRESSION TAG	UNP P04526
E	0	SER	-	EXPRESSION TAG	UNP P04526

- Molecule 2 is a protein called DNA polymerase accessory protein 62.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	A	178	1416	909	233	268	6	0	0	0

There are 13 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	188	GLY	-	EXPRESSION TAG	UNP P04527
A	189	LEU	-	EXPRESSION TAG	UNP P04527
A	190	GLU	-	EXPRESSION TAG	UNP P04527
A	191	HIS	-	EXPRESSION TAG	UNP P04527
A	192	HIS	-	EXPRESSION TAG	UNP P04527
A	193	HIS	-	EXPRESSION TAG	UNP P04527
A	194	HIS	-	EXPRESSION TAG	UNP P04527
A	195	HIS	-	EXPRESSION TAG	UNP P04527
A	196	HIS	-	EXPRESSION TAG	UNP P04527
A	197	HIS	-	EXPRESSION TAG	UNP P04527
A	198	HIS	-	EXPRESSION TAG	UNP P04527
A	199	HIS	-	EXPRESSION TAG	UNP P04527
A	200	HIS	-	EXPRESSION TAG	UNP P04527

- Molecule 3 is a protein called DNA polymerase processivity component.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	Se			
3	G	228	1750	1113	288	343	6	0	0	0
3	H	228	1750	1113	288	343	6	0	0	0
3	F	222	1699	1083	280	330	6	0	0	0


- Molecule 4 is a DNA chain called Template DNA strand.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
4	I	8	163	79	29	48	7	0	0	0

- Molecule 5 is a DNA chain called Primer DNA strand.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
5	J	8	162	77	31	46	8	0	0	0







### 3 Residue-property plots i

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA polymerase accessory protein 44




- Molecule 1: DNA polymerase accessory protein 44






W318  
LYS


• Molecule 1: DNA polymerase accessory protein 44




• Molecule 1: DNA polymerase accessory protein 44




• Molecule 2: DNA polymerase accessory protein 62







• Molecule 3: DNA polymerase processivity component



• Molecule 3: DNA polymerase processivity component



• Molecule 3: DNA polymerase processivity component



- Molecule 4: Template DNA strand



- Molecule 5: Primer DNA strand



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	82.30 Å 137.27 Å 222.78 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.65 – 3.20 47.65 – 2.92	Depositor EDS
% Data completeness (in resolution range)	96.2 (47.65-3.20) 89.9 (47.65-2.92)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.12	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.81 (at 2.91 Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.7.1_743)	Depositor
R, $R_{free}$	0.239 , 0.305 0.230 , 0.297	Depositor DCC
$R_{free}$ test set	3572 reflections (7.12%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	73.1	Xtrriage
Anisotropy	0.483	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.28 , 59.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	16748	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	106.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.99% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ADP, 08T

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	B	0.52	0/2450	0.71	0/3303
1	C	0.58	0/2546	0.75	1/3434 (0.0%)
1	D	0.54	0/2558	0.77	3/3449 (0.1%)
1	E	0.52	0/2331	0.73	1/3144 (0.0%)
2	A	0.48	0/1440	0.74	0/1938
3	F	0.52	0/1721	0.65	0/2324
3	G	0.59	0/1774	0.67	0/2395
3	H	0.50	0/1774	0.63	0/2395
4	I	1.03	0/182	1.86	7/280 (2.5%)
5	J	1.03	0/181	2.09	13/276 (4.7%)
All	All	0.55	0/16957	0.77	25/22938 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1
1	D	0	1
1	E	0	2
2	A	0	2
3	F	0	1
All	All	0	7

There are no bond length outliers.

The worst 5 of 25 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	J	3	DT	O4'-C1'-N1	9.74	114.82	108.00

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	J	3	DT	C5-C4-O4	-8.25	119.12	124.90
5	J	7	DC	O4'-C1'-N1	8.20	113.74	108.00
5	J	8	DA	O4'-C1'-N9	7.47	113.23	108.00
5	J	5	DG	O4'-C1'-C2'	-7.41	99.97	105.90

There are no chirality outliers.

5 of 7 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	A	2	SER	Peptide
1	C	128	TYR	Mainchain
1	D	217	LYS	Mainchain
1	E	157	PHE	Peptide
1	E	49	SER	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	2407	0	2424	182	0
1	C	2503	0	2524	231	0
1	D	2514	0	2543	245	0
1	E	2292	0	2302	198	0
2	A	1416	0	1431	146	0
3	F	1699	0	1716	72	1
3	G	1750	0	1752	112	1
3	H	1750	0	1752	66	0
4	I	163	0	93	18	0
5	J	162	0	90	6	0
6	B	27	0	12	10	0
7	B	1	0	0	0	0
7	C	1	0	0	0	0
7	D	1	0	0	0	0
8	C	31	0	13	5	0
8	D	31	0	13	1	0
All	All	16748	0	16665	1056	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 32.

The worst 5 of 1056 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:97:PHE:CD1	3:G:1204:GLN:HB3	1.57	1.40
1:C:9:HIS:CD2	1:D:41:LYS:HB3	1.73	1.22
1:D:174:ARG:HH11	1:D:174:ARG:CB	1.53	1.20
1:D:174:ARG:HB3	1:D:174:ARG:NH1	1.55	1.20
1:C:174:ARG:HB3	1:C:174:ARG:NH1	1.55	1.18

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:G:1193:ASN:OD1	3:F:3176:ASN:ND2[4_445]	2.12	0.08

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	B	301/324 (93%)	282 (94%)	18 (6%)	1 (0%)	41 74
1	C	317/324 (98%)	302 (95%)	11 (4%)	4 (1%)	12 47
1	D	318/324 (98%)	300 (94%)	13 (4%)	5 (2%)	9 43
1	E	290/324 (90%)	277 (96%)	12 (4%)	1 (0%)	41 74
2	A	174/199 (87%)	140 (80%)	33 (19%)	1 (1%)	25 64
3	F	220/228 (96%)	217 (99%)	3 (1%)	0	100 100
3	G	226/228 (99%)	223 (99%)	3 (1%)	0	100 100
3	H	226/228 (99%)	222 (98%)	4 (2%)	0	100 100
All	All	2072/2179 (95%)	1963 (95%)	97 (5%)	12 (1%)	25 64

5 of 12 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	219	VAL
1	C	97	PHE
1	D	234	ALA
1	B	259	ALA
1	C	259	ALA

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	266/279 (95%)	239 (90%)	27 (10%)	7	29
1	C	276/279 (99%)	244 (88%)	32 (12%)	5	24
1	D	278/279 (100%)	248 (89%)	30 (11%)	6	27
1	E	250/279 (90%)	225 (90%)	25 (10%)	7	30
2	A	154/174 (88%)	143 (93%)	11 (7%)	14	47
3	F	183/183 (100%)	170 (93%)	13 (7%)	14	47
3	G	189/183 (103%)	177 (94%)	12 (6%)	18	52
3	H	189/183 (103%)	177 (94%)	12 (6%)	18	52
All	All	1785/1839 (97%)	1623 (91%)	162 (9%)	9	34

5 of 162 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	121	LEU
1	E	32	THR
3	F	3025	SER
1	D	146	LYS
1	D	232	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 26 such sidechains are listed below:



Mol	Chain	Res	Type
1	D	101	GLN
1	D	295	HIS
3	H	2190	GLN
1	D	230	ASN
1	D	293	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 3 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
6	ADP	B	700	-	24,29,29	0.94	1 (4%)	29,45,45	1.54	6 (20%)
8	08T	C	700	7	26,33,33	2.91	11 (42%)	25,52,52	5.01	5 (20%)
8	08T	D	700	7	26,33,33	2.93	11 (42%)	25,52,52	5.48	8 (32%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	ADP	B	700	-	-	7/12/32/32	0/3/3/3
8	08T	C	700	7	-	1/12/38/38	0/3/3/3
8	08T	D	700	7	-	2/12/38/38	0/3/3/3

The worst 5 of 23 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	C	700	08T	F1-BE	-7.58	1.35	1.54
8	D	700	08T	F1-BE	-6.99	1.37	1.54
8	C	700	08T	F3-BE	-5.86	1.39	1.54
8	D	700	08T	F2-BE	-5.75	1.40	1.54
8	D	700	08T	F3-BE	-5.64	1.40	1.54

The worst 5 of 19 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	D	700	08T	C1'-N9-C4	-25.14	82.46	126.64
8	C	700	08T	C1'-N9-C4	-22.41	87.26	126.64
8	C	700	08T	C3'-C2'-C1'	6.79	111.20	100.98
8	D	700	08T	O4'-C1'-C2'	-6.30	97.72	106.93
8	C	700	08T	O4'-C1'-C2'	-5.05	99.54	106.93

There are no chirality outliers.

5 of 10 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	B	700	ADP	PA-O3A-PB-O2B
6	B	700	ADP	PA-O3A-PB-O3B
6	B	700	ADP	C5'-O5'-PA-O1A
6	B	700	ADP	C5'-O5'-PA-O2A
6	B	700	ADP	O4'-C4'-C5'-O5'


There are no ring outliers.


3 monomers are involved in 16 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	B	700	ADP	10	0
8	C	700	08T	5	0
8	D	700	08T	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In

addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å <sup>2</sup> )	Q < 0.9
1	B	305/324 (94%)	-0.06	6 (1%) 65 51	81, 104, 150, 221	0
1	C	319/324 (98%)	-0.15	1 (0%) 94 92	57, 95, 144, 211	0
1	D	320/324 (98%)	-0.04	5 (1%) 72 59	54, 89, 179, 232	0
1	E	294/324 (90%)	0.06	12 (4%) 37 24	69, 117, 189, 246	0
2	A	178/199 (89%)	0.07	4 (2%) 62 48	81, 120, 181, 223	0
3	F	216/228 (94%)	-0.12	3 (1%) 75 63	70, 96, 159, 252	0
3	G	222/228 (97%)	-0.26	3 (1%) 75 63	58, 84, 130, 188	0
3	H	222/228 (97%)	-0.23	1 (0%) 91 86	71, 95, 143, 200	0
4	I	8/20 (40%)	-0.23	0 100 100	83, 92, 123, 127	0
5	J	8/10 (80%)	-0.08	0 100 100	107, 118, 140, 160	0
All	All	2092/2209 (94%)	-0.09	35 (1%) 70 57	54, 100, 167, 252	0

The worst 5 of 35 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	A	73	PRO	6.3
2	A	177	GLN	4.9
1	D	253	LEU	4.1
1	B	214	TYR	4.1
1	D	285	TYR	4.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
7	MG	B	800	1/1	0.79	0.29	83,83,83,83	0
7	MG	D	800	1/1	0.90	0.52	83,83,83,83	0
6	ADP	B	700	27/27	0.92	0.18	83,83,83,83	0
8	08T	D	700	31/31	0.94	0.24	83,83,83,83	0
7	MG	C	800	1/1	0.95	0.46	83,83,83,83	0
8	08T	C	700	31/31	0.95	0.22	83,83,83,83	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around ADP B 700:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around 08T D 700:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.