



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 6, 2023 – 11:49 AM EST

PDB ID : 3U6B  
Title : Ef-tu (escherichia coli) in complex with nvp-ldi028  
Authors : Palestrant, D.J.  
Deposited on : 2011-10-12  
Resolution : 2.12 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.12 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	6241 (2.14-2.10)
Clashscore	141614	6778 (2.14-2.10)
Ramachandran outliers	138981	6705 (2.14-2.10)
Sidechain outliers	138945	6706 (2.14-2.10)
RSRZ outliers	127900	6112 (2.14-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	394	
1	B	394	
2	C	12	
2	D	12	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	MH6	C	11	-	X	-	-
2	BB9	C	9	-	X	-	-
2	MH6	D	11	-	X	-	-

## 2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 6044 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Elongation factor Tu 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	369	2837	1802	482	540	13	0	0	0
1	B	376	2870	1817	492	548	13	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	expression tag	UNP P0CE47
A	1	ALA	-	expression tag	UNP P0CE47
B	0	MET	-	expression tag	UNP P0CE47
B	1	ALA	-	expression tag	UNP P0CE47

- Molecule 2 is a protein called Thiocillin GE2270 analogue NVP-LDI028.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	C	12	84	55	13	10	6	0	0	0
2	D	12	84	55	13	10	6	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	12	8BB	CYS	SEE REMARK 999	UNP Q7M0J8
D	12	8BB	CYS	SEE REMARK 999	UNP Q7M0J8

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mg	0	0
			1	1		

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Mg	0	0
			1	1		

- Molecule 4 is GUANOSINE-5'-DIPHOSPHATE (three-letter code: GDP) (formula:  $C_{10}H_{15}N_5O_{11}P_2$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
4	A	1	Total	C	N	O	P	0	0
			28	10	5	11	2		
4	B	1	Total	C	N	O	P	0	0
			28	10	5	11	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	62	Total	O	0	0
			62	62		
5	B	47	Total	O	0	0
			47	47		
5	C	2	Total	O	0	0
			2	2		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Elongation factor Tu 1





- Molecule 1: Elongation factor Tu 1



- Molecule 2: Thiocillin GE2270 analogue NVP-LDI028



- Molecule 2: Thiocillin GE2270 analogue NVP-LDI028



## 4 Data and refinement statistics i

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	97.25Å 100.38Å 156.48Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	8.00 – 2.12 42.24 – 2.12	Depositor EDS
% Data completeness (in resolution range)	85.8 (8.00-2.12) 88.2 (42.24-2.12)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.98 (at 2.12Å)	Xtrriage
Refinement program	CNX, BUSTER 2.11.2	Depositor
R, $R_{free}$	0.196 , 0.257 0.205 , 0.264	Depositor DCC
$R_{free}$ test set	1955 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	30.9	Xtrriage
Anisotropy	0.416	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 28.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.28$	Xtrriage
Estimated twinning fraction	0.118 for -k,-h,-l	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	6044	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	37.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.00% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BB9, MG, BB7, MEN, BB6, 8BB, GDP, MH6, BB8

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.48	0/2889	0.71	0/3913
1	B	0.49	0/2921	0.76	0/3956
2	C	2.63	1/10 (10.0%)	1.53	0/9
2	D	2.93	1/10 (10.0%)	1.45	0/9
All	All	0.51	2/5830 (0.0%)	0.74	0/7887

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	C	1	0
2	D	1	0
All	All	2	0

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	1	SER	CA-CB	-8.92	1.39	1.52
2	C	1	SER	CA-CB	-7.84	1.41	1.52

There are no bond angle outliers.

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	C	1	SER	CA
2	D	1	SER	CA

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2837	0	2845	25	0
1	B	2870	0	2858	25	0
2	C	84	0	53	2	0
2	D	84	0	53	4	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	28	0	12	0	0
4	B	28	0	12	0	0
5	A	62	0	0	1	0
5	B	47	0	0	0	0
5	C	2	0	0	0	0
All	All	6044	0	5833	52	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (52) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:163:PRO:O	1:A:167:THR:HG23	1.88	0.73
1:B:32:THR:HG21	1:B:67:VAL:HG11	1.76	0.68
1:B:74:ARG:NH2	1:B:199:ILE:O	2.20	0.68
1:B:301:HIS:HD2	1:B:393:GLY:HA2	1.63	0.64
1:B:329:GLN:HE21	1:B:336:ASP:HB3	1.63	0.63
1:A:66:HIS:CE1	1:A:79:VAL:HG22	2.39	0.57
1:B:157:LEU:HD13	1:B:167:THR:HG21	1.87	0.57
1:B:176:LYS:HB3	1:B:184:TRP:CD1	2.41	0.56
1:A:167:THR:HG22	5:A:518:HOH:O	2.07	0.55
1:A:327:ARG:HG3	1:A:340:THR:HG22	1.90	0.54
1:A:329:GLN:HE21	1:A:336:ASP:HB3	1.71	0.54
1:A:28:THR:O	1:A:32:THR:HG23	2.08	0.53
1:B:69:TYR:OH	1:B:78:HIS:HD2	1.93	0.52
1:A:273:ASN:O	2:D:3:MEN:HE21	2.10	0.52
1:A:150:GLU:HG2	1:A:169:ILE:HG21	1.93	0.51
2:D:3:MEN:HE23	2:D:10:BB9:SG	2.51	0.51

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:237:LYS:HG2	1:B:267:GLU:HB3	1.92	0.50
1:B:163:PRO:O	1:B:167:THR:HG23	2.12	0.50
1:A:217:VAL:HG22	1:A:227:VAL:HG12	1.95	0.49
1:B:277:LEU:HB2	2:C:12:8BB:CB	2.43	0.49
1:A:288:ARG:HG2	1:A:335:THR:HB	1.94	0.48
1:A:157:LEU:HD13	1:A:167:THR:HG21	1.95	0.48
1:A:212:LEU:HD12	1:A:231:VAL:HG22	1.94	0.48
1:B:301:HIS:CD2	1:B:393:GLY:HA2	2.48	0.48
1:B:342:GLU:HB2	1:B:359:VAL:HB	1.97	0.46
1:A:69:TYR:OH	1:A:78:HIS:HD2	1.98	0.46
1:A:132:VAL:HB	1:A:169:ILE:HG12	1.96	0.46
1:B:70:ASP:OD1	1:B:75:HIS:HD2	1.98	0.46
1:B:95:ALA:HA	1:B:96:ALA:HA	1.82	0.46
1:B:28:THR:O	1:B:32:THR:HG23	2.16	0.46
1:A:308:VAL:O	1:A:355:ASN:HA	2.17	0.45
1:B:210:PHE:CE1	1:B:236:ILE:HB	2.51	0.45
1:B:247:ILE:HD11	1:B:289:GLY:HA3	1.99	0.45
1:A:125:VAL:HG12	1:A:387:VAL:HG11	1.99	0.45
1:A:338:THR:HB	1:A:363:ILE:HG12	1.99	0.45
1:A:66:HIS:ND1	1:A:79:VAL:HG22	2.32	0.44
1:B:244:ILE:HG21	1:B:286:ILE:HG21	1.98	0.44
1:A:260:MET:HG3	2:D:3:MEN:HE22	2.00	0.44
1:B:273:ASN:O	2:C:3:MEN:HE21	2.18	0.44
1:A:19:HIS:HA	1:A:114:GLN:HB2	2.00	0.44
1:B:306:SER:HB2	1:B:387:VAL:O	2.18	0.43
1:A:305:GLU:HG2	1:A:359:VAL:HG22	2.01	0.43
1:B:305:GLU:HG3	1:B:392:LEU:HD11	2.00	0.43
1:A:32:THR:HG21	1:A:67:VAL:HG21	2.00	0.43
1:B:247:ILE:HD12	1:B:290:GLN:HG3	2.00	0.42
1:A:251:GLN:HE22	1:A:285:GLU:HB3	1.84	0.42
1:A:300:PRO:HB2	1:A:365:PRO:HB2	2.01	0.42
2:D:2:BB9:N	2:D:10:BB9:N	2.67	0.42
1:B:300:PRO:HB2	1:B:365:PRO:HB2	2.01	0.42
1:A:214:ILE:HD12	1:A:227:VAL:HB	2.02	0.42
1:B:107:ALA:HB2	1:B:134:LEU:HD22	2.02	0.41
1:B:146:LEU:HD13	1:B:171:ARG:HD3	2.01	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	363/394 (92%)	353 (97%)	8 (2%)	2 (1%)	25	20
1	B	370/394 (94%)	350 (95%)	18 (5%)	2 (0%)	29	25
2	C	2/12 (17%)	1 (50%)	1 (50%)	0	100	100
2	D	2/12 (17%)	1 (50%)	0	1 (50%)	0	0
All	All	737/812 (91%)	705 (96%)	27 (4%)	5 (1%)	22	17

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	D	5	VAL
1	A	333	ARG
1	B	7	ARG
1	A	247	ILE
1	B	247	ILE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	305/325 (94%)	292 (96%)	13 (4%)	29	28
1	B	304/325 (94%)	291 (96%)	13 (4%)	29	28
2	C	1/2 (50%)	1 (100%)	0	100	100
2	D	1/2 (50%)	1 (100%)	0	100	100
All	All	611/654 (93%)	585 (96%)	26 (4%)	29	28

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	32	THR
1	A	134	LEU
1	A	145	LEU
1	A	183	GLU
1	A	242	VAL
1	A	256	THR
1	A	266	ASP
1	A	288	ARG
1	A	311	LEU
1	A	327	ARG
1	A	335	THR
1	A	377	ARG
1	A	381	ARG
1	B	9	LYS
1	B	20	VAL
1	B	32	THR
1	B	58	ARG
1	B	67	VAL
1	B	134	LEU
1	B	145	LEU
1	B	232	GLU
1	B	249	GLU
1	B	262	ARG
1	B	284	GLU
1	B	313	LYS
1	B	391	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	78	HIS
1	A	124	GLN
1	A	251	GLN
1	A	329	GLN
1	B	11	HIS
1	B	75	HIS
1	B	78	HIS
1	B	301	HIS
1	B	329	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

18 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	BB6	D	4	2	4,6,7	2.50	2 (50%)	2,7,9	1.63	1 (50%)
2	BB9	C	2	2	3,5,6	2.53	2 (66%)	1,5,7	5.73	1 (100%)
2	BB8	C	8	2	11,11,13	2.37	3 (27%)	12,14,17	1.44	3 (25%)
2	8BB	D	12	2	15,16,16	1.82	3 (20%)	17,21,21	1.22	2 (11%)
2	BB9	D	2	2	3,5,6	2.70	1 (33%)	1,5,7	4.30	1 (100%)
2	BB8	D	8	2	11,11,13	2.35	4 (36%)	12,14,17	0.99	1 (8%)
2	BB6	C	4	2	4,6,7	2.49	2 (50%)	2,7,9	1.89	1 (50%)
2	BB9	D	9	2	2,4,6	3.63	1 (50%)	3,4,7	2.99	2 (66%)
2	BB9	D	10	2	2,4,6	3.60	1 (50%)	3,4,7	5.14	2 (66%)
2	MH6	D	11	2	3,3,6	4.24	3 (100%)	1,3,7	0.60	0
2	MEN	C	3	2	7,7,9	0.66	0	6,8,11	1.19	0
2	MEN	D	3	2	7,7,9	0.74	0	6,8,11	1.51	2 (33%)
2	BB7	D	6	2	6,8,9	2.17	2 (33%)	3,9,11	1.59	1 (33%)
2	MH6	C	11	2	3,3,6	3.83	3 (100%)	1,3,7	1.14	0
2	BB9	C	9	2	2,4,6	3.53	1 (50%)	3,4,7	2.69	3 (100%)
2	BB9	C	10	2	2,4,6	4.09	1 (50%)	3,4,7	5.13	2 (66%)
2	8BB	C	12	2	15,16,16	1.95	3 (20%)	17,21,21	1.02	1 (5%)
2	BB7	C	6	2	6,8,9	2.20	2 (33%)	3,9,11	1.54	1 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BB6	D	4	2	-	0/0/6/8	-
2	BB9	C	2	2	-	0/0/4/6	-
2	BB8	C	8	2	-	2/8/8/12	0/1/1/1
2	8BB	D	12	2	-	3/10/24/24	0/1/1/1
2	BB9	D	2	2	-	0/0/4/6	-
2	BB8	D	8	2	-	1/8/8/12	0/1/1/1
2	BB6	C	4	2	-	0/0/6/8	-
2	BB9	D	9	2	-	0/0/2/6	-
2	BB9	D	10	2	-	0/0/2/6	-
2	MEN	C	3	2	-	0/6/6/10	-
2	MEN	D	3	2	-	0/6/6/10	-
2	BB7	D	6	2	-	1/1/9/11	-
2	BB9	C	9	2	-	0/0/2/6	-
2	BB9	C	10	2	-	0/0/2/6	-
2	8BB	C	12	2	-	5/10/24/24	0/1/1/1
2	BB7	C	6	2	-	0/1/9/11	-

All (34) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	11	MH6	CB-CA	-6.02	1.40	1.49
2	C	10	BB9	CA-N	5.73	1.47	1.33
2	C	12	8BB	CA-N	5.56	1.46	1.34
2	D	12	8BB	CA-N	5.39	1.45	1.34
2	C	8	BB8	OB-CB	-5.37	1.32	1.42
2	C	11	MH6	CB-CA	-5.33	1.41	1.49
2	D	8	BB8	OB-CB	-5.18	1.32	1.42
2	D	10	BB9	CA-N	4.99	1.45	1.33
2	D	9	BB9	CA-N	4.90	1.45	1.33
2	C	9	BB9	CA-N	4.69	1.45	1.33
2	D	2	BB9	CA-N	4.45	1.46	1.35
2	D	6	BB7	CA-N	4.44	1.47	1.36
2	C	4	BB6	CA-N	4.30	1.47	1.36
2	C	6	BB7	CA-N	4.28	1.47	1.36
2	D	4	BB6	CA-N	4.21	1.46	1.36
2	C	12	8BB	CA-N1	3.90	1.49	1.40
2	C	2	BB9	CA-N	3.80	1.44	1.35
2	D	12	8BB	CA-N1	3.47	1.48	1.40
2	D	8	BB8	CD1-CG	3.32	1.44	1.39
2	D	11	MH6	C-CA	-3.22	1.44	1.49
2	C	8	BB8	CG-CB	-3.21	1.47	1.51
2	C	6	BB7	C-CA	3.10	1.50	1.45
2	C	11	MH6	CA-N	2.91	1.35	1.27

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	11	MH6	CA-N	2.71	1.34	1.27
2	C	11	MH6	C-CA	-2.67	1.45	1.49
2	D	6	BB7	C-CA	2.61	1.49	1.45
2	D	4	BB6	C-CA	2.57	1.49	1.45
2	C	8	BB8	CD1-CG	2.50	1.43	1.39
2	C	4	BB6	C-CA	2.42	1.49	1.45
2	C	12	8BB	O98-C96	-2.42	1.22	1.30
2	D	8	BB8	CD2-CG	2.42	1.43	1.39
2	D	12	8BB	O98-C96	-2.31	1.23	1.30
2	C	2	BB9	C-CA	2.18	1.48	1.45
2	D	8	BB8	CE2-CZ	2.13	1.43	1.38

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	10	BB9	C-CA-N	8.30	125.81	116.53
2	D	10	BB9	C-CA-N	8.23	125.73	116.53
2	C	2	BB9	O-C-CA	-5.73	118.10	125.39
2	D	2	BB9	O-C-CA	-4.30	119.93	125.39
2	D	9	BB9	C-CA-N	4.23	121.25	116.53
2	D	10	BB9	CB-CA-N	-3.39	111.67	122.50
2	C	9	BB9	C-CA-N	3.24	120.14	116.53
2	C	10	BB9	CB-CA-N	-3.15	112.41	122.50
2	C	8	BB8	C-CA-CB	-2.88	107.58	112.00
2	D	6	BB7	O-C-CA	-2.71	120.49	125.54
2	C	6	BB7	O-C-CA	-2.66	120.58	125.54
2	D	12	8BB	C94-C-N1	-2.57	110.64	114.22
2	C	9	BB9	CB-CA-N	-2.53	114.40	122.50
2	D	9	BB9	CB-CA-N	-2.52	114.44	122.50
2	C	4	BB6	O-C-CA	-2.48	120.91	125.54
2	C	8	BB8	CG-CB-CA	-2.45	108.45	112.34
2	D	3	MEN	CB-CG-ND2	-2.43	112.22	115.48
2	C	9	BB9	C-CA-CB	2.21	125.42	121.39
2	C	8	BB8	CD2-CG-CD1	2.21	121.04	118.29
2	D	3	MEN	C-CA-CB	2.12	114.39	111.94
2	D	4	BB6	O-C-CA	-2.11	121.61	125.54
2	D	8	BB8	CG-CB-CA	-2.10	109.01	112.34
2	D	12	8BB	O-C-N1	2.04	125.79	122.57
2	C	12	8BB	O98-C96-C91	2.04	119.56	114.21

There are no chirality outliers.

All (12) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
2	C	8	BB8	C-CA-CB-CG
2	D	8	BB8	C-CA-CB-CG
2	C	12	8BB	O-C-C94-C93
2	D	6	BB7	CB-CB1-OB2-CB3
2	D	12	8BB	C92-C91-C96-O97
2	C	12	8BB	C92-C91-C96-O97
2	C	12	8BB	C92-C91-C96-O98
2	D	12	8BB	C92-C91-C96-O98
2	C	12	8BB	N1-C-C94-C93
2	C	12	8BB	N1-C-C94-C95
2	C	8	BB8	OB-CB-CG-CD1
2	D	12	8BB	O-C-C94-C93

There are no ring outliers.

5 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	2	BB9	1	0
2	D	10	BB9	2	0
2	C	3	MEN	1	0
2	D	3	MEN	3	0
2	C	12	8BB	1	0

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	GDP	B	401	3	24,30,30	1.05	2 (8%)	30,47,47	0.58	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	GDP	A	402	3	24,30,30	0.90	1 (4%)	30,47,47	0.62	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GDP	B	401	3	-	2/12/32/32	0/3/3/3
4	GDP	A	402	3	-	3/12/32/32	0/3/3/3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	401	GDP	C5-C6	-3.15	1.41	1.47
4	B	401	GDP	C8-N7	-2.32	1.31	1.35
4	A	402	GDP	C8-N7	-2.28	1.31	1.35

There are no bond angle outliers.

There are no chirality outliers.

All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	402	GDP	PA-O3A-PB-O2B
4	A	402	GDP	PA-O3A-PB-O3B
4	B	401	GDP	PA-O3A-PB-O2B
4	B	401	GDP	PA-O3A-PB-O3B
4	A	402	GDP	PA-O3A-PB-O1B

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and

any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	369/394 (93%)	-0.04	1 (0%) 94 95	19, 34, 58, 70	0
1	B	376/394 (95%)	0.10	9 (2%) 59 64	17, 35, 62, 86	0
2	C	3/12 (25%)	0.62	1 (33%) 0 0	50, 50, 50, 66	0
2	D	3/12 (25%)	0.39	0 100 100	45, 45, 60, 69	0
All	All	751/812 (92%)	0.04	11 (1%) 73 77	17, 35, 60, 86	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	264	LEU	3.0
1	B	286	ILE	2.8
1	B	96	ALA	2.4
1	B	42	ALA	2.3
1	B	204	ARG	2.3
1	B	379	GLY	2.2
1	B	280	GLY	2.2
1	B	393	GLY	2.2
1	B	264	LEU	2.1
2	C	5	VAL	2.1
1	B	57	ALA	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	BB7	D	6	9/10	0.84	0.22	65,73,78,79	0

*Continued on next page...*

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	8BB	C	12	16/16	0.84	0.34	61,85,91,91	0
2	8BB	D	12	16/16	0.84	0.29	56,77,83,85	0
2	MH6	C	11	4/7	0.90	0.16	54,54,57,61	0
2	BB9	C	2	6/7	0.91	0.13	43,44,46,47	0
2	BB7	C	6	9/10	0.92	0.18	59,65,71,71	0
2	BB8	D	8	11/13	0.93	0.14	40,45,51,51	0
2	BB6	C	4	7/8	0.93	0.16	54,55,61,63	0
2	MEN	C	3	8/10	0.93	0.10	48,52,54,56	0
2	BB6	D	4	7/8	0.94	0.13	52,54,59,59	0
2	BB9	C	9	5/7	0.95	0.10	42,43,43,45	0
2	BB9	C	10	5/7	0.95	0.10	43,48,50,50	0
2	BB8	C	8	11/13	0.95	0.13	38,41,45,46	0
2	MH6	D	11	4/7	0.95	0.10	42,46,47,55	0
2	BB9	D	2	6/7	0.95	0.12	44,46,46,47	0
2	MEN	D	3	8/10	0.95	0.12	46,48,49,50	0
2	BB9	D	9	5/7	0.97	0.09	40,40,42,42	0
2	BB9	D	10	5/7	0.98	0.10	37,39,41,42	0

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	MG	A	401	1/1	0.97	0.06	25,25,25,25	0
3	MG	B	402	1/1	0.99	0.08	22,22,22,22	0
4	GDP	A	402	28/28	0.99	0.10	21,24,29,31	0
4	GDP	B	401	28/28	0.99	0.10	15,20,25,32	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around GDP A 402:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around GDP B 401:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.