

wwPDB EM Validation Summary Report (i)

Oct 8, 2022 – 07:12 PM EDT

PDB ID : 7U6E

EMDB ID : EMD-26364

Title: Head region of insulin receptor ectodomain (A-isoform) bound to the non-

insulin agonist IM462

Authors: Kirk, N.S.; Lawrence, M.C.

Deposited on : 2022-03-03

Resolution : 3.00 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev43

Mogul : 1.8.5 (274361), CSD as541be (2020)

MolProbity: 4.02b-467

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

MapQ: 1.9.9

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.31.2

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $ELECTRON\ MICROSCOPY$

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	${ m EM\ structures} \ (\#{ m Entries})$
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion <40%). The numeric value is given above the bar.

Mol	Chain	Length		Qua	lity of chain		
1	A	21	19%		100%		
2	В	30	13%	80%			17%
3	Е	930		52%	11%	37%	
3	F	930	28%	5%	67%		
4	G	23	39%	83%			17%
4	Н	23	39% 43%		39%		17%
5	С	2			100%		



2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 8004 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Insulin A chain.

Mol	Chain	Residues		${f Atoms}$					Trace
1	A	21	Total 163	C 99	11	O 35	S 4	0	0

• Molecule 2 is a protein called Insulin B chain.

Mol	Chain	Residues		Ato	ms			AltConf	Trace
9	D	25	Total	С	N	О	S	0	0
	Б	29	200	129	34	35	2		U

• Molecule 3 is a protein called Isoform Short of Insulin receptor.

I	Mol	Chain	Residues	Atoms					AltConf	Trace
	3	Е	584	Total 4708	C 2980		O 869	S 42	0	0
	3	F	308	Total 2507	C 1605	N 427	O 467	S 8	0	0

There are 118 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
Е	144	HIS	TYR	conflict	UNP P06213-2
Е	421	THR	ILE	conflict	UNP P06213-2
Е	465	LYS	GLN	conflict	UNP P06213-2
Е	?	-	VAL	deletion	UNP P06213-2
Е	?	-	THR	deletion	UNP P06213-2
Е	?	-	VAL	deletion	UNP P06213-2
Е	?	-	ALA	deletion	UNP P06213-2
Е	?	-	VAL	deletion	UNP P06213-2
Е	?	-	PRO	deletion	UNP P06213-2
Е	?	-	THR	deletion	UNP P06213-2
Е	?	-	VAL	deletion	UNP P06213-2
Е	?	-	ALA	deletion	UNP P06213-2
Е	?	-	ALA	deletion	UNP P06213-2



 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled	Actual	Comment	Reference
E	?	-	PHE	deletion	UNP P06213-2
Е	?	-	PRO	deletion	UNP P06213-2
Е	?	-	ASN	deletion	UNP P06213-2
Е	?	-	THR	deletion	UNP P06213-2
Е	?	-	SER	deletion	UNP P06213-2
Е	?	-	SER	deletion	UNP P06213-2
Е	?	-	THR	deletion	UNP P06213-2
Е	?	-	SER	deletion	UNP P06213-2
Е	?	-	VAL	deletion	UNP P06213-2
E	731	ALA	PRO	conflict	UNP P06213-2
E	732	GLY	THR	conflict	UNP P06213-2
Е	733	ASN	SER	conflict	UNP P06213-2
E	734	ASN	PRO	conflict	UNP P06213-2
E	898	ARG	-	expression tag	UNP P06213-2
E	899	MET	-	expression tag	UNP P06213-2
E	900	LYS	-	expression tag	UNP P06213-2
E	901	GLN	-	expression tag	UNP P06213-2
E	902	LEU	-	expression tag	UNP P06213-2
E	903	GLU	-	expression tag	UNP P06213-2
E	904	ASP	-	expression tag	UNP P06213-2
E	905	LYS	-	expression tag	UNP P06213-2
Ε	906	VAL	-	expression tag	UNP P06213-2
E	907	GLU	-	expression tag	UNP P06213-2
Е	908	GLU	-	expression tag	UNP P06213-2
E	909	LEU	-	expression tag	UNP P06213-2
Е	910	LEU	-	expression tag	UNP P06213-2
Е	911	SER	-	expression tag	UNP P06213-2
E	912	LYS	-	expression tag	UNP P06213-2
E	913	ASN	-	expression tag	UNP P06213-2
Е	914	TYR	-	expression tag	UNP P06213-2
Е	915	HIS	-	expression tag	UNP P06213-2
E	916	LEU	-	expression tag	UNP P06213-2
E	917	GLU	-	expression tag	UNP P06213-2
E	918	ASN	-	expression tag	UNP P06213-2
E	919	GLU	-	expression tag	UNP P06213-2
Е	920	VAL	-	expression tag	UNP P06213-2
E	921	ALA	-	expression tag	UNP P06213-2
E	922	ARG	-	expression tag	UNP P06213-2
E	923	LEU	-	expression tag	UNP P06213-2
E	924	LYS	-	expression tag	UNP P06213-2
E	925	LYS	-	expression tag	UNP P06213-2
E	926	LEU	-	expression tag	UNP P06213-2



 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled	Actual	Comment	Reference
Е	927	VAL	-	expression tag	UNP P06213-2
Е	928	GLY	-	expression tag	UNP P06213-2
Е	929	GLU	-	expression tag	UNP P06213-2
Е	930	ARG	-	expression tag	UNP P06213-2
F	144	HIS	TYR	conflict	UNP P06213-2
F	421	THR	ILE	conflict	UNP P06213-2
F	465	LYS	GLN	conflict	UNP P06213-2
F	?	-	VAL	deletion	UNP P06213-2
F	?	-	THR	deletion	UNP P06213-2
F	?	-	VAL	deletion	UNP P06213-2
F	?	-	ALA	deletion	UNP P06213-2
F	?	-	VAL	deletion	UNP P06213-2
F	?	-	PRO	deletion	UNP P06213-2
F	?	-	THR	deletion	UNP P06213-2
F	?	-	VAL	deletion	UNP P06213-2
F	?	-	ALA	deletion	UNP P06213-2
F	?	-	ALA	deletion	UNP P06213-2
F	?	-	PHE	deletion	UNP P06213-2
F	?	-	PRO	deletion	UNP P06213-2
F	?	-	ASN	deletion	UNP P06213-2
F	?	-	THR	deletion	UNP P06213-2
F	?	-	SER	deletion	UNP P06213-2
F	?	-	SER	deletion	UNP P06213-2
F	?	-	THR	deletion	UNP P06213-2
F	?	-	SER	deletion	UNP P06213-2
F	?	-	VAL	deletion	UNP P06213-2
F	731	ALA	PRO	conflict	UNP P06213-2
F	732	GLY	THR	conflict	UNP P06213-2
F	733	ASN	SER	conflict	UNP P06213-2
F	734	ASN	PRO	conflict	UNP P06213-2
F	898	ARG	-	expression tag	UNP P06213-2
F	899	MET	-	expression tag	UNP P06213-2
F	900	LYS	-	expression tag	UNP P06213-2
F	901	GLN	-	expression tag	UNP P06213-2
F	902	LEU	-	expression tag	UNP P06213-2
F	903	GLU	-	expression tag	UNP P06213-2
F	904	ASP	-	expression tag	UNP P06213-2
F	905	LYS	-	expression tag	UNP P06213-2
F	906	VAL	-	expression tag	UNP P06213-2
F	907	GLU	-	expression tag	UNP P06213-2
F	908	GLU	-	expression tag	UNP P06213-2
F	909	LEU	-	expression tag	UNP P06213-2



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	910	LEU	-	expression tag	UNP P06213-2
F	911	SER	-	expression tag	UNP P06213-2
F	912	LYS	-	expression tag	UNP P06213-2
F	913	ASN	-	expression tag	UNP P06213-2
F	914	TYR	-	expression tag	UNP P06213-2
F	915	HIS	-	expression tag	UNP P06213-2
F	916	LEU	-	expression tag	UNP P06213-2
F	917	GLU	-	expression tag	UNP P06213-2
F	918	ASN	-	expression tag	UNP P06213-2
F	919	GLU	-	expression tag	UNP P06213-2
F	920	VAL	-	expression tag	UNP P06213-2
F	921	ALA	-	expression tag	UNP P06213-2
F	922	ARG	-	expression tag	UNP P06213-2
F	923	LEU	-	expression tag	UNP P06213-2
F	924	LYS	-	expression tag	UNP P06213-2
F	925	LYS	-	expression tag	UNP P06213-2
F	926	LEU	-	expression tag	UNP P06213-2
F	927	VAL	-	expression tag	UNP P06213-2
F	928	GLY	-	expression tag	UNP P06213-2
F	929	GLU	-	expression tag	UNP P06213-2
F	930	ARG	-	expression tag	UNP P06213-2

 \bullet Molecule 4 is a protein called IM462.

Mol	Chain	Residues		${f Atoms}$					Trace
4	G	19	Total	_		_		0	0
_		10	157	100	23	32	2		
1 4	П	19	Total	С	N	Ο	S	0	0
4	11	19	157	100	23	32	2	U	U

• Molecule 5 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-a cetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				AltConf	Trace
5	С	9	Total	С	N	О	0	0
)		2	28	16	2	10	U	U

• Molecule 6 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula:



 $\mathrm{C_8H_{15}NO_6}).$




Mol	Chain	Residues	A	ton	ns		AltConf
6	Е	1	Total	С	N	О	0
U	12	1	42	24	3	15	U
6	E	1	Total	С	N	O	0
0	<u> 1</u> 2	1	42	24	3	15	0
6	E	1	Total	С	N	Ο	0
0	<u> 1</u> 2	1	42	24	3	15	0
6	F	1	Total	С	N	О	0
0	I'	1	42	24	3	15	
6	F	1	Total	С	N	О	0
U	I.	1	42	24	3	15	U
6	F	1	Total	С	N	О	0
	I.	1	42	24	3	15	U




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Insulin A chain













4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	346000	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE	Depositor
	CORRECTION	
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{Å}^2)$	60	Depositor
Minimum defocus (nm)	500	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	0.167	Depositor
Minimum map value	-0.091	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.002	Depositor
Recommended contour level	0.037	Depositor
Map size (Å)	407.03992, 407.03992, 407.03992	wwPDB
Map dimensions	288, 288, 288	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.413333, 1.413333, 1.413333	Depositor



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, HY1

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain		Bond	lengths	Bond angles		
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.47	0/164	0.54	0/220	
2	В	0.46	0/205	0.53	0/276	
3	Е	0.44	0/4821	0.59	0/6532	
3	F	0.41	0/2565	0.61	0/3469	
4	G	0.35	0/151	0.44	0/204	
4	Н	0.44	0/151	0.55	0/204	
All	All	0.43	0/8057	0.59	0/10905	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	163	0	149	0	0
2	В	200	0	184	1	0
3	Ε	4708	0	4568	60	0
3	F	2507	0	2465	29	0
4	G	157	0	136	0	0
4	Н	157	0	136	6	0
5	С	28	0	25	0	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	$\mathbf{H}(\mathbf{added})$	Clashes	Symm-Clashes
6	Е	42	0	39	1	0
6	F	42	0	39	0	0
All	All	8004	0	7741	91	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 91 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$\begin{array}{c} \text{Clash} \\ \text{overlap } (\text{\AA}) \end{array}$
3:E:78:ASN:HA	3:E:108:ASN:HD22	1.44	0.83
3:E:211:GLU:OE2	3:E:230:ASN:ND2	2.17	0.77
3:E:80:THR:HA	3:E:109:LEU:HA	1.74	0.69
3:E:146:VAL:HG21	3:F:702:ARG:HH12	1.62	0.64
3:F:410:GLN:HB2	3:F:518:PHE:CD2	2.34	0.63

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	19/21 (90%)	19 (100%)	0	0	100 100
2	В	23/30 (77%)	22 (96%)	1 (4%)	0	100 100
3	E	578/930 (62%)	548 (95%)	29 (5%)	1 (0%)	47 82
3	F	302/930 (32%)	289 (96%)	13 (4%)	0	100 100
4	G	17/23~(74%)	16 (94%)	1 (6%)	0	100 100
4	Н	17/23 (74%)	17 (100%)	0	0	100 100
All	All	956/1957 (49%)	911 (95%)	44 (5%)	1 (0%)	54 85



All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type	
3	${ m E}$	193	PRO	

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Analysed Rotameric Outliers		Percentiles		
1	A	20/20 (100%)	20 (100%)	0	100	100	
2	В	21/26 (81%)	21 (100%)	0	100	100	
3	E	532/843 (63%)	525 (99%)	7 (1%)	69	89	
3	F	276/843 (33%)	272 (99%)	4 (1%)	67	88	
4	G	16/20 (80%)	16 (100%)	0	100	100	
4	Н	16/20 (80%)	16 (100%)	0	100	100	
All	All	881/1772 (50%)	870 (99%)	11 (1%)	72	90	

5 of 11 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	F	412	TRP
3	F	417	HIS
3	F	524	CYS
3	F	457	ILE
3	Е	295	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
3	F	561	GLN
3	F	462	ASN
3	Е	417	HIS
3	Ε	348	ASN
3	F	405	ASN



5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Tuno	Type Chain Res Link			Bo	Bond lengths			Bond angles		
IVIOI	Type	Chain	nes	Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2	
5	NAG	С	1	5,3	14,14,15	0.94	1 (7%)	17,19,21	0.77	1 (5%)	
5	NAG	С	2	5	14,14,15	0.32	0	17,19,21	1.20	1 (5%)	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	С	1	5,3	-	2/6/23/26	0/1/1/1
5	NAG	С	2	5	-	0/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	Observed(A)	Ideal(A)
5	С	1	NAG	O5-C1	-3.17	1.38	1.43

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	${f Z}$	$\operatorname{Observed}(^{o})$	$\operatorname{Ideal}({}^o)$
5	С	2	NAG	C1-O5-C5	4.83	118.74	112.19



Continued from previous page...

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
5	С	1	NAG	O4-C4-C5	-2.40	103.34	109.30

There are no chirality outliers.


All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	С	1	NAG	C4-C5-C6-O6
5	С	1	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry (i)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Вс	ond leng	ths	Bond angles		
MIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z >2
6	NAG	F	1002	3	14,14,15	0.71	1 (7%)	17,19,21	1.07	1 (5%)
6	NAG	F	1003	3	14,14,15	0.24	0	17,19,21	0.41	0
6	NAG	Е	1001	3	14,14,15	0.40	0	17,19,21	0.79	1 (5%)
6	NAG	F	1001	3	14,14,15	0.36	0	17,19,21	0.86	1 (5%)
6	NAG	E	1002	3	14,14,15	0.45	0	17,19,21	0.48	0
6	NAG	Е	1003	3	14,14,15	0.65	1 (7%)	17,19,21	0.75	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	NAG	F	1002	3	-	2/6/23/26	0/1/1/1
6	NAG	F	1003	3	-	0/6/23/26	0/1/1/1
6	NAG	E	1001	3	-	3/6/23/26	0/1/1/1
6	NAG	F	1001	3	-	2/6/23/26	0/1/1/1
6	NAG	Е	1002	3	-	3/6/23/26	0/1/1/1
6	NAG	E	1003	3	-	1/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(A)	$\operatorname{Ideal}(\text{\AA})$
6	F	1002	NAG	O5-C1	2.44	1.47	1.43
6	Ε	1003	NAG	C1-C2	2.21	1.55	1.52

All (4) bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
6	F	1002	NAG	C1-O5-C5	4.18	117.86	112.19
6	F	1001	NAG	C1-O5-C5	3.35	116.73	112.19
6	Е	1003	NAG	C1-O5-C5	2.53	115.61	112.19
6	Е	1001	NAG	C1-O5-C5	2.26	115.25	112.19

There are no chirality outliers.

5 of 11 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	Е	1002	NAG	C1-C2-N2-C7
6	Е	1001	NAG	O5-C5-C6-O6
6	Е	1001	NAG	C4-C5-C6-O6
6	F	1002	NAG	O5-C5-C6-O6
6	F	1002	NAG	C4-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes	
6	E	1002	NAG	1	0	

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.




6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-26364. These allow visual inspection of the internal detail of the map and identification of artifacts.

Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections (i)

6.1.1 Primary map



6.1.2 Raw map




The images above show the map projected in three orthogonal directions.




6.2 Central slices (i)

6.2.1 Primary map







Y Index: 144




Z Index: 144


6.2.2 Raw map



X Index: 144



Y Index: 144




Z Index: 144

The images above show central slices of the map in three orthogonal directions.




6.3 Largest variance slices (i)

6.3.1 Primary map



6.3.2 Raw map




The images above show the largest variance slices of the map in three orthogonal directions.




6.4 Orthogonal surface views (i)

6.4.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.037. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.4.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.

6.5 Mask visualisation (i)


This section was not generated. No masks/segmentation were deposited.



7 Map analysis (i)

This section contains the results of statistical analysis of the map.


7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



7.2 Volume estimate (i)




The volume at the recommended contour level is $36~\mathrm{nm^3}$; this corresponds to an approximate mass of $32~\mathrm{kDa}$.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



7.3 Rotationally averaged power spectrum (i)




^{*}Reported resolution corresponds to spatial frequency of 0.333 $\rm \mathring{A}^{-1}$



8 Fourier-Shell correlation (i)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC (i)



*Reported resolution corresponds to spatial frequency of 0.333 $\rm \mathring{A}^{-1}$



8.2 Resolution estimates (i)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)				
rtesolution estimate (A)	0.143	0.5	Half-bit		
Reported by author	3.00	-	-		
Author-provided FSC curve	-	-	-		
Unmasked-calculated*	3.72	6.33	3.78		


^{*}Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.72 differs from the reported value 3.0 by more than 10 %



9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-26364 and PDB model 7U6E. Per-residue inclusion information can be found in section 3 on page 8.


9.1 Map-model overlay (i)



The images above show the 3D surface view of the map at the recommended contour level 0.037 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.




9.2 Q-score mapped to coordinate model (i)



The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


9.3 Atom inclusion mapped to coordinate model (i)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.037).



9.4 Atom inclusion (i)



At the recommended contour level, 75% of all backbone atoms, 61% of all non-hydrogen atoms, are inside the map.



9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (0.037) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.6139	0.4800
A	0.5652	0.4850
В	0.7143	0.5040
С	0.6071	0.5350
Е	0.6452	0.4900
F	0.5739	0.4750
G	0.4183	0.3100
H	0.4314	0.3690



