



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 24, 2023 – 08:01 AM EDT

PDB ID : 5UCM  
Title : Crystal Structure of Prolyl-tRNA Synthetase from *Pseudomonas aeruginosa*  
Authors : Seattle Structural Genomics Center for Infectious Disease; Seattle Structural Genomics Center for Infectious Disease (SSGCID)  
Deposited on : 2016-12-22  
Resolution : 2.60 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	579	 88% 9% .
1	B	579	 % 84% 13% ..

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 9131 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Proline-tRNA ligase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	564	4283	2707	746	813	17	0	1	0
1	B	565	4358	2752	762	827	17	0	6	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	MET	-	initiating methionine	UNP Q9I502
A	-6	ALA	-	expression tag	UNP Q9I502
A	-5	HIS	-	expression tag	UNP Q9I502
A	-4	HIS	-	expression tag	UNP Q9I502
A	-3	HIS	-	expression tag	UNP Q9I502
A	-2	HIS	-	expression tag	UNP Q9I502
A	-1	HIS	-	expression tag	UNP Q9I502
A	0	HIS	-	expression tag	UNP Q9I502
B	-7	MET	-	initiating methionine	UNP Q9I502
B	-6	ALA	-	expression tag	UNP Q9I502
B	-5	HIS	-	expression tag	UNP Q9I502
B	-4	HIS	-	expression tag	UNP Q9I502
B	-3	HIS	-	expression tag	UNP Q9I502
B	-2	HIS	-	expression tag	UNP Q9I502
B	-1	HIS	-	expression tag	UNP Q9I502
B	0	HIS	-	expression tag	UNP Q9I502

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mg	0	0
			1	1		
2	B	1	Total	Mg	0	0
			1	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	226	Total 226	O 226	0	0
3	B	262	Total 262	O 262	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Proline-tRNA ligase



- Molecule 1: Proline-tRNA ligase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	85.83Å 101.54Å 187.24Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.00 – 2.60 49.00 – 2.60	Depositor EDS
% Data completeness (in resolution range)	92.4 (49.00-2.60) 91.5 (49.00-2.60)	Depositor EDS
$R_{merge}$	0.13	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.86 (at 2.61Å)	Xtrriage
Refinement program	PHENIX	Depositor
R, $R_{free}$	0.169 , 0.232 0.169 , 0.233	Depositor DCC
$R_{free}$ test set	2000 reflections (4.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	28.3	Xtrriage
Anisotropy	0.118	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 51.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	9131	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	35.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.12% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.37	0/4369	0.55	0/5931
1	B	0.39	0/4459	0.58	1/6047 (0.0%)
All	All	0.38	0/8828	0.57	1/11978 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	478	PRO	C-N-CA	5.11	134.48	121.70

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4283	0	4182	36	0
1	B	4358	0	4300	47	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	226	0	0	3	0
3	B	262	0	0	3	0
All	All	9131	0	8482	79	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (79) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:257:THR:HB	1:A:260:THR:HB	1.60	0.83
1:A:123:ASN:HD22	1:A:123:ASN:H	1.35	0.74
1:A:250:GLU:O	1:A:375:ARG:NH2	2.25	0.70
1:A:67:LEU:HD11	1:B:5:GLN:HB3	1.78	0.65
1:A:239:VAL:HG23	1:A:393:PRO:HG2	1.79	0.64
1:A:2:ARG:NH2	1:A:494:TYR:OH	2.31	0.64
1:B:239:VAL:HG23	1:B:393:PRO:HG2	1.79	0.64
1:B:525:ILE:HG13	1:B:527:ILE:HG13	1.81	0.62
1:B:478:PRO:HA	1:B:479:LEU:HB2	1.81	0.61
1:A:121:GLU:HA	1:B:9:SER:HB2	1.85	0.57
1:A:373:TRP:CZ3	1:A:380:PRO:HD3	2.41	0.56
1:B:415:LEU:HD22	1:B:441[A]:CYS:SG	2.46	0.56
1:B:254:LEU:HD13	1:B:366:LYS:HD3	1.89	0.54
1:A:92:GLU:HG2	1:A:140:ARG:NH2	2.23	0.53
1:A:155:PHE:HE2	1:A:157:MET:HE3	1.73	0.53
1:B:2:ARG:HB3	1:B:505:LEU:HD12	1.91	0.53
1:A:528:PRO:HB2	1:A:529:HIS:ND1	2.25	0.52
1:B:479:LEU:HD23	1:B:534:SER:HB3	1.92	0.51
1:A:155:PHE:CE2	1:A:157:MET:HE3	2.44	0.51
1:A:310:GLN:O	3:A:701:HOH:O	2.19	0.51
1:B:193[A]:ARG:HD2	3:B:839:HOH:O	2.11	0.51
1:B:155:PHE:HE2	1:B:157:MET:HE3	1.76	0.50
1:A:375:ARG:HD2	1:A:376:ASP:OD1	2.12	0.50
1:B:92:GLU:OE2	1:B:140:ARG:NH2	2.45	0.49
1:B:347:VAL:HG13	1:B:380:PRO:HG2	1.94	0.49
1:B:188:LEU:HD22	1:B:469:LEU:HD21	1.94	0.49
1:B:170:LEU:HD21	1:B:414:GLN:HB3	1.94	0.49
1:B:155:PHE:CE2	1:B:157:MET:HE3	2.48	0.49
1:B:372:ASN:HD22	1:B:375:ARG:NH2	2.11	0.49
1:B:92:GLU:OE1	1:B:140:ARG:NE	2.47	0.48
1:B:478:PRO:HG3	1:B:490:THR:HG21	1.96	0.48
1:B:483:THR:HB	1:B:486:VAL:HG12	1.96	0.48
1:A:47:LEU:HD12	1:A:50:LEU:HD23	1.95	0.48
1:B:564:PHE:CZ	1:B:568:LYS:HE2	2.49	0.47
1:A:121:GLU:OE2	1:B:33:ARG:NH1	2.45	0.47
1:B:119:ARG:HD3	1:B:423:MET:O	2.15	0.46
1:A:456:GLN:HG2	1:A:457:ASN:ND2	2.30	0.46

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:257:THR:HB	1:B:260:THR:HB	1.98	0.46
1:A:29:ALA:HB1	1:A:525:ILE:HG22	1.97	0.46
1:A:415:LEU:HD21	1:A:419:TYR:HE2	1.80	0.45
1:A:529:HIS:HD2	1:A:545:TYR:OH	1.99	0.45
1:B:118:ALA:HB1	1:B:122:LEU:HD12	1.99	0.45
1:B:277:THR:OG1	1:B:298:GLY:HA3	2.16	0.45
1:B:543:LEU:HG	1:B:558:ILE:HG12	1.97	0.45
1:A:123:ASN:HD22	1:A:123:ASN:N	2.07	0.45
1:B:425:LEU:HD21	1:B:437:LEU:HD12	1.98	0.45
1:B:313:VAL:O	3:B:701:HOH:O	2.21	0.44
1:A:241:ARG:HG2	1:A:393:PRO:HD3	1.98	0.44
1:A:565:ILE:HD12	1:A:565:ILE:HA	1.87	0.44
1:B:2:ARG:HD2	1:B:494:TYR:OH	2.16	0.44
1:B:43:LEU:HD23	1:B:43:LEU:HA	1.78	0.44
1:B:473:GLN:HE21	1:B:569:LEU:HD22	1.83	0.44
1:B:47:LEU:HD12	1:B:50:LEU:HD23	2.00	0.43
1:A:33:ARG:NH2	1:B:121:GLU:OE1	2.51	0.43
1:A:273:PRO:HG2	1:A:276:LYS:HG2	2.00	0.43
1:B:235:LYS:HE2	3:B:797:HOH:O	2.17	0.43
1:A:119:ARG:HD3	1:A:423:MET:O	2.19	0.43
1:A:208:HIS:O	1:A:411:HIS:HD2	2.02	0.43
1:B:239:VAL:CG2	1:B:393:PRO:HG2	2.49	0.43
1:B:242:GLU:CD	1:B:242:GLU:H	2.22	0.43
1:B:251:ASP:O	1:B:253[B]:ARG:NH2	2.51	0.43
1:B:534:SER:O	1:B:538:LEU:HG	2.19	0.43
1:B:3:THR:OG1	1:B:507:ASP:HA	2.19	0.42
1:B:394:SER:HA	1:B:401:LEU:HD21	2.02	0.42
1:B:37:SER:OG	1:B:145:PRO:HG2	2.19	0.41
1:B:158:LYS:O	1:B:443:GLY:HA2	2.20	0.41
1:B:379:LEU:HD12	1:B:380:PRO:HD2	2.01	0.41
1:A:123:ASN:ND2	1:A:127:GLN:OE1	2.54	0.41
1:B:3:THR:HB	1:B:7:LEU:HD22	2.03	0.41
1:A:68:GLU:HA	1:A:133:TYR:O	2.21	0.41
1:A:346:ILE:HG22	1:A:385:LEU:HD11	2.03	0.41
1:A:375:ARG:NH1	1:A:376:ASP:OD1	2.54	0.41
1:A:420:SER:HA	1:A:425:LEU:HB3	2.03	0.41
1:A:448:ARG:HH11	1:A:448:ARG:HD3	1.69	0.41
1:B:33:ARG:CZ	1:B:33:ARG:HB2	2.52	0.40
1:A:221:ILE:HG13	1:A:223:PHE:CE1	2.56	0.40
1:A:301:GLU:HG3	3:A:875:HOH:O	2.21	0.40
1:A:313:VAL:O	3:A:701:HOH:O	2.21	0.40

There are no symmetry-related clashes.

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	561/579 (97%)	545 (97%)	16 (3%)	0	100	100
1	B	567/579 (98%)	551 (97%)	16 (3%)	0	100	100
All	All	1128/1158 (97%)	1096 (97%)	32 (3%)	0	100	100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	442/483 (92%)	435 (98%)	7 (2%)	62	82
1	B	457/483 (95%)	448 (98%)	9 (2%)	55	78
All	All	899/966 (93%)	883 (98%)	16 (2%)	59	80

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	123	ASN
1	A	158	LYS
1	A	254	LEU
1	A	456	GLN
1	A	536	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	542	VAL
1	A	556	LEU
1	B	45	MET
1	B	124	SER
1	B	158	LYS
1	B	205	SER
1	B	485	SER
1	B	542	VAL
1	B	551	SER
1	B	558	ILE
1	B	566	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	123	ASN
1	A	457	ASN
1	A	529	HIS
1	B	372	ASN
1	B	529	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	564/579 (97%)	-0.57	1 (0%) 95   95	18, 34, 69, 91	0
1	B	565/579 (97%)	-0.65	3 (0%) 91   89	18, 29, 65, 106	0
All	All	1129/1158 (97%)	-0.61	4 (0%) 92   91	18, 31, 67, 106	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	542	VAL	2.7
1	B	16	ALA	2.5
1	B	1	MET	2.2
1	A	8	LEU	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	MG	B	600	1/1	0.90	0.16	52,52,52,52	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	MG	A	600	1/1	0.97	0.15	44,44,44,44	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.