



wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 24, 2023 – 06:31 AM EDT

PDB ID : 5UEM
Title : Crystal structure of 354NC37 Fab in complex with HIV-1 clade AE strain 93TH057 gp120
Authors : Sievers, S.A.; Gristick, H.B.; Bjorkman, P.J.
Deposited on : 2017-01-02
Resolution : 2.70 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	G	361	<div style="display: flex; align-items: center;"> <div style="width: 9%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 85%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 10%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 5%; height: 10px; background-color: grey;"></div> </div>
2	H	230	<div style="display: flex; align-items: center;"> <div style="width: 5%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 84%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 15%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 0%; height: 10px; background-color: grey;"></div> </div>
3	L	213	<div style="display: flex; align-items: center;"> <div style="width: 0%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 90%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 10%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 0%; height: 10px; background-color: grey;"></div> </div>
4	A	2	<div style="display: flex; align-items: center;"> <div style="width: 100%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 0%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 0%; height: 10px; background-color: grey;"></div> </div>
5	B	5	<div style="display: flex; align-items: center;"> <div style="width: 80%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 20%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 0%; height: 10px; background-color: grey;"></div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
6	C	7	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	NAG	A	2	-	-	-	X
5	MAN	B	4	-	-	-	X
5	MAN	B	5	-	-	-	X
6	MAN	C	7	-	-	-	X
7	NAG	G	603	-	-	-	X

2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 6296 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called clade A/E 93TH057 HIV-1 gp120 core.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	G	342	2637	1658	456	499	24	0	0	1

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
G	65	CYS	VAL	conflict	UNP A0A0M3KKW9
G	115	CYS	SER	conflict	UNP A0A0M3KKW9
G	493	GLY	-	expression tag	UNP A0A0M3KKW9
G	494	SER	-	expression tag	UNP A0A0M3KKW9
G	495	HIS	-	expression tag	UNP A0A0M3KKW9
G	496	HIS	-	expression tag	UNP A0A0M3KKW9
G	497	HIS	-	expression tag	UNP A0A0M3KKW9
G	498	HIS	-	expression tag	UNP A0A0M3KKW9
G	499	HIS	-	expression tag	UNP A0A0M3KKW9
G	500	HIS	-	expression tag	UNP A0A0M3KKW9

- Molecule 2 is a protein called 354NC37 Fab Heavy Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	H	230	1719	1084	296	331	8	0	0	0

- Molecule 3 is a protein called 354NC37 Fab Light Chain.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	L	213	1599	1007	274	314	4	0	0	0

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	A	2	28	16	2	10	0	0	0

- Molecule 5 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
5	B	5	61	34	2	25	0	0	0

- Molecule 6 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
6	C	7	83	46	2	35	0	0	0

- Molecule 7 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
7	G	1	Total 14	8	1	5	0	0
7	G	1	Total 14	8	1	5	0	0
7	G	1	Total 14	8	1	5	0	0
7	G	1	Total 14	8	1	5	0	0
7	G	1	Total 14	8	1	5	0	0
7	G	1	Total 14	8	1	5	0	0

- Molecule 8 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (three-letter code: EPE) (formula: C₈H₁₈N₂O₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
8	G	1	15	8	2	4	1	0	0


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	G	31	Total	O	0	0
			31	31		
9	H	22	Total	O	0	0
			22	22		
9	L	17	Total	O	0	0
			17	17		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: clade A/E 93TH057 HIV-1 gp120 core



- Molecule 2: 354NC37 Fab Heavy Chain



- Molecule 3: 354NC37 Fab Light Chain




- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



MAG1
MAG2

- Molecule 5: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain B:  80% 20%

MAG1
MAG2
BMA3
MAN4
MAN5

- Molecule 6: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C:  43% 57%

MAG1
MAG2
BMA3
MAN4
MAN5
MAN6
MAN7

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	63.63Å 67.51Å 210.66Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	35.74 – 2.70 35.74 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.7 (35.74-2.70) 99.7 (35.74-2.70)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.54 (at 2.68Å)	Xtrriage
Refinement program	PHENIX 1.12_2829	Depositor
R, R_{free}	0.220 , 0.266 0.220 , 0.266	Depositor DCC
R_{free} test set	1431 reflections (5.57%)	wwPDB-VP
Wilson B-factor (Å ²)	42.3	Xtrriage
Anisotropy	0.403	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 33.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	6296	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.24% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EPE, NAG, MAN, BMA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	G	0.25	0/2693	0.44	0/3660
2	H	0.26	0/1765	0.48	0/2409
3	L	0.25	0/1638	0.45	0/2233
All	All	0.26	0/6096	0.45	0/8302

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	G	2637	0	2535	21	0
2	H	1719	0	1661	20	0
3	L	1599	0	1541	13	0
4	A	28	0	25	0	0
5	B	61	0	52	0	0
6	C	83	0	70	0	0
7	G	84	0	78	1	0
8	G	15	0	17	0	0
9	G	31	0	0	0	0
9	H	22	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
9	L	17	0	0	0	0
All	All	6296	0	5979	52	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 52 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:101:VAL:HG21	1:G:480:ARG:HG2	1.71	0.73
1:G:124:GLY:H	2:H:74:HIS:HA	1.59	0.67
2:H:127:SER:H	2:H:130:SER:HB3	1.59	0.66
3:L:145:LYS:HB3	3:L:197:THR:HB	1.78	0.65
1:G:275:GLU:OE2	1:G:282:LYS:NZ	2.34	0.60

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	G	336/361 (93%)	324 (96%)	12 (4%)	0	100	100
2	H	228/230 (99%)	217 (95%)	11 (5%)	0	100	100
3	L	211/213 (99%)	201 (95%)	10 (5%)	0	100	100
All	All	775/804 (96%)	742 (96%)	33 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	G	293/318 (92%)	293 (100%)	0	100	100
2	H	186/190 (98%)	180 (97%)	6 (3%)	39	68
3	L	175/184 (95%)	173 (99%)	2 (1%)	73	90
All	All	654/692 (94%)	646 (99%)	8 (1%)	71	88

5 of 8 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	L	28	LEU
3	L	24	ARG
2	H	100(B)	VAL
2	H	82(B)	ARG
2	H	183	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	G	425	ASN
1	G	444	ASN
3	L	90	ASN
3	L	124	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates i

14 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	A	1	1,4	14,14,15	0.31	0	17,19,21	0.73	0
4	NAG	A	2	4	14,14,15	0.28	0	17,19,21	0.68	0
5	NAG	B	1	1,5	14,14,15	0.32	0	17,19,21	0.77	0
5	NAG	B	2	5	14,14,15	0.30	0	17,19,21	0.70	0
5	BMA	B	3	5	11,11,12	0.29	0	15,15,17	1.09	1 (6%)
5	MAN	B	4	5	11,11,12	0.26	0	15,15,17	0.86	0
5	MAN	B	5	5	11,11,12	0.28	0	15,15,17	0.72	0
6	NAG	C	1	1,6	14,14,15	0.31	0	17,19,21	0.81	1 (5%)
6	NAG	C	2	6	14,14,15	0.31	0	17,19,21	0.96	2 (11%)
6	BMA	C	3	6	11,11,12	0.24	0	15,15,17	0.83	0
6	MAN	C	4	6	11,11,12	0.27	0	15,15,17	0.82	0
6	MAN	C	5	6	11,11,12	0.27	0	15,15,17	0.86	0
6	MAN	C	6	6	11,11,12	0.30	0	15,15,17	1.22	1 (6%)
6	MAN	C	7	6	11,11,12	0.33	0	15,15,17	0.77	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	A	2	4	-	2/6/23/26	0/1/1/1
5	NAG	B	1	1,5	-	2/6/23/26	0/1/1/1
5	NAG	B	2	5	-	0/6/23/26	0/1/1/1
5	BMA	B	3	5	-	1/2/19/22	0/1/1/1
5	MAN	B	4	5	-	2/2/19/22	0/1/1/1
5	MAN	B	5	5	-	1/2/19/22	0/1/1/1
6	NAG	C	1	1,6	-	0/6/23/26	0/1/1/1
6	NAG	C	2	6	-	0/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	BMA	C	3	6	-	1/2/19/22	0/1/1/1
6	MAN	C	4	6	-	0/2/19/22	0/1/1/1
6	MAN	C	5	6	-	2/2/19/22	0/1/1/1
6	MAN	C	6	6	-	1/2/19/22	0/1/1/1
6	MAN	C	7	6	-	1/2/19/22	0/1/1/1

There are no bond length outliers.

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	C	6	MAN	O3-C3-C2	-3.24	103.78	109.99
5	B	3	BMA	O3-C3-C2	-2.48	105.24	109.99
6	C	1	NAG	C1-O5-C5	2.23	115.21	112.19
6	C	7	MAN	O5-C1-C2	-2.22	107.35	110.77
6	C	2	NAG	O5-C1-C2	-2.20	107.82	111.29

There are no chirality outliers.


5 of 13 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	2	NAG	C8-C7-N2-C2
4	A	2	NAG	O7-C7-N2-C2
5	B	4	MAN	O5-C5-C6-O6
5	B	4	MAN	C4-C5-C6-O6
6	C	5	MAN	O5-C5-C6-O6


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.







5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	NAG	G	618	1	14,14,15	0.31	0	17,19,21	0.80	1 (5%)
7	NAG	G	609	1	14,14,15	0.30	0	17,19,21	0.82	0
8	EPE	G	621	-	15,15,15	0.76	1 (6%)	18,20,20	1.92	5 (27%)
7	NAG	G	620	1	14,14,15	0.26	0	17,19,21	0.84	0
7	NAG	G	619	1	14,14,15	0.31	0	17,19,21	0.74	0
7	NAG	G	603	1	14,14,15	0.30	0	17,19,21	0.70	0
7	NAG	G	617	1	14,14,15	0.32	0	17,19,21	0.81	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	NAG	G	618	1	-	0/6/23/26	0/1/1/1
7	NAG	G	609	1	-	2/6/23/26	0/1/1/1
8	EPE	G	621	-	-	6/9/19/19	0/1/1/1
7	NAG	G	620	1	-	0/6/23/26	0/1/1/1
7	NAG	G	619	1	-	0/6/23/26	0/1/1/1
7	NAG	G	603	1	-	0/6/23/26	0/1/1/1
7	NAG	G	617	1	-	0/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	G	621	EPE	C10-S	2.44	1.81	1.77

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	G	621	EPE	C5-N4-C3	4.26	118.41	108.83
8	G	621	EPE	C7-N4-C5	3.80	120.94	111.23
8	G	621	EPE	C7-N4-C3	3.37	119.84	111.23
8	G	621	EPE	O3S-S-C10	2.30	109.49	105.77
8	G	621	EPE	O1S-S-C10	2.20	109.56	106.92

There are no chirality outliers.

5 of 8 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
8	G	621	EPE	C8-C7-N4-C5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
8	G	621	EPE	C9-C10-S-O1S
8	G	621	EPE	C9-C10-S-O2S
8	G	621	EPE	C9-C10-S-O3S
7	G	609	NAG	O5-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	G	617	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	G	342/361 (94%)	0.46	31 (9%) 9 7	28, 38, 59, 76	0
2	H	230/230 (100%)	0.24	11 (4%) 30 28	23, 36, 59, 71	0
3	L	213/213 (100%)	0.10	3 (1%) 75 77	22, 36, 53, 60	0
All	All	785/804 (97%)	0.30	45 (5%) 23 22	22, 37, 58, 76	0

The worst 5 of 45 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	462	ASN	7.7
1	G	461	ASN	5.3
1	G	460	ALA	4.3
2	H	128	SER	3.7
1	G	87	GLU	3.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	MAN	C	7	11/12	0.50	0.42	59,59,59,59	0
5	MAN	B	5	11/12	0.66	0.58	94,94,94,94	0
5	MAN	B	4	11/12	0.73	0.49	66,66,66,66	0


Continued on next page...


Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	NAG	A	2	14/15	0.74	0.41	49,59,68,72	14
5	BMA	B	3	11/12	0.79	0.27	53,70,85,87	0
6	MAN	C	5	11/12	0.80	0.28	46,46,46,46	0
6	MAN	C	6	11/12	0.84	0.40	56,56,56,56	0
5	NAG	B	2	14/15	0.84	0.19	42,58,79,84	0
5	NAG	B	1	14/15	0.86	0.21	32,42,60,65	0
6	NAG	C	2	14/15	0.87	0.28	39,50,73,77	14
4	NAG	A	1	14/15	0.88	0.21	23,50,61,71	0
6	MAN	C	4	11/12	0.89	0.23	51,51,51,51	0
6	BMA	C	3	11/12	0.89	0.20	32,49,56,62	11
6	NAG	C	1	14/15	0.90	0.20	37,50,57,62	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.







6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
7	NAG	G	603	14/15	0.73	0.55	66,66,66,66	0
7	NAG	G	620	14/15	0.77	0.30	40,45,50,52	14
7	NAG	G	617	14/15	0.83	0.35	43,59,74,75	0
7	NAG	G	619	14/15	0.84	0.40	54,60,65,67	14
7	NAG	G	618	14/15	0.89	0.22	35,54,75,82	0
7	NAG	G	609	14/15	0.90	0.17	32,37,49,50	0
8	EPE	G	621	15/15	0.96	0.17	25,30,45,51	0

6.5 Other polymers [i](#)

There are no such residues in this entry.