



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 5, 2024 – 12:16 PM EDT

PDB ID : 8UFY
Title : High Resolution structure of A. viridans Lactate Oxidase
Authors : Lodowski, D.T.
Deposited on : 2023-10-05
Resolution : 1.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.37.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1223 (1.22-1.18)
Clashscore	141614	1286 (1.22-1.18)
Ramachandran outliers	138981	1240 (1.22-1.18)
Sidechain outliers	138945	1239 (1.22-1.18)
RSRZ outliers	127900	1200 (1.22-1.18)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	380	
1	B	380	

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 12267 atoms, of which 5866 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called L-lactate oxidase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	358	Total	C	H	N	O	S	0	51	0
			6063	1928	3005	537	586	7			
1	B	347	Total	C	H	N	O	S	0	38	0
			5673	1816	2796	495	559	7			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	HIS	-	expression tag	UNP Q44467
A	-4	HIS	-	expression tag	UNP Q44467
A	-3	HIS	-	expression tag	UNP Q44467
A	-2	HIS	-	expression tag	UNP Q44467
A	-1	HIS	-	expression tag	UNP Q44467
A	0	HIS	-	expression tag	UNP Q44467
A	232	ALA	GLY	conflict	UNP Q44467
A	255	ALA	ARG	conflict	UNP Q44467
B	-5	HIS	-	expression tag	UNP Q44467
B	-4	HIS	-	expression tag	UNP Q44467
B	-3	HIS	-	expression tag	UNP Q44467
B	-2	HIS	-	expression tag	UNP Q44467
B	-1	HIS	-	expression tag	UNP Q44467
B	0	HIS	-	expression tag	UNP Q44467
B	232	ALA	GLY	conflict	UNP Q44467
B	255	ALA	ARG	conflict	UNP Q44467

- Molecule 2 is FLAVIN MONONUCLEOTIDE (three-letter code: FMN) (formula: C₁₇H₂₁N₄O₉P) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
2	A	1	Total	C	H	N	O	P	0	0
			50	17	19	4	9	1		
2	B	1	Total	C	H	N	O	P	0	0
			50	17	19	4	9	1		

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
3	A	1	Total	C	H	O	0	0
			10	2	6	2		
3	A	1	Total	C	H	O	0	0
			10	2	6	2		

- Molecule 4 is LACTIC ACID (three-letter code: LAC) (formula: $C_3H_6O_3$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
4	A	1	11	3	5	3	0	1
4	B	1	10	3	4	3	0	1

- Molecule 5 is PYRUVIC ACID (three-letter code: PYR) (formula: $C_3H_4O_3$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	H	O	0	1
			9	3	3	3		
5	B	1	Total	C	H	O	0	1
			9	3	3	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	183	Total	O	0	0
			183	183		
6	B	189	Total	O	0	0
			189	189		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: L-lactate oxidase



- Molecule 1: L-lactate oxidase



4 Data and refinement statistics i

Property	Value	Source
Space group	I 4	Depositor
Cell constants a, b, c, α , β , γ	132.26Å 132.26Å 91.68Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.70 – 1.20 49.70 – 1.17	Depositor EDS
% Data completeness (in resolution range)	98.6 (49.70-1.20) 99.9 (49.70-1.17)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.51 (at 1.17Å)	Xtrriage
Refinement program	PHENIX 1.21rc1_5084	Depositor
R, R_{free}	0.146 , 0.162 0.138 , 0.156	Depositor DCC
R_{free} test set	13436 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	13.3	Xtrriage
Anisotropy	0.007	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.46 , 48.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.016 for -k,-h,-l	Xtrriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	12267	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.98% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FMN, EDO, PYR, LAC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.30	0/3255	0.59	0/4401
1	B	0.31	0/3042	0.59	0/4121
All	All	0.31	0/6297	0.59	0/8522

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3058	3005	2857	10	0
1	B	2877	2796	2665	6	0
2	A	31	19	19	0	0
2	B	31	19	19	1	0
3	A	8	12	12	0	0
4	A	6	5	0	1	0
4	B	6	4	0	1	0
5	A	6	3	0	0	0
5	B	6	3	0	0	0
6	A	183	0	0	1	0
6	B	189	0	0	0	0
All	All	6401	5866	5572	15	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 15 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:143[A]:PHE:CD1	1:B:166:ALA:HB2	2.35	0.61
1:A:198[A]:VAL:HG12	1:A:202:LEU:HD12	1.84	0.60
1:A:198[A]:VAL:CG1	1:A:202:LEU:HD12	2.35	0.57
1:B:265:HIS:NE2	4:B:403[A]:LAC:OXT	2.44	0.49
1:B:268[A]:ARG:HD2	2:B:401:FMN:C8M	2.43	0.48

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	403/380 (106%)	395 (98%)	7 (2%)	1 (0%)	47	19
1	B	380/380 (100%)	372 (98%)	7 (2%)	1 (0%)	41	16
All	All	783/760 (103%)	767 (98%)	14 (2%)	2 (0%)	41	16

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	297	SER
1	B	297	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	331/303 (109%)	329 (99%)	2 (1%)	86	63
1	B	310/303 (102%)	308 (99%)	2 (1%)	86	63
All	All	641/606 (106%)	637 (99%)	4 (1%)	84	63

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	141	ARG
1	A	190	VAL
1	B	141	ARG
1	B	149	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	LAC	A	404[A]	-	5,5,5	1.11	0	4,6,6	0.84	0
5	PYR	B	402[B]	-	5,5,5	1.07	1 (20%)	3,6,6	1.51	1 (33%)
4	LAC	B	403[A]	-	5,5,5	1.25	1 (20%)	4,6,6	0.62	0
3	EDO	A	402	-	3,3,3	0.28	0	2,2,2	0.12	0
2	FMN	A	401	-	33,33,33	1.02	2 (6%)	48,50,50	1.11	3 (6%)
5	PYR	A	405[B]	-	5,5,5	1.07	1 (20%)	3,6,6	1.58	1 (33%)
2	FMN	B	401	-	33,33,33	1.02	2 (6%)	48,50,50	1.09	4 (8%)
3	EDO	A	403	-	3,3,3	0.29	0	2,2,2	0.29	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	LAC	A	404[A]	-	-	2/4/4/4	-
5	PYR	B	402[B]	-	-	2/4/4/4	-
4	LAC	B	403[A]	-	-	2/4/4/4	-
3	EDO	A	402	-	-	1/1/1/1	-
2	FMN	A	401	-	-	1/18/18/18	0/3/3/3
5	PYR	A	405[B]	-	-	2/4/4/4	-
2	FMN	B	401	-	-	1/18/18/18	0/3/3/3
3	EDO	A	403	-	-	0/1/1/1	-

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	FMN	C4A-N5	3.94	1.38	1.30
2	B	401	FMN	C4A-N5	3.87	1.38	1.30
2	B	401	FMN	C10-N1	2.20	1.37	1.33
2	A	401	FMN	C10-N1	2.08	1.37	1.33
5	A	405[B]	PYR	OXT-C	-2.06	1.24	1.30

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	FMN	C4A-C10-N10	3.01	120.88	116.48
2	B	401	FMN	C4A-C10-N10	2.97	120.83	116.48
5	A	405[B]	PYR	OXT-C-O	2.48	129.27	123.61
2	A	401	FMN	C4-N3-C2	-2.41	121.20	125.64

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	402[B]	PYR	OXT-C-O	2.37	129.04	123.61

There are no chirality outliers.

5 of 11 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	A	405[B]	PYR	O-C-CA-CB
5	A	405[B]	PYR	OXT-C-CA-CB
5	B	402[B]	PYR	O-C-CA-CB
5	B	402[B]	PYR	OXT-C-CA-CB
4	A	404[A]	LAC	OXT-C-CA-OHN


There are no ring outliers.


3 monomers are involved in 3 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	404[A]	LAC	1	0
4	B	403[A]	LAC	1	0
2	B	401	FMN	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	358/380 (94%)	0.11	22 (6%) 21 20	10, 18, 37, 71	0
1	B	347/380 (91%)	0.42	42 (12%) 4 5	10, 16, 44, 64	0
All	All	705/760 (92%)	0.26	64 (9%) 9 9	10, 17, 41, 71	0

The worst 5 of 64 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	198	VAL	14.5
1	A	214	ILE	10.7
1	B	157	ILE	10.1
1	A	215	TYR	9.3
1	B	197	ILE	8.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	PYR	A	405[B]	6/6	0.69	0.29	28,30,35,35	9
4	LAC	B	403[A]	6/6	0.78	0.25	20,27,32,33	10
5	PYR	B	402[B]	6/6	0.80	0.23	50,52,60,60	9
4	LAC	A	404[A]	6/6	0.83	0.14	32,41,51,51	11
3	EDO	A	402	4/4	0.85	0.20	42,50,55,57	0
3	EDO	A	403	4/4	0.94	0.22	29,35,38,40	0
2	FMN	A	401	31/31	0.98	0.08	11,14,24,25	0
2	FMN	B	401	31/31	0.98	0.07	10,13,23,25	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around LAC B 403 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around PYR B 402 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around LAC A 404 (A):


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around FMN A 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.