



wwPDB EM Validation Summary Report ⓘ

Apr 30, 2024 – 09:21 pm BST

PDB ID : 4UUD
EMDB ID : EMD-2701
Title : Human dynamin 1 K44A superconstricted polymer stabilized with GTP
Authors : Sundborger, A.C.; Fang, S.; Heymann, J.A.; Ray, P.; Chappie, J.S.; Hinshaw, J.E.
Deposited on : 2014-07-25
Resolution : 12.50 Å (reported)
Based on initial models : 3ZYC, 1DYN, 3SNH

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev92
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 12.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.







Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	864	
1	D	864	
1	G	864	
1	K	864	
2	B	864	
2	C	864	
2	E	864	
2	F	864	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	H	864	 7% . . 87%
2	I	864	 8% 10% . . 76%
2	J	864	 7% 10% . . 76%
2	L	864	 7% . . . 87%

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 20992 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DYNAMIN-1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	329	2567	1615	453	489	10	0	0
1	D	337	2643	1664	466	503	10	0	0
1	G	329	2567	1615	453	489	10	0	0
1	K	337	2643	1664	466	503	10	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	744	ASN	ASP	variant	UNP Q05193
D	744	ASN	ASP	variant	UNP Q05193
G	744	ASN	ASP	variant	UNP Q05193
K	744	ASN	ASP	variant	UNP Q05193

- Molecule 2 is a protein called DYNAMIN-1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	B	204	1697	1079	297	307	14	0	0
2	C	113	946	609	158	175	4	0	0
2	E	204	1697	1079	297	307	14	0	0
2	F	113	946	609	158	175	4	0	0
2	H	113	946	609	158	175	4	0	0
2	I	204	1697	1079	297	307	14	0	0
2	J	204	1697	1079	297	307	14	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	L	113	946	609	158	175	4	0	0

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: DYNAMIN-1




ARG
SER
GLY
GLN
ALA
GLY
SER
PRO
PRO
SER
ARG
ARG
PRO
PRO
PHE
ASP
LEU

● Molecule 1: DYNAMIN-1



● Molecule 1: DYNAMIN-1



F83	L187	L285	GLU
L84	K188	T286	LEU
H85	V189	L289	LEU
C86	A190	R290	GLY
K87	K191	L296	ILE
G88	E192	L300	ASP
K89	V193	Q301	ILE
K90	P195	I307	ASN
F94	I201	E308	THR
V97	T205	K309	GLY
R98	K206	E310	GLU
I101	L207	V311	PHE
T105	D208	GLU	PRO
D106	L209	GLU	ILE
R107	R217	TYR	THR
I115	K223	LYS	VAL
P119	R228	ASN	LEU
H120	I232	ASN	GLY
N121	G233	ASN	THR
R123	V234	PRO	ILE
V124	V235	ALA	LEU
L130	V236	ALA	TYR
T133	R237	ASN	PRO
L134	S238	PRO	VAL
V135	Q239	LEU	GLN
D136	D241	ALA	GLU
L137	I242	LEU	LYS
P138	D243	LEU	TYR
G139	G244	GLN	ARG
M140	K245	MET	ILE
Q148	E255	VAL	ARG
M159	F259	GLN	ASN
L160	H262	GLU	ASN
M161	P263	GLU	GLU
V164	S264	GLU	ILE
E167	Y265	LYS	ARG
M168	D270	ILE	THR
G169	R271	GLY	PRO
L170	M272	GLU	ASP
I171	G273	SER	GLY
L172	T274	GLY	ALA
A173	P275	ASP	GLY
V174	Y276	GLN	PHE
S175	L277	ILE	THR
P176	V280	ASP	ILE
A177	L281	TYR	GLN

● Molecule 1: DYNAMIN-1



MET	V97	Q198	R297
GLY	R98	I201	L304
ASN	L99	I204	I307
ARG	E100	T205	E310
GLY	I101	K206	V311
M6	K113	L207	E312
I10	V118	D208	E313
P11	P119	L209	Y314
L12	I120	D218	K315
V13	M121	V219	K316
L16	L122	L220	F317
Q17	R123	E221	R318
L23	L130	N222	F319
G24	V135	K223	ASP
Q25	V136	L224	ASP
L29	D136	L224	PRO
D30	L137	L224	ALA
L31	P296	L224	ALA
P32	R140	L227	ARG
Q33	T141	R228	LYS
I34	K142	R229	LYS
K35	V145	I232	ALA
V36	P149	G233	LEU
Q40	P150	V234	LEU
A42	D151	V235	GLN
G43	I152	N236	VAL
L48	I157	R237	GLN
D55	L160	I242	PHE
F56	M161	G244	ALA
L57	F163	K245	VAL
P58	Q162	I248	VAL
R59	V164	A251	ASP
G60	L170	A254	PHE
S61	I171	E255	ALA
G62	I172	F258	ALA
T63	A173	F259	ALA
V64	V174	L260	GLY
T65	M178	D270	GLN
R66	S179	R271	ILE
R67	L181	T274	ASP
V74	S184	P275	THR
F78	A190	L277	TYR
E79	K191	V280	TYR
Y80	E192	L281	GLU
F83	C86	I289	LEU
L84	R87		SER
H85			GLY
C86			GLY
K87			ALA
			ARG
			ILE

4 Experimental information

Property	Value	Source
EM reconstruction method	HELICAL	Depositor
Imposed symmetry	HELICAL, twist=Not provided°, rise=Not provided Å, axial sym=Not provided	Depositor
Number of segments used	7525	Depositor
Resolution determination method	Not provided	
CTF correction method	INDIVIDUAL IMAGES	Depositor
Microscope	FEI/PHILIPS CM300FEG/HE	Depositor
Voltage (kV)	200	Depositor
Electron dose ($e^-/\text{Å}^2$)	10	Depositor
Minimum defocus (nm)	500	Depositor
Maximum defocus (nm)	2500	Depositor
Magnification	49000	Depositor
Image detector	KODAK SO-163 FILM	Depositor
Maximum map value	18.352	Depositor
Minimum map value	-18.065	Depositor
Average map value	0.361	Depositor
Map value standard deviation	3.624	Depositor
Recommended contour level	0.6	Depositor
Map size (Å)	510.0, 510.0, 510.0	wwPDB
Map dimensions	200, 200, 200	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	2.55, 2.55, 2.55	Depositor

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.72	2/2604 (0.1%)	1.19	19/3524 (0.5%)
1	D	0.86	8/2683 (0.3%)	1.71	42/3630 (1.2%)
1	G	0.73	2/2604 (0.1%)	1.19	19/3524 (0.5%)
1	K	0.86	8/2683 (0.3%)	1.71	43/3630 (1.2%)
2	B	1.03	4/1718 (0.2%)	2.22	46/2293 (2.0%)
2	C	1.01	3/966 (0.3%)	1.74	34/1298 (2.6%)
2	E	1.18	6/1718 (0.3%)	2.25	67/2293 (2.9%)
2	F	1.01	4/966 (0.4%)	1.73	27/1298 (2.1%)
2	H	1.01	3/966 (0.3%)	1.74	34/1298 (2.6%)
2	I	1.03	4/1718 (0.2%)	2.22	46/2293 (2.0%)
2	J	1.18	6/1718 (0.3%)	2.25	67/2293 (2.9%)
2	L	1.02	3/966 (0.3%)	1.73	26/1298 (2.0%)
All	All	0.95	53/21310 (0.2%)	1.80	470/28672 (1.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	1	12
1	D	1	15
1	G	1	11
1	K	1	15
2	B	7	28
2	C	6	13
2	E	6	37
2	F	4	16
2	H	6	13
2	I	7	28
2	J	6	37
2	L	4	16
All	All	50	241

The worst 5 of 53 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	J	699	SER	CB-OG	-22.93	1.12	1.42
2	E	699	SER	CB-OG	-22.90	1.12	1.42
1	G	245	LYS	CA-CB	-9.85	1.32	1.53
1	A	245	LYS	CA-CB	-9.84	1.32	1.53
1	K	317	PHE	C-O	-9.47	1.05	1.23

The worst 5 of 470 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	492	ASP	CB-CG-OD2	50.51	163.76	118.30
2	I	492	ASP	CB-CG-OD2	50.42	163.68	118.30
2	I	492	ASP	CB-CG-OD1	-45.51	77.34	118.30
2	B	492	ASP	CB-CG-OD1	-45.47	77.37	118.30
1	K	318	ARG	NE-CZ-NH2	39.72	140.16	120.30

5 of 50 chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	A	726	ASP	CA
2	B	366	PHE	CA
2	B	373	GLU	CA
2	B	441	GLN	CA
2	B	442	CYS	CA

5 of 241 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	105	THR	Mainchain
1	A	123	ARG	Sidechain
1	A	28	ASP	Mainchain
1	A	41	SER	Mainchain
1	A	50	ASN	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2567	0	2629	75	0
1	D	2643	0	2691	117	0
1	G	2567	0	2629	76	0
1	K	2643	0	2691	130	0
2	B	1697	0	1748	123	0
2	C	946	0	934	23	0
2	E	1697	0	1749	169	0
2	F	946	0	936	30	0
2	H	946	0	934	24	0
2	I	1697	0	1747	162	0
2	J	1697	0	1749	193	0
2	L	946	0	936	29	0
All	All	20992	0	21373	984	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 23.

The worst 5 of 984 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:E:454:GLU:OE1	1:K:271:ARG:NH1	1.72	1.21
1:D:317:PHE:CZ	2:E:332:MET:O	1.94	1.20
2:J:332:MET:O	1:K:317:PHE:CZ	1.94	1.18
2:J:453:ARG:N	1:K:318:ARG:HD2	1.55	1.17
1:D:318:ARG:HD2	2:E:453:ARG:N	1.55	1.14

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	325/864 (38%)	313 (96%)	10 (3%)	2 (1%)	25 66

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	333/864 (38%)	319 (96%)	13 (4%)	1 (0%)	41	77
1	G	325/864 (38%)	313 (96%)	10 (3%)	2 (1%)	25	66
1	K	333/864 (38%)	319 (96%)	13 (4%)	1 (0%)	41	77
2	B	194/864 (22%)	158 (81%)	28 (14%)	8 (4%)	3	23
2	C	111/864 (13%)	94 (85%)	12 (11%)	5 (4%)	2	22
2	E	194/864 (22%)	166 (86%)	23 (12%)	5 (3%)	5	31
2	F	111/864 (13%)	93 (84%)	16 (14%)	2 (2%)	8	40
2	H	111/864 (13%)	94 (85%)	12 (11%)	5 (4%)	2	22
2	I	194/864 (22%)	158 (81%)	29 (15%)	7 (4%)	3	25
2	J	194/864 (22%)	166 (86%)	23 (12%)	5 (3%)	5	31
2	L	111/864 (13%)	93 (84%)	16 (14%)	2 (2%)	8	40
All	All	2536/10368 (24%)	2286 (90%)	205 (8%)	45 (2%)	12	40

5 of 45 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	88	GLY
2	B	442	CYS
2	B	490	HIS
2	B	690	ILE
2	C	523	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	287/761 (38%)	275 (96%)	12 (4%)	30	54
1	D	295/761 (39%)	281 (95%)	14 (5%)	26	51
1	G	287/761 (38%)	275 (96%)	12 (4%)	30	54
1	K	295/761 (39%)	281 (95%)	14 (5%)	26	51
2	B	191/761 (25%)	158 (83%)	33 (17%)	2	11

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	C	102/761 (13%)	91 (89%)	11 (11%)	6	23
2	E	191/761 (25%)	151 (79%)	40 (21%)	1	6
2	F	102/761 (13%)	89 (87%)	13 (13%)	4	18
2	H	102/761 (13%)	91 (89%)	11 (11%)	6	23
2	I	191/761 (25%)	158 (83%)	33 (17%)	2	11
2	J	191/761 (25%)	151 (79%)	40 (21%)	1	6
2	L	102/761 (13%)	89 (87%)	13 (13%)	4	18
All	All	2336/9132 (26%)	2090 (90%)	246 (10%)	10	24

5 of 246 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	F	569	ASN
1	K	87	LYS
2	H	594	ARG
1	K	34	ILE
2	L	560	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 65 such sidechains are listed below:

Mol	Chain	Res	Type
1	K	148	GLN
1	K	198	GLN
2	E	448	GLN
2	E	335	GLN
1	K	236	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	D	2
1	K	2

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	D	315:LYS	C	316:ASN	N	1.19
1	K	315:LYS	C	316:ASN	N	1.19
1	D	316:ASN	C	317:PHE	N	1.17
1	K	316:ASN	C	317:PHE	N	1.17


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-2701. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map




The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [\(i\)](#)

6.3.1 Primary map



X Index: 145



Y Index: 55




Z Index: 197


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [\(i\)](#)


6.4.1 Primary map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.6. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [\(i\)](#)



The volume at the recommended contour level is 33840 nm^3 ; this corresponds to an approximate mass of 30569 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [i](#)



*Reported resolution corresponds to spatial frequency of 0.080\AA^{-1}


8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-2701 and PDB model 4UUD. Per-residue inclusion information can be found in section [3](#) on page [6](#).

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.6 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [\(i\)](#)




The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [\(i\)](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.6).



























9.4 Atom inclusion [i](#)



At the recommended contour level, 99% of all backbone atoms, 98% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.6) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.9820	 0.0550
A	 0.9940	 0.0640
B	 0.9930	 0.0490
C	 0.9840	 0.0640
D	 0.9550	 0.0480
E	 0.9970	 0.0530
F	 0.9740	 0.0520
G	 0.9940	 0.0660
H	 0.9830	 0.0660
I	 0.9930	 0.0480
J	 0.9970	 0.0540
K	 0.9550	 0.0480
L	 0.9740	 0.0530

