

wwPDB EM Validation Summary Report (i)

Nov 23, 2022 – 05:07 PM JST

PDB ID : 7V9X

EMDB ID : EMD-31829

Title: Cryo-EM structure of E.coli retron-Ec86 in complex with its effector at 2.8

angstrom

Authors: Wang, Y.J.; Guan, Z.Y.; Zou, T.T.

Deposited on : 2021-08-27

Resolution : 2.82 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev43

MolProbity : 4.02b-467

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

MapQ: 1.9.9

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.31.3

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $ELECTRON\ MICROSCOPY$

The reported resolution of this entry is 2.82 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	${ m EM\ structures} \ (\#{ m Entries})$
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826
RNA backbone	4643	859

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion <40%). The numeric value is given above the bar.

Mol	Chain	Length			Quality of c	hain		
1	A	330	<u> </u>		84%		11%	5%
1	В	330	<u> </u>		81%		15%	5%
2	С	307	36%		7%	57%		
3	D	105	23%	14%		63%		
3	G	105	22%	15%		63%		
4	Е	81	28%		35%	9%	28%	
4	Н	81	19%		35%	·	28%	

Continued on next page...



 $Continued\ from\ previous\ page...$

Mol	Chain	Length		Quality of chain						
			14%							
5	F	14	21%	29%	7%	43%				
			21%							
5	I	14		50%	7%	43%				



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 10550 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called RNA-directed DNA polymerase from retron EC86.

\mathbf{Mol}	Chain	Residues		\mathbf{At}	oms		AltConf	Trace	
1	A	314		C 1613		_	 0	0	
1	В	314	Total 2518	C 1613		O 446	0	0	

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	321	LEU	-	expression tag	UNP P23070
A	322	GLU	-	expression tag	UNP P23070
A	323	HIS	=	expression tag	UNP P23070
A	324	HIS	-	expression tag	UNP P23070
A	325	HIS	-	expression tag	UNP P23070
A	326	HIS	-	expression tag	UNP P23070
A	327	HIS	-	expression tag	UNP P23070
A	328	HIS	-	expression tag	UNP P23070
A	329	HIS	-	expression tag	UNP P23070
A	330	HIS	-	expression tag	UNP P23070
В	321	LEU	-	expression tag	UNP P23070
В	322	GLU	-	expression tag	UNP P23070
В	323	HIS	-	expression tag	UNP P23070
В	324	HIS	-	expression tag	UNP P23070
В	325	HIS	ı	expression tag	UNP P23070
В	326	HIS	-	expression tag	UNP P23070
В	327	HIS	-	expression tag	UNP P23070
В	328	HIS	=	expression tag	UNP P23070
В	329	HIS		expression tag	UNP P23070
В	330	HIS	-	expression tag	UNP P23070

• Molecule 2 is a protein called retron St85 family effector protein.



Mol	Chain	Residues		At	oms			AltConf	Trace
2	С	132	Total	С	N	О	S	0	0
		132	1096	704	195	192	5	0	

• Molecule 3 is a DNA chain called DNA (105-MER).

Mol	Chain	Residues	Atoms				AltConf	Trace	
3	D	39	Total 808	_	N 163	_	P 39	0	0
3	G	39	Total 808	_	N 163	_	P 39	0	0

 $\bullet\,$ Molecule 4 is a RNA chain called RNA (81-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace	
1	E	58	Total	С	N	О	Р	0	0	
4	ш	30	1225	547	205	415	58	0		
1	П	58	Total	С	N	О	Р	0	0	
4	11	00	1225	547	205	415	58			

• Molecule 5 is a RNA chain called RNA (14-MER).


Mol	Chain	Residues	${f Atoms}$				AltConf	Trace		
5	Ŀ	Q	Total	С	N	О	Р	0	0	
9	Г	8	176	78	35	55	8	0		
5	Т	Q	Total	С	N	О	Р	0	0	
9	1	0	176	78	35	55	8	U	U	



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: RNA-directed DNA polymerase from retron EC86







• Molecule 3: DNA (105-MER)








• Molecule 3: DNA (105-MER)







• Molecule 4: RNA (81-MER)







• Molecule 4: RNA (81-MER)








• Molecule 5: RNA (14-MER)







4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	137409	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{Å}^2)$	53.68	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	2.761	Depositor
Minimum map value	-1.295	Depositor
Average map value	0.005	Depositor
Map value standard deviation	0.071	Depositor
Recommended contour level	0.4	Depositor
Map size (Å)	238.0, 238.0, 238.0	wwPDB
Map dimensions	280, 280, 280	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.85, 0.85, 0.85	Depositor



5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Во	ond angles
WIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.31	0/2568	0.52	0/3456
1	В	0.31	0/2568	0.52	0/3456
2	С	0.27	0/1112	0.52	0/1489
3	D	0.63	0/910	0.87	0/1400
3	G	0.64	0/910	0.87	0/1400
4	Е	0.35	0/1364	0.87	3/2119 (0.1%)
4	Н	0.32	0/1364	0.85	1/2119 (0.0%)
5	F	0.24	0/197	0.75	0/306
5	I	0.22	0/197	0.72	0/306
All	All	0.38	0/11190	0.70	4/16051 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
4	Е	65	С	C2-N1-C1'	5.78	125.16	118.80
4	Е	65	С	N1-C2-O2	5.76	122.36	118.90
4	Е	65	С	N3-C2-O2	-5.32	118.18	121.90
4	Н	65	С	C2-N1-C1'	5.04	124.35	118.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2518	0	2621	27	0
1	В	2518	0	2621	32	0
2	С	1096	0	1139	12	0
3	D	808	0	437	11	0
3	G	808	0	437	12	0
4	Е	1225	0	621	16	0
4	Н	1225	0	621	15	0
5	F	176	0	88	4	0
5	I	176	0	88	1	0
All	All	10550	0	8673	115	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 115 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$\begin{array}{c} \text{Clash} \\ \text{overlap } (\text{\AA}) \end{array}$
1:B:312:LYS:HG3	1:B:314:PRO:HD3	1.66	0.78
1:B:8:ASN:OD1	1:B:11:ARG:NH2	2.24	0.70
2:C:211:TYR:O	1:B:31:ARG:NH2	2.25	0.69
1:A:312:LYS:HG3	1:A:314:PRO:HD3	1.77	0.67
1:B:60:LYS:NZ	1:B:121:GLU:OE2	2.27	0.67

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	A	312/330 (94%)	295 (95%)	17 (5%)	0	100	100
1	В	312/330 (94%)	300 (96%)	12 (4%)	0	100	100
2	С	126/307 (41%)	125 (99%)	1 (1%)	0	100	100

Continued on next page...



Continued from previous page...

Mo	l Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
All	All	750/967 (78%)	720 (96%)	30 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percen	ntiles
1	A	281/296 (95%)	280 (100%)	1 (0%)	91	97
1	В	281/296 (95%)	281 (100%)	0	100	100
2	С	120/278 (43%)	120 (100%)	0	100	100
All	All	682/870 (78%)	681 (100%)	1 (0%)	93	98

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	11	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
4	Е	57/81 (70%)	21 (36%)	2 (3%)
4	Н	56/81~(69%)	17 (30%)	1 (1%)
5	F	7/14~(50%)	1 (14%)	0
5	I	7/14 (50%)	0	0
All	All	127/190~(66%)	39 (30%)	3 (2%)

5 of 39 RNA backbone outliers are listed below:



Mol	Chain	Res	Type
4	Е	9	С
4	Е	15	С
4	Е	16	G
4	Е	20	G
4	Е	21	G

All (3) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
4	Е	8	С
4	Е	66	A
4	Н	66	A

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.




6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-31829. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections (i)


6.1.1 Primary map




The images above show the map projected in three orthogonal directions.

6.2 Central slices (i)


6.2.1 Primary map







Y Index: 140




Z Index: 140



The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices (i)


6.3.1 Primary map



The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal surface views (i)

6.4.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.4. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.



6.5 Mask visualisation (i)


This section was not generated. No masks/segmentation were deposited.



7 Map analysis (i)

This section contains the results of statistical analysis of the map.


7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



7.2 Volume estimate (i)




The volume at the recommended contour level is 54 nm^3 ; this corresponds to an approximate mass of 49 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



7.3 Rotationally averaged power spectrum (i)



*Reported resolution corresponds to spatial frequency of 0.355 $\rm \mathring{A}^{-1}$



8 Fourier-Shell correlation (i)


This section was not generated. No FSC curve or half-maps provided.



9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-31829 and PDB model 7V9X. Per-residue inclusion information can be found in section 3 on page 6.


9.1 Map-model overlay (i)



The images above show the 3D surface view of the map at the recommended contour level 0.4 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.




9.2 Q-score mapped to coordinate model (i)



The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


9.3 Atom inclusion mapped to coordinate model (i)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.4).



9.4 Atom inclusion (i)



At the recommended contour level, 82% of all backbone atoms, 78% of all non-hydrogen atoms, are inside the map.



9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (0.4) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.7815	0.5490
A	0.8723	0.5880
В	0.8714	0.5900
С	0.6288	0.5470
D	0.8032	0.5530
E	0.7469	0.5010
F	0.6193	0.4390
G	0.7970	0.5610
Н	0.6400	0.4640
I	0.3977	0.3900



