



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2024 – 02:33 AM EST

PDB ID : 8V9Q  
Title : Crystal structure of mGalNAc-T1 in complex with the mucin glycopeptide Muc5AC-13, Mn<sup>2+</sup>, and UDP.  
Authors : Samara, N.L.; Collette, A.M.  
Deposited on : 2023-12-08  
Resolution : 2.29 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.29 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	559	
1	B	559	
2	D	16	
2	F	16	
2	H	16	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	X	5	
4	C	2	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	MAN	X	4	-	-	-	X
4	NAG	C	2	-	-	-	X

## 2 Entry composition [i](#)

There are 12 unique types of molecules in this entry. The entry contains 16274 atoms, of which 7964 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Polypeptide N-acetylgalactosaminyltransferase 1 soluble form.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	500	7974	2543	3946	717	739	29	0	0	0
1	B	449	7227	2302	3579	656	664	26	0	0	0

- Molecule 2 is a protein called Mucin-5AC.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	H	N	O			
2	D	16	207	63	103	16	25	0	0	0
2	F	15	193	59	96	15	23	0	0	0
2	H	7	73	26	28	7	12	0	0	0

- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	H	N	O			
3	X	5	115	34	54	2	25	0	0	0

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	H	N	O			
4	C	2	53	16	25	2	10	0	0	0

- Molecule 5 is URIDINE-5'-DIPHOSPHATE (three-letter code: UDP) (formula: C<sub>9</sub>H<sub>14</sub>N<sub>2</sub>O<sub>12</sub>P<sub>2</sub>) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	H	N	O			P
5	A	1	36	9	11	2	12	2	0	0
5	B	1	36	9	11	2	12	2	0	0

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	H	O	0	0
			13	3	7	3		
6	A	1	Total	C	H	O	0	0
			14	3	8	3		

- Molecule 7 is SUCCINIC ACID (three-letter code: SIN) (formula: C<sub>4</sub>H<sub>6</sub>O<sub>4</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	A	1	Total	C	H	O	0	0
			12	4	4	4		

- Molecule 8 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	1	Total C H O 10 2 6 2	0	0
8	A	1	Total C H O 10 2 6 2	0	0
8	A	1	Total C H O 9 2 5 2	0	0
8	D	1	Total C H O 10 2 6 2	0	0
8	D	1	Total C O 4 2 2	0	0

- Molecule 9 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn) (labeled as "Ligand of Interest" by depositor).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	1	Total Mn 1 1	0	0
9	B	1	Total Mn 1 1	0	0

- Molecule 10 is 2-acetamido-2-deoxy-alpha-D-galactopyranose (three-letter code: A2G) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf		
			Total	C	H	N			O	
10	D	1	Total	28	8	14	1	5	0	0
10	F	1	Total	28	8	14	1	5	0	0
10	H	1	Total	27	8	13	1	5	0	0

- Molecule 11 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).






Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
11	B	1	Total	C	H	N	O	0	0
			28	8	14	1	5		
11	B	1	Total	C	H	N	O	0	0
			28	8	14	1	5		


- Molecule 12 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
12	A	132	Total	O	0	0
			132	132		
12	D	4	Total	O	0	0
			4	4		
12	B	1	Total	O	0	0
			1	1		








- Molecule 2: Mucin-5AC



- Molecule 2: Mucin-5AC



- Molecule 2: Mucin-5AC



- Molecule 3: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C:  100%

3AG2  
3AG2

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	59.63Å 72.76Å 148.50Å 90.00° 95.44° 90.00°	Depositor
Resolution (Å)	29.90 – 2.29 29.90 – 2.29	Depositor EDS
% Data completeness (in resolution range)	96.6 (29.90-2.29) 96.6 (29.90-2.29)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.30 (at 2.29Å)	Xtrriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, $R_{free}$	0.248 , 0.278 0.245 , 0.276	Depositor DCC
$R_{free}$ test set	2757 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	40.2	Xtrriage
Anisotropy	0.096	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 55.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	16274	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	89.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.65% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: A2G, BMA, MAN, GOL, SIN, EDO, MN, NAG, UDP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.26	0/4119	0.56	0/5570
1	B	0.25	0/3705	0.54	0/4970
2	D	0.34	0/107	0.53	0/151
2	F	0.33	0/99	0.51	0/138
2	H	0.24	0/45	0.43	0/62
All	All	0.26	0/8075	0.55	0/10891

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
All	All	0	2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	551	ARG	Sidechain
1	B	551	ARG	Sidechain

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4028	3946	3983	41	1
1	B	3648	3579	3604	75	0
2	D	104	103	103	0	0
2	F	97	96	95	3	0
2	H	45	28	41	6	0
3	X	61	54	52	0	0
4	C	28	25	25	0	0
5	A	25	11	10	2	0
5	B	25	11	10	3	0
6	A	12	15	16	0	0
7	A	8	4	4	0	0
8	A	12	17	18	0	0
8	D	8	6	12	0	0
9	A	1	0	0	0	0
9	B	1	0	0	0	0
10	D	14	14	12	0	0
10	F	14	14	12	2	0
10	H	14	13	12	3	0
11	B	28	28	26	0	0
12	A	132	0	0	3	0
12	B	1	0	0	0	0
12	D	4	0	0	0	0
All	All	8310	7964	8035	124	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 124 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:158:SER:HB3	1:B:163:LEU:HD13	1.52	0.89
1:B:258:GLY:N	1:B:290:THR:HG1	1.69	0.88
1:A:509:ASP:OD2	1:A:512:LYS:HE2	1.87	0.74
1:B:298:ILE:HD11	1:B:302:TYR:HD2	1.54	0.73
1:B:404:ARG:HE	1:B:404:ARG:HA	1.60	0.67

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:80:LYS:HZ3	1:A:198:ALA:O[2_656]	1.55	0.05

## 5.3 Torsion angles [\(i\)](#)

### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	498/559 (89%)	479 (96%)	19 (4%)	0	100	100
1	B	399/559 (71%)	380 (95%)	19 (5%)	0	100	100
2	D	14/16 (88%)	14 (100%)	0	0	100	100
2	F	11/16 (69%)	11 (100%)	0	0	100	100
2	H	5/16 (31%)	3 (60%)	2 (40%)	0	100	100
All	All	927/1166 (80%)	887 (96%)	40 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	447/501 (89%)	433 (97%)	14 (3%)	40	55
1	B	409/501 (82%)	393 (96%)	16 (4%)	32	46
2	D	14/14 (100%)	14 (100%)	0	100	100

*Continued on next page...*



Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	F	13/14 (93%)	13 (100%)	0	100	100
2	H	6/14 (43%)	4 (67%)	2 (33%)	0	0
All	All	889/1044 (85%)	857 (96%)	32 (4%)	35	49

5 of 32 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	488	LEU
1	B	513	LEU
1	A	537	ILE
1	A	534	VAL
2	H	10	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	72	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

7 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	NAG	C	1	1,4	14,14,15	0.28	0	17,19,21	0.49	0
4	NAG	C	2	4	14,14,15	0.29	0	17,19,21	0.62	0
3	NAG	X	1	3,1	14,14,15	0.18	0	17,19,21	0.40	0
3	NAG	X	2	3	14,14,15	0.20	0	17,19,21	0.68	1 (5%)
3	BMA	X	3	3	11,11,12	0.72	0	15,15,17	0.87	0
3	MAN	X	4	3	11,11,12	0.56	0	15,15,17	1.03	2 (13%)
3	MAN	X	5	3	11,11,12	0.68	0	15,15,17	1.04	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	C	1	1,4	-	2/6/23/26	0/1/1/1
4	NAG	C	2	4	-	0/6/23/26	0/1/1/1
3	NAG	X	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	X	2	3	-	2/6/23/26	0/1/1/1
3	BMA	X	3	3	-	2/2/19/22	0/1/1/1
3	MAN	X	4	3	-	0/2/19/22	0/1/1/1
3	MAN	X	5	3	-	1/2/19/22	0/1/1/1

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	X	5	MAN	C1-O5-C5	2.85	116.05	112.19
3	X	4	MAN	C1-O5-C5	2.72	115.88	112.19
3	X	5	MAN	O2-C2-C3	-2.20	105.72	110.14
3	X	2	NAG	C1-O5-C5	2.13	115.08	112.19
3	X	4	MAN	O2-C2-C3	-2.01	106.10	110.14

There are no chirality outliers.

5 of 7 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	X	2	NAG	C8-C7-N2-C2
3	X	2	NAG	O7-C7-N2-C2
4	C	1	NAG	C8-C7-N2-C2
4	C	1	NAG	O7-C7-N2-C2

*Continued on next page...*


Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	X	5	MAN	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





## 5.6 Ligand geometry [i](#)

Of 17 ligands modelled in this entry, 2 are monoatomic - leaving 15 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
8	EDO	A	605	-	3,3,3	0.51	0	2,2,2	0.20	0
8	EDO	D	103	-	3,3,3	0.48	0	2,2,2	0.26	0
10	A2G	H	101	2	14,14,15	1.78	4 (28%)	17,19,21	1.21	1 (5%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
10	A2G	F	101	2	14,14,15	1.83	5 (35%)	17,19,21	2.23	7 (41%)
11	NAG	B	603	-	14,14,15	0.21	0	17,19,21	0.50	0
8	EDO	A	606	-	3,3,3	0.48	0	2,2,2	0.31	0
5	UDP	A	601	9	24,26,26	3.70	13 (54%)	37,40,40	1.47	6 (16%)
5	UDP	B	601	9	24,26,26	3.74	13 (54%)	37,40,40	1.63	6 (16%)
6	GOL	A	608	-	5,5,5	0.91	0	5,5,5	0.89	0
6	GOL	A	602	-	5,5,5	0.87	0	5,5,5	0.96	0
8	EDO	A	604	-	3,3,3	0.45	0	2,2,2	0.38	0
10	A2G	D	101	2	14,14,15	1.74	3 (21%)	17,19,21	0.89	0
11	NAG	B	602	1	14,14,15	0.25	0	17,19,21	0.52	0
7	SIN	A	603	-	7,7,7	1.15	0	8,8,8	1.56	1 (12%)
8	EDO	D	102	-	3,3,3	0.47	0	2,2,2	0.30	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	EDO	A	605	-	-	1/1/1/1	-
8	EDO	D	103	-	-	0/1/1/1	-
10	A2G	H	101	2	-	4/6/23/26	0/1/1/1
10	A2G	F	101	2	-	3/6/23/26	0/1/1/1
11	NAG	B	603	-	-	2/6/23/26	0/1/1/1
8	EDO	A	606	-	-	0/1/1/1	-
5	UDP	A	601	9	-	2/16/32/32	0/2/2/2
5	UDP	B	601	9	-	0/16/32/32	0/2/2/2
6	GOL	A	608	-	-	0/4/4/4	-
6	GOL	A	602	-	-	1/4/4/4	-
8	EDO	A	604	-	-	1/1/1/1	-
10	A2G	D	101	2	-	0/6/23/26	0/1/1/1
11	NAG	B	602	1	-	4/6/23/26	0/1/1/1
7	SIN	A	603	-	-	2/5/5/5	-
8	EDO	D	102	-	-	0/1/1/1	-

The worst 5 of 38 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	601	UDP	O4'-C4'	7.31	1.61	1.45
5	A	601	UDP	O4'-C4'	7.28	1.61	1.45

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	601	UDP	C3'-C4'	-7.21	1.34	1.53
5	B	601	UDP	C3'-C4'	-7.08	1.34	1.53
5	B	601	UDP	C2-N1	6.91	1.49	1.38

The worst 5 of 21 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	601	UDP	C4-N3-C2	-5.10	119.86	126.58
5	A	601	UDP	C4-N3-C2	-4.94	120.07	126.58
10	F	101	A2G	O5-C1-C2	-4.53	104.14	111.29
10	F	101	A2G	C8-C7-N2	4.14	123.11	116.10
5	A	601	UDP	N3-C2-N1	3.82	119.95	114.89

There are no chirality outliers.

5 of 20 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
11	B	603	NAG	O5-C5-C6-O6
10	H	101	A2G	C4-C5-C6-O6
11	B	603	NAG	C4-C5-C6-O6
10	F	101	A2G	O7-C7-N2-C2
10	F	101	A2G	C8-C7-N2-C2


There are no ring outliers.



4 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
10	H	101	A2G	3	0
10	F	101	A2G	2	0
5	A	601	UDP	2	0
5	B	601	UDP	3	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.

The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	500/559 (89%)	0.96	60 (12%) 4 6	22, 38, 63, 113	0
1	B	449/559 (80%)	3.56	338 (75%) 0 0	84, 124, 158, 209	0
2	D	16/16 (100%)	0.99	2 (12%) 3 5	30, 39, 64, 72	0
2	F	15/16 (93%)	2.99	12 (80%) 0 0	110, 130, 146, 170	0
2	H	7/16 (43%)	2.37	4 (57%) 0 0	72, 93, 121, 137	0
All	All	987/1166 (84%)	2.18	416 (42%) 0 0	22, 64, 149, 209	0

The worst 5 of 416 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	66	VAL	10.6
1	B	457	PHE	9.8
1	B	296	PHE	9.7
1	B	540	CYS	9.5
1	B	292	ALA	9.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	MAN	X	4	11/12	0.42	0.48	101,119,142,151	0


*Continued on next page...*

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	MAN	X	5	11/12	0.45	0.40	86,103,118,122	0
3	BMA	X	3	11/12	0.61	0.39	93,106,127,131	0
4	NAG	C	2	14/15	0.67	0.48	75,90,108,111	0
3	NAG	X	2	14/15	0.75	0.25	58,77,98,105	0
4	NAG	C	1	14/15	0.84	0.18	51,60,72,80	0
3	NAG	X	1	14/15	0.92	0.11	48,62,85,85	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
9	MN	B	604	1/1	0.61	0.12	143,143,143,143	0
10	A2G	F	101	14/15	0.62	0.32	116,134,152,158	0
6	GOL	A	602	6/6	0.63	0.24	54,77,96,96	0
11	NAG	B	603	14/15	0.64	0.26	84,110,138,138	0
5	UDP	B	601	25/25	0.71	0.23	118,132,152,158	0
6	GOL	A	608	6/6	0.74	0.30	34,56,87,104	0
8	EDO	D	102	4/4	0.77	0.20	41,50,77,77	0
7	SIN	A	603	8/8	0.81	0.23	59,78,93,93	0

*Continued on next page...*




*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
10	A2G	H	101	14/15	0.82	0.21	42,57,72,82	0
11	NAG	B	602	14/15	0.83	0.22	93,113,126,139	0
8	EDO	A	605	4/4	0.83	0.25	31,37,45,45	0
8	EDO	D	103	4/4	0.88	0.21	24,34,35,37	0
8	EDO	A	606	4/4	0.88	0.24	37,45,55,66	0
8	EDO	A	604	4/4	0.88	0.16	39,46,53,60	0
5	UDP	A	601	25/25	0.92	0.25	18,42,56,63	0
10	A2G	D	101	14/15	0.96	0.20	19,26,33,34	0
9	MN	A	607	1/1	0.99	0.13	25,25,25,25	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around MN B 604:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around A2G F 101:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)













**Electron density around A2G H 101:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







**Electron density around UDP A 601:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around A2G D 101:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers ⓘ

There are no such residues in this entry.