



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 18, 2024 – 10:07 AM EDT

PDB ID : 3VEP
Title : Crystal structure of SigD4 in complex with its negative regulator RsdA
Authors : Jaiswal, R.K.; Gopal, B.
Deposited on : 2012-01-09
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 2.37.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	C	80	
1	G	80	
1	J	80	
1	X	80	
2	A	86	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	D	86	
2	E	86	
2	H	86	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	SO4	E	302	-	-	X	-
3	SO4	E	303	-	-	X	-

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3680 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Uncharacterized protein Rv3413c/MT3522.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
1	X	46	372	234	64	74	0	0	0
1	C	47	379	239	65	75	0	0	0
1	G	47	377	237	65	75	0	0	0
1	J	47	376	236	65	75	0	0	0

- Molecule 2 is a protein called Probable RNA polymerase sigma-D factor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	Se			
2	D	68	505	313	93	97	2	0	0	0
2	A	70	511	316	92	101	2	0	0	0
2	E	70	517	319	95	101	2	0	0	0
2	H	71	526	325	97	102	2	0	0	0

There are 56 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	127	MSE	-	EXPRESSION TAG	UNP P66811
D	128	GLY	-	EXPRESSION TAG	UNP P66811
D	129	SER	-	EXPRESSION TAG	UNP P66811
D	130	SER	-	EXPRESSION TAG	UNP P66811
D	131	HIS	-	EXPRESSION TAG	UNP P66811
D	132	HIS	-	EXPRESSION TAG	UNP P66811
D	133	HIS	-	EXPRESSION TAG	UNP P66811
D	134	HIS	-	EXPRESSION TAG	UNP P66811

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
D	135	HIS	-	EXPRESSION TAG	UNP P66811
D	136	HIS	-	EXPRESSION TAG	UNP P66811
D	137	SER	-	EXPRESSION TAG	UNP P66811
D	138	GLN	-	EXPRESSION TAG	UNP P66811
D	139	ASP	-	EXPRESSION TAG	UNP P66811
D	140	PRO	-	EXPRESSION TAG	UNP P66811
A	127	MSE	-	EXPRESSION TAG	UNP P66811
A	128	GLY	-	EXPRESSION TAG	UNP P66811
A	129	SER	-	EXPRESSION TAG	UNP P66811
A	130	SER	-	EXPRESSION TAG	UNP P66811
A	131	HIS	-	EXPRESSION TAG	UNP P66811
A	132	HIS	-	EXPRESSION TAG	UNP P66811
A	133	HIS	-	EXPRESSION TAG	UNP P66811
A	134	HIS	-	EXPRESSION TAG	UNP P66811
A	135	HIS	-	EXPRESSION TAG	UNP P66811
A	136	HIS	-	EXPRESSION TAG	UNP P66811
A	137	SER	-	EXPRESSION TAG	UNP P66811
A	138	GLN	-	EXPRESSION TAG	UNP P66811
A	139	ASP	-	EXPRESSION TAG	UNP P66811
A	140	PRO	-	EXPRESSION TAG	UNP P66811
E	127	MSE	-	EXPRESSION TAG	UNP P66811
E	128	GLY	-	EXPRESSION TAG	UNP P66811
E	129	SER	-	EXPRESSION TAG	UNP P66811
E	130	SER	-	EXPRESSION TAG	UNP P66811
E	131	HIS	-	EXPRESSION TAG	UNP P66811
E	132	HIS	-	EXPRESSION TAG	UNP P66811
E	133	HIS	-	EXPRESSION TAG	UNP P66811
E	134	HIS	-	EXPRESSION TAG	UNP P66811
E	135	HIS	-	EXPRESSION TAG	UNP P66811
E	136	HIS	-	EXPRESSION TAG	UNP P66811
E	137	SER	-	EXPRESSION TAG	UNP P66811
E	138	GLN	-	EXPRESSION TAG	UNP P66811
E	139	ASP	-	EXPRESSION TAG	UNP P66811
E	140	PRO	-	EXPRESSION TAG	UNP P66811
H	127	MSE	-	EXPRESSION TAG	UNP P66811
H	128	GLY	-	EXPRESSION TAG	UNP P66811
H	129	SER	-	EXPRESSION TAG	UNP P66811
H	130	SER	-	EXPRESSION TAG	UNP P66811
H	131	HIS	-	EXPRESSION TAG	UNP P66811
H	132	HIS	-	EXPRESSION TAG	UNP P66811
H	133	HIS	-	EXPRESSION TAG	UNP P66811
H	134	HIS	-	EXPRESSION TAG	UNP P66811

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
H	135	HIS	-	EXPRESSION TAG	UNP P66811
H	136	HIS	-	EXPRESSION TAG	UNP P66811
H	137	SER	-	EXPRESSION TAG	UNP P66811
H	138	GLN	-	EXPRESSION TAG	UNP P66811
H	139	ASP	-	EXPRESSION TAG	UNP P66811
H	140	PRO	-	EXPRESSION TAG	UNP P66811

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	X	1	Total O S 5 4 1	0	0
3	D	1	Total O S 5 4 1	0	0
3	D	1	Total O S 5 4 1	0	0
3	C	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	E	1	Total O S 5 4 1	0	0
3	E	1	Total O S 5 4 1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	E	1	Total 5	O 4	S 1	0	0
3	H	1	Total 5	O 4	S 1	0	0
3	H	1	Total 5	O 4	S 1	0	0
3	H	1	Total 5	O 4	S 1	0	0
3	H	1	Total 5	O 4	S 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	X	5	Total 5	O 5	0	0
4	D	6	Total 6	O 6	0	0
4	C	9	Total 9	O 9	0	0
4	A	10	Total 10	O 10	0	0
4	G	6	Total 6	O 6	0	0
4	E	3	Total 3	O 3	0	0
4	J	6	Total 6	O 6	0	0
4	H	7	Total 7	O 7	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Uncharacterized protein Rv3413c/MT3522





- Molecule 1: Uncharacterized protein Rv3413c/MT3522



- Molecule 1: Uncharacterized protein Rv3413c/MT3522




- Molecule 1: Uncharacterized protein Rv3413c/MT3522




- Molecule 2: Probable RNA polymerase sigma-D factor








● Molecule 2: Probable RNA polymerase sigma-D factor



● Molecule 2: Probable RNA polymerase sigma-D factor



● Molecule 2: Probable RNA polymerase sigma-D factor



ALA

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	99.74Å 110.72Å 73.13Å 90.00° 133.00° 90.00°	Depositor
Resolution (Å)	38.47 – 2.50 38.46 – 2.50	Depositor EDS
% Data completeness (in resolution range)	86.6 (38.47-2.50) 86.4 (38.46-2.50)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.97 (at 2.51Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.6.4_486)	Depositor
R, R_{free}	0.243 , 0.287 0.253 , 0.270	Depositor DCC
R_{free} test set	935 reflections (5.16%)	wwPDB-VP
Wilson B-factor (Å ²)	46.7	Xtrriage
Anisotropy	0.450	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 30.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.022 for -h,-h-2*1,1/2*h-1/2*k 0.017 for -h,h+2*1,1/2*h+1/2*k 0.428 for h,-k,-h-l	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	3680	wwPDB-VP
Average B, all atoms (Å ²)	53.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.82% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	C	0.50	0/388	0.86	0/531
1	G	0.55	0/385	0.87	0/527
1	J	0.47	0/385	0.76	1/527 (0.2%)
1	X	0.56	0/380	0.77	0/519
2	A	0.56	0/511	0.72	1/691 (0.1%)
2	D	0.61	0/505	0.84	3/682 (0.4%)
2	E	0.57	1/517 (0.2%)	0.71	1/698 (0.1%)
2	H	0.57	0/526	0.79	1/709 (0.1%)
All	All	0.55	1/3597 (0.0%)	0.79	7/4884 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	J	0	1
2	A	0	2
2	D	0	1
All	All	0	4

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	141	MSE	CG-SE	-5.02	1.78	1.95

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	J	12	PRO	C-N-CD	-8.30	102.34	120.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	150	ARG	NE-CZ-NH1	-5.70	117.45	120.30
2	D	183	VAL	CG1-CB-CG2	5.55	119.78	110.90
2	H	163	ARG	NE-CZ-NH2	-5.53	117.53	120.30
2	D	150	ARG	CB-CG-CD	-5.13	98.27	111.60

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	A	208	ALA	Peptide
2	A	209	GLY	Peptide
2	D	207	ALA	Peptide
1	J	14	LEU	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	379	0	361	27	0
1	G	377	0	355	23	0
1	J	376	0	352	30	0
1	X	372	0	354	15	0
2	A	511	0	527	35	0
2	D	505	0	531	37	0
2	E	517	0	538	37	0
2	H	526	0	551	38	0
3	A	10	0	0	0	0
3	C	5	0	0	1	0
3	D	10	0	0	1	0
3	E	15	0	0	5	0
3	H	20	0	0	0	0
3	X	5	0	0	0	0
4	A	10	0	0	0	0
4	C	9	0	0	1	0
4	D	6	0	0	1	0
4	E	3	0	0	0	0
4	G	6	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	H	7	0	0	1	0
4	J	6	0	0	0	0
4	X	5	0	0	0	0
All	All	3680	0	3569	197	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 27.

The worst 5 of 197 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:205:ILE:HD13	2:H:205:ILE:O	1.37	1.25
1:J:12:PRO:N	1:J:13:PRO:CD	1.99	1.24
1:J:14:LEU:HD23	1:J:14:LEU:C	1.54	1.24
1:J:12:PRO:N	1:J:13:PRO:HD3	1.51	1.15
2:A:141:MSE:N	2:A:141:MSE:SE	2.30	1.15

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	45/80 (56%)	43 (96%)	1 (2%)	1 (2%)	6	10
1	G	45/80 (56%)	44 (98%)	1 (2%)	0	100	100
1	J	45/80 (56%)	42 (93%)	2 (4%)	1 (2%)	6	10
1	X	44/80 (55%)	42 (96%)	2 (4%)	0	100	100
2	A	68/86 (79%)	67 (98%)	1 (2%)	0	100	100
2	D	66/86 (77%)	65 (98%)	1 (2%)	0	100	100
2	E	68/86 (79%)	67 (98%)	1 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	H	69/86 (80%)	68 (99%)	1 (1%)	0	100	100
All	All	450/664 (68%)	438 (97%)	10 (2%)	2 (0%)	34	54

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	13	PRO
1	J	57	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	39/61 (64%)	35 (90%)	4 (10%)	7	14
1	G	38/61 (62%)	32 (84%)	6 (16%)	2	4
1	J	38/61 (62%)	34 (90%)	4 (10%)	7	13
1	X	38/61 (62%)	32 (84%)	6 (16%)	2	4
2	A	52/65 (80%)	45 (86%)	7 (14%)	4	7
2	D	52/65 (80%)	45 (86%)	7 (14%)	4	7
2	E	53/65 (82%)	46 (87%)	7 (13%)	4	7
2	H	54/65 (83%)	44 (82%)	10 (18%)	1	2
All	All	364/504 (72%)	313 (86%)	51 (14%)	3	6

5 of 51 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	23	LEU
2	E	201	LEU
2	H	206	VAL
1	G	27	LEU
2	E	174	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	X	49	GLN
1	C	49	GLN
1	G	49	GLN
1	J	49	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

13 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	A	301	-	4,4,4	0.27	0	6,6,6	0.44	0
3	SO4	E	303	-	4,4,4	0.39	0	6,6,6	0.55	0
3	SO4	H	302	-	4,4,4	0.37	0	6,6,6	0.31	0
3	SO4	H	303	-	4,4,4	0.20	0	6,6,6	0.38	0
3	SO4	H	304	-	4,4,4	0.26	0	6,6,6	0.20	0
3	SO4	C	101	-	4,4,4	0.35	0	6,6,6	0.10	0
3	SO4	X	101	-	4,4,4	0.27	0	6,6,6	0.14	0
3	SO4	D	302	-	4,4,4	0.35	0	6,6,6	0.28	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	E	301	-	4,4,4	0.28	0	6,6,6	0.29	0
3	SO4	H	301	-	4,4,4	0.28	0	6,6,6	0.19	0
3	SO4	D	301	-	4,4,4	0.24	0	6,6,6	0.44	0
3	SO4	E	302	-	4,4,4	0.38	0	6,6,6	0.35	0
3	SO4	A	302	-	4,4,4	0.25	0	6,6,6	0.18	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	E	303	SO4	2	0
3	C	101	SO4	1	0
3	D	302	SO4	1	0
3	E	302	SO4	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	C	47/80 (58%)	0.42	2 (4%) 35 38	20, 49, 76, 89	0
1	G	47/80 (58%)	0.69	5 (10%) 6 5	34, 50, 78, 88	0
1	J	47/80 (58%)	0.37	5 (10%) 6 5	20, 49, 75, 87	0
1	X	46/80 (57%)	0.47	1 (2%) 62 65	34, 48, 76, 88	0
2	A	68/86 (79%)	1.10	13 (19%) 1 1	20, 51, 77, 82	0
2	D	66/86 (76%)	0.86	9 (13%) 3 2	28, 51, 73, 83	0
2	E	68/86 (79%)	1.15	16 (23%) 0 0	31, 51, 77, 82	0
2	H	69/86 (80%)	0.79	6 (8%) 10 10	33, 53, 77, 83	0
All	All	458/664 (68%)	0.77	57 (12%) 4 3	20, 51, 77, 89	0

The worst 5 of 57 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	E	208	ALA	6.2
2	A	206	VAL	5.6
2	E	171	VAL	5.1
2	H	180	ALA	4.6
2	A	156	GLU	4.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	SO4	E	302	5/5	0.74	0.20	58,60,69,86	0
3	SO4	H	303	5/5	0.76	0.24	63,65,77,87	0
3	SO4	H	302	5/5	0.85	0.16	72,81,108,108	0
3	SO4	D	301	5/5	0.86	0.21	50,52,83,84	0
3	SO4	A	302	5/5	0.86	0.16	68,72,76,89	0
3	SO4	E	301	5/5	0.86	0.21	60,66,76,86	0
3	SO4	E	303	5/5	0.88	0.13	62,70,86,86	0
3	SO4	C	101	5/5	0.89	0.24	30,30,30,30	0
3	SO4	A	301	5/5	0.91	0.14	53,61,71,77	0
3	SO4	H	301	5/5	0.91	0.09	62,68,79,87	0
3	SO4	X	101	5/5	0.92	0.12	67,80,88,92	0
3	SO4	D	302	5/5	0.94	0.15	65,74,85,90	0
3	SO4	H	304	5/5	0.94	0.14	66,72,87,89	0

6.5 Other polymers [i](#)

There are no such residues in this entry.