



wwPDB NMR Structure Validation Summary Report ⓘ

Jun 3, 2023 – 04:25 AM EDT

PDB ID : 5VKG
BMRB ID : 30285
Title : Solution-state NMR structural ensemble of human Tsg101 UEV in complex with tenatoprazole
Authors : Strickland, M.; Ehrlich, L.S.; Watanabe, S.; Khan, M.; Strub, M.-P.; Luan, C.H.; Powell, M.D.; Leis, J.; Tjandra, N.; Carter, C.
Deposited on : 2017-04-21

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
BMRB Restraints Analysis : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.33


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 86%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	145	 72% 18% 8%

2 Ensemble composition and analysis

This entry contains 20 models. Model 2 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:3-A:42, A:50-A:142 (133)	0.47	2

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 4 single-model clusters were found.

Cluster number	Models
1	2, 4, 5, 10, 14, 16, 17, 20
2	3, 6, 9, 11, 15, 18
3	1, 7
Single-model clusters	8; 12; 13; 19

3 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 2386 atoms, of which 1201 are hydrogens and 0 are deuteriums.


- Molecule 1 is a protein called Tumor susceptibility gene 101 protein.

Mol	Chain	Residues	Atoms					Trace	
			Total	C	H	N	O		S
1	A	144	2345	757	1183	188	211	6	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	MET	engineered mutation	UNP Q99816

- Molecule 2 is 4-methoxy-1-(5-methoxy-3H-imidazo[4,5-b]pyridin-2-yl)-3,5-dimethyl-2-(sulfa nylmethyl)pyridin-1-ium (three-letter code: 4N1) (formula: C₁₆H₁₉N₄O₂S).




Mol	Chain	Residues	Atoms					
			Total	C	H	N	O	S
2	A	1	41	16	18	4	2	1

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


- Molecule 1: Tumor susceptibility gene 101 protein



4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 2. Colouring as in section 4.1 above.

- Molecule 1: Tumor susceptibility gene 101 protein



5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR NIH	refinement	2.37.7
Gaussian	geometry optimization	9
X-PLOR NIH	structure calculation	2.37.7

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	1742
Number of shifts mapped to atoms	1739
Number of unparsed shifts	0
Number of shifts with mapping errors	3
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	86%

6 Model quality [i](#)

6.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:
4N1

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	1082	1112	1109	16±4
All	All	22100	22600	22180	325

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

5 of 86 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:8:LEU:HD23	1:A:22:VAL:HG13	0.84	1.49	13	20
1:A:111:LEU:HD22	1:A:134:VAL:HG11	0.77	1.56	1	6
1:A:8:LEU:O	1:A:12:VAL:HG22	0.77	1.80	10	20
1:A:11:MET:SD	1:A:39:LEU:HD13	0.72	2.24	7	14
1:A:31:LEU:HD23	1:A:125:LEU:HD21	0.70	1.62	18	6

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	133/145 (92%)	125±1 (94±1%)	7±1 (5±1%)	2±1 (1±0%)	17	64
All	All	2660/2900 (92%)	2493 (94%)	136 (5%)	31 (1%)	17	64

All 3 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	120	PRO	20
1	A	123	ASP	9
1	A	139	PRO	2

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	126/135 (93%)	109±3 (87±2%)	17±3 (13±2%)	7	48
All	All	2520/2700 (93%)	2186 (87%)	334 (13%)	7	48

5 of 62 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	29	ILE	20
1	A	83	ASN	20
1	A	74	LEU	19
1	A	63	TYR	18
1	A	96	THR	13

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mol	Type	Chain	Res	Link	Bond lengths		
					Counts	RMSZ	#Z>2
2	4N1	A	201	1	21,25,25	1.36±0.02	4±0 (19±0%)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mol	Type	Chain	Res	Link	Bond angles		
					Counts	RMSZ	#Z>2
2	4N1	A	201	1	23,36,36	2.47±0.06	13±0 (54±2%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means

no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	4N1	A	201	1	-	0±0,4,10,10	0±0,3,3,3

All unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
2	A	201	4N1	C1-N2	2.85	1.34	1.30	11	20
2	A	201	4N1	O14-C14	2.69	1.33	1.38	6	20
2	A	201	4N1	C3-N7	2.58	1.39	1.34	6	20
2	A	201	4N1	C3-N2	2.55	1.31	1.36	14	20

5 of 13 unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.


Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
2	A	201	4N1	C16-N11-C8	4.94	114.62	119.83	3	20
2	A	201	4N1	C12-N11-C8	4.11	124.36	119.83	3	20
2	A	201	4N1	C21-O14-C14	4.00	125.75	114.78	11	20
2	A	201	4N1	O1-C1-C6	3.93	123.55	116.71	10	20
2	A	201	4N1	O14-C14-C13	3.58	113.80	118.83	12	20

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 86% for the well-defined parts and 86% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: *Tsg101_N16_chemical_shifts_IUPAC.str*

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1742
Number of shifts mapped to atoms	1739
Number of unparsed shifts	0
Number of shifts with mapping errors	3
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	8

The following assigned chemical shifts were not mapped to the molecules present in the coordinate file.

- No matching atom found in the structure. All 3 occurrences are reported below.

List ID	Chain	Res	Type	Atom	Shift Data		
					Value	Uncertainty	Ambiguity
1	A	1	GLY	HA2	3.816	.	.
1	A	1	GLY	HA3	3.816	.	.
1	A	1	GLY	CA	43.219	.	.

7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	143	0.02 ± 0.15	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	135	0.22 ± 0.11	None needed (< 0.5 ppm)
$^{13}\text{C}'$	0	—	None (insufficient data)
^{15}N	131	-0.00 ± 0.35	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 86%, i.e. 1635 atoms were assigned a chemical shift out of a possible 1900. 0 out of 28 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	511/647 (79%)	258/260 (99%)	132/266 (50%)	121/121 (100%)
Sidechain	996/1085 (92%)	685/712 (96%)	302/341 (89%)	9/32 (28%)
Aromatic	128/168 (76%)	70/79 (89%)	56/81 (69%)	2/8 (25%)
Overall	1635/1900 (86%)	1013/1051 (96%)	490/688 (71%)	132/161 (82%)

7.1.4 Statistically unusual chemical shifts [i](#)


The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	25	THR	HG1	3.92	0.08 – 2.19	13.2
1	A	21	THR	HG1	3.85	0.08 – 2.19	12.9
1	A	122	SER	HB2	1.20	2.61 – 5.13	-10.6
1	A	81	PRO	HG2	-0.33	0.41 – 3.45	-7.4
1	A	122	SER	HB3	1.91	2.49 – 5.20	-7.2
1	A	120	PRO	HA	2.52	2.78 – 6.00	-5.8
1	A	81	PRO	HA	2.54	2.78 – 6.00	-5.7
1	A	78	ASP	HA	2.98	3.04 – 6.12	-5.2

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:



8 NMR restraints analysis

8.1 Conformationally restricting restraints

The following table provides the summary of experimentally observed NMR restraints in different categories. Restraints are classified into different categories based on the sequence separation of the atoms involved.

Description	Value
Total distance restraints	4737
Intra-residue ($ i-j =0$)	2206
Sequential ($ i-j =1$)	960
Medium range ($ i-j >1$ and $ i-j <5$)	627
Long range ($ i-j \geq 5$)	858
Inter-chain	0
Hydrogen bond restraints	86
Disulfide bond restraints	0
Total dihedral-angle restraints	276
Number of unmapped restraints	1
Number of restraints per residue	34.6
Number of long range restraints per residue ¹	6.2

¹Long range hydrogen bonds and disulfide bonds are counted as long range restraints while calculating the number of long range restraints per residue

8.2 Residual restraint violations

This section provides the overview of the restraint violations analysis. The violations are binned as small, medium and large violations based on its absolute value. Average number of violations per model is calculated by dividing the total number of violations in each bin by the size of the ensemble.

8.2.1 Average number of distance violations per model

Distance violations less than 0.1 Å are not included in the calculation.

Bins (Å)	Average number of violations per model	Max (Å)
0.1-0.2 (Small)	402.4	0.2
0.2-0.5 (Medium)	400.6	0.5
>0.5 (Large)	0.8	0.62

8.2.2 Average number of dihedral-angle violations per model [i](#)

Dihedral-angle violations less than 1° are not included in the calculation.

Bins (°)	Average number of violations per model	Max (°)
1.0-10.0 (Small)	12.8	8.6
10.0-20.0 (Medium)	None	None
>20.0 (Large)	None	None

9 Distance violation analysis


9.1 Summary of distance violations

The following table shows the summary of distance violations in different restraint categories based on the sequence separation of the atoms involved. Each category is further sub-divided into three sub-categories based on the atoms involved. Violations less than 0.1 Å are not included in the statistics.

Restrains type	Count	% ¹	Violated ³			Consistently Violated ⁴		
			Count	% ²	% ¹	Count	% ²	% ¹
Intra-residue ($i-j =0$)	2206	46.6	701	31.8	14.8	329	14.9	6.9
Backbone-Backbone	139	2.9	33	23.7	0.7	29	20.9	0.6
Backbone-Sidechain	1100	23.2	345	31.4	7.3	113	10.3	2.4
Sidechain-Sidechain	967	20.4	323	33.4	6.8	187	19.3	3.9
Sequential ($i-j =1$)	960	20.3	213	22.2	4.5	54	5.6	1.1
Backbone-Backbone	331	7.0	43	13.0	0.9	25	7.6	0.5
Backbone-Sidechain	504	10.6	127	25.2	2.7	26	5.2	0.5
Sidechain-Sidechain	125	2.6	43	34.4	0.9	3	2.4	0.1
Medium range ($i-j >1$ & $i-j <5$)	627	13.2	186	29.7	3.9	28	4.5	0.6
Backbone-Backbone	136	2.9	25	18.4	0.5	5	3.7	0.1
Backbone-Sidechain	276	5.8	81	29.3	1.7	12	4.3	0.3
Sidechain-Sidechain	215	4.5	80	37.2	1.7	11	5.1	0.2
Long range ($i-j \geq 5$)	858	18.1	314	36.6	6.6	20	2.3	0.4
Backbone-Backbone	91	1.9	25	27.5	0.5	7	7.7	0.1
Backbone-Sidechain	297	6.3	95	32.0	2.0	6	2.0	0.1
Sidechain-Sidechain	470	9.9	194	41.3	4.1	7	1.5	0.1
Inter-chain	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Hydrogen bond	86	1.8	12	14.0	0.3	0	0.0	0.0
Disulfide bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Total	4737	100.0	1426	30.1	30.1	431	9.1	9.1
Backbone-Backbone	783	16.5	138	17.6	2.9	66	8.4	1.4
Backbone-Sidechain	2177	46.0	648	29.8	13.7	157	7.2	3.3
Sidechain-Sidechain	1777	37.5	640	36.0	13.5	208	11.7	4.4

¹ percentage calculated with respect to the total number of distance restraints, ² percentage calculated with respect to the number of restraints in a particular restraint category, ³ violated in at least one model, ⁴ violated in all the models

9.1.1 Bar chart : Distribution of distance restraints and violations [i](#)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories. The hydrogen bonds and disulfid bonds are counted in their appropriate category on the x-axis

9.2 Distance violation statistics for each model [i](#)

The following table provides the distance violation statistics for each model in the ensemble. Violations less than 0.1 Å are not included in the statistics.

Model ID	Number of violations						Mean (Å)	Max (Å)	SD ⁶ (Å)	Median (Å)
	IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total				
1	481	112	91	102	0	786	0.22	0.52	0.09	0.2
2	496	114	84	123	0	817	0.22	0.62	0.1	0.2
3	488	115	87	110	0	800	0.22	0.56	0.09	0.2
4	481	108	86	117	0	792	0.22	0.49	0.09	0.21
5	501	114	79	99	0	793	0.22	0.5	0.09	0.21
6	474	121	88	112	0	795	0.23	0.49	0.09	0.21
7	480	117	83	120	0	800	0.22	0.59	0.09	0.21
8	487	119	90	125	0	821	0.22	0.51	0.09	0.2
9	483	117	94	99	0	793	0.22	0.54	0.09	0.2
10	483	125	91	113	0	812	0.22	0.5	0.09	0.21
11	483	119	85	113	0	800	0.22	0.56	0.09	0.21


Continued on next page...

Continued from previous page...

Model ID	Number of violations					Total	Mean (Å)	Max (Å)	SD ⁶ (Å)	Median (Å)
	IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵					
12	499	115	89	104	0	807	0.22	0.51	0.09	0.2
13	485	109	85	115	0	794	0.23	0.49	0.09	0.21
14	489	108	86	109	0	792	0.22	0.55	0.09	0.2
15	489	120	94	131	0	834	0.22	0.51	0.09	0.2
16	485	112	87	120	0	804	0.22	0.5	0.09	0.2
17	498	122	85	110	0	815	0.22	0.55	0.09	0.21
18	479	118	98	117	0	812	0.22	0.49	0.09	0.2
19	493	106	91	111	0	801	0.23	0.61	0.09	0.21
20	498	109	85	115	0	807	0.22	0.5	0.09	0.21

¹Intra-residue restraints, ²Sequential restraints, ³Medium range restraints, ⁴Long range restraints, ⁵Inter-chain restraints, ⁶Standard deviation

9.2.1 Bar graph : Distance Violation statistics for each model [i](#)



The mean(dot),median(x) and the standard deviation are shown in blue with respect to the y axis on the right


9.3 Distance violation statistics for the ensemble

Violation analysis may find that some restraints are violated in few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of the ensemble. In total, 3237(IR:1505, SQ:747, MR:441, LR:544, IC:0) restraints are not violated in the ensemble.

Number of violated restraints						Fraction of the ensemble	
IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total	Count ⁶	%
44	23	31	72	0	170	1	5.0
31	14	21	37	0	103	2	10.0
30	15	5	37	0	87	3	15.0
24	8	13	20	0	65	4	20.0
25	7	8	15	0	55	5	25.0
16	12	7	9	0	44	6	30.0
20	7	8	9	0	44	7	35.0
18	6	9	9	0	42	8	40.0
13	13	2	12	0	40	9	45.0
9	9	4	7	0	29	10	50.0
11	6	4	5	0	26	11	55.0
14	8	3	10	0	35	12	60.0
16	4	5	8	0	33	13	65.0
21	3	5	4	0	33	14	70.0
14	2	4	6	0	26	15	75.0
14	3	11	8	0	36	16	80.0
16	5	3	7	0	31	17	85.0
13	4	2	8	0	27	18	90.0
23	10	13	11	0	57	19	95.0
329	54	28	20	0	431	20	100.0

¹Intra-residue restraints, ²Sequential restraints, ³Medium range restraints, ⁴Long range restraints, ⁵Inter-chain restraints, ⁶ Number of models with violations


9.3.1 Bar graph : Distance violation statistics for the ensemble [i](#)



9.4 Most violated distance restraints in the ensemble [i](#)

9.4.1 Histogram : Distribution of mean distance violations [i](#)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models in the ensemble



9.4.2 Table: Most violated distance restraints [i](#)

The following table provides the mean and the standard deviation of the violations for the 10 worst performing restraints, sorted by number of violated models and the mean violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.


Key	Atom-1	Atom-2	Models ¹	Mean (Å)	SD ¹ (Å)	Median (Å)
(4,236)	1:A:139:PRO:HB3	1:A:139:PRO:HB2	20	0.49	0.0	0.49
(4,880)	1:A:104:ASP:HB2	1:A:104:ASP:HB3	20	0.49	0.0	0.49
(4,598)	1:A:64:ARG:HB3	1:A:64:ARG:HB2	20	0.48	0.01	0.49
(4,599)	1:A:64:ARG:HB2	1:A:64:ARG:HB3	20	0.48	0.01	0.49
(4,3104)	1:A:145:PRO:HG2	1:A:145:PRO:HD2	20	0.48	0.01	0.48
(4,3104)	1:A:145:PRO:HG2	1:A:145:PRO:HD3	20	0.48	0.01	0.48
(4,3104)	1:A:145:PRO:HG3	1:A:145:PRO:HD2	20	0.48	0.01	0.48
(4,3104)	1:A:145:PRO:HG3	1:A:145:PRO:HD3	20	0.48	0.01	0.48
(4,687)	1:A:82:TYR:HB3	1:A:82:TYR:HB2	20	0.48	0.0	0.48
(4,688)	1:A:82:TYR:HB2	1:A:82:TYR:HB3	20	0.48	0.0	0.48
(6,280)	1:A:142:PHE:HA	1:A:142:PHE:H	20	0.46	0.01	0.46
(6,439)	1:A:123:ASP:H	1:A:123:ASP:HA	20	0.45	0.01	0.45
(4,1030)	1:A:128:ILE:HG21	1:A:128:ILE:HB	20	0.44	0.0	0.44

¹Number of violated models, ²Standard deviation

9.5 All violated distance restraints [i](#)

9.5.1 Histogram : Distribution of distance violations [i](#)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.



9.5.2 Table : All distance violations [i](#)

The following table provides the 10 worst performing restraints, sorted by the violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(4,478)	1:A:39:LEU:HA	1:A:39:LEU:HD11	2	0.62
(4,478)	1:A:39:LEU:HA	1:A:39:LEU:HD12	2	0.62
(4,478)	1:A:39:LEU:HA	1:A:39:LEU:HD13	2	0.62
(8,81)	1:A:67:THR:H	1:A:63:TYR:HE1	19	0.61
(8,81)	1:A:67:THR:H	1:A:63:TYR:HE2	19	0.61
(6,66)	1:A:9:LYS:HD2	1:A:8:LEU:H	7	0.59
(6,66)	1:A:9:LYS:HD3	1:A:8:LEU:H	7	0.59
(4,1747)	1:A:39:LEU:HD11	1:A:51:GLU:HG2	7	0.59
(4,1747)	1:A:39:LEU:HD11	1:A:51:GLU:HG3	7	0.59
(4,1747)	1:A:39:LEU:HD12	1:A:51:GLU:HG2	7	0.59
(4,1747)	1:A:39:LEU:HD12	1:A:51:GLU:HG3	7	0.59

Continued on next page...

Continued from previous page...

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(4,1747)	1:A:39:LEU:HD13	1:A:51:GLU:HG2	7	0.59
(4,1747)	1:A:39:LEU:HD13	1:A:51:GLU:HG3	7	0.59
(4,1747)	1:A:39:LEU:HD21	1:A:51:GLU:HG2	7	0.59
(4,1747)	1:A:39:LEU:HD21	1:A:51:GLU:HG3	7	0.59
(4,1747)	1:A:39:LEU:HD22	1:A:51:GLU:HG2	7	0.59
(4,1747)	1:A:39:LEU:HD22	1:A:51:GLU:HG3	7	0.59
(4,1747)	1:A:39:LEU:HD23	1:A:51:GLU:HG2	7	0.59
(4,1747)	1:A:39:LEU:HD23	1:A:51:GLU:HG3	7	0.59
(6,1194)	1:A:39:LEU:HD11	1:A:54:ASN:H	2	0.58
(6,1194)	1:A:39:LEU:HD12	1:A:54:ASN:H	2	0.58
(6,1194)	1:A:39:LEU:HD13	1:A:54:ASN:H	2	0.58
(6,1359)	1:A:108:LYS:HD2	1:A:105:ALA:H	3	0.56
(6,1359)	1:A:108:LYS:HD3	1:A:105:ALA:H	3	0.56
(4,1747)	1:A:39:LEU:HD11	1:A:51:GLU:HG2	11	0.56
(4,1747)	1:A:39:LEU:HD11	1:A:51:GLU:HG3	11	0.56
(4,1747)	1:A:39:LEU:HD12	1:A:51:GLU:HG2	11	0.56
(4,1747)	1:A:39:LEU:HD12	1:A:51:GLU:HG3	11	0.56
(4,1747)	1:A:39:LEU:HD13	1:A:51:GLU:HG2	11	0.56
(4,1747)	1:A:39:LEU:HD13	1:A:51:GLU:HG3	11	0.56
(4,1747)	1:A:39:LEU:HD21	1:A:51:GLU:HG2	11	0.56
(4,1747)	1:A:39:LEU:HD21	1:A:51:GLU:HG3	11	0.56
(4,1747)	1:A:39:LEU:HD22	1:A:51:GLU:HG2	11	0.56
(4,1747)	1:A:39:LEU:HD22	1:A:51:GLU:HG3	11	0.56
(4,1747)	1:A:39:LEU:HD23	1:A:51:GLU:HG2	11	0.56
(4,1747)	1:A:39:LEU:HD23	1:A:51:GLU:HG3	11	0.56
(6,148)	1:A:15:TYR:H	1:A:50:ARG:HD2	14	0.55
(6,148)	1:A:15:TYR:H	1:A:50:ARG:HD3	14	0.55
(6,148)	1:A:15:TYR:H	1:A:50:ARG:HD2	17	0.55
(6,148)	1:A:15:TYR:H	1:A:50:ARG:HD3	17	0.55
(6,200)	1:A:23:ARG:H	1:A:19:ASP:H	9	0.54

10 Dihedral-angle violation analysis [i](#)


10.1 Summary of dihedral-angle violations [i](#)

The following table provides the summary of dihedral-angle violations in different dihedral-angle types. Violations less than 1° are not included in the calculation.

Angle type	Count	% ¹	Violated ³			Consistently Violated ⁴		
			Count	% ²	% ¹	Count	% ²	% ¹
PSI	138	50.0	24	17.4	8.7	1	0.7	0.4
PHI	138	50.0	23	16.7	8.3	1	0.7	0.4
Total	276	100.0	47	17.0	17.0	2	0.7	0.7

¹ percentage calculated with respect to total number of dihedral-angle restraints, ² percentage calculated with respect to number of restraints in a particular dihedral-angle type, ³ violated in at least one model, ⁴ violated in all the models

10.1.1 Bar chart : Distribution of dihedral-angles and violations [i](#)




Violated and consistently violated restraints are shown using different hatch patterns in their respective categories

10.2 Dihedral-angle violation statistics for each model [i](#)

The following table provides the dihedral-angle violation statistics for each model in the ensemble. Violations less than 1° are not included in the statistics.

Model ID	Number of violations			Mean (°)	Max (°)	SD (°)	Median (°)
	PSI	PHI	Total				
1	3	5	8	1.99	2.5	0.34	2.05
2	6	5	11	1.75	3.2	0.65	1.4
3	11	6	17	2.05	3.8	0.76	1.8
4	9	7	16	1.94	3.9	0.8	1.65
5	9	4	13	1.91	2.9	0.46	1.9
6	8	8	16	2.12	4.3	0.88	2.0
7	6	7	13	1.91	3.7	0.72	1.8
8	6	9	15	2.01	3.8	0.76	1.8
9	7	4	11	2.95	3.8	0.56	2.8
10	6	4	10	2.19	2.9	0.59	2.45
11	4	6	10	2.62	4.7	0.9	2.4
12	8	7	15	2.15	4.0	0.88	1.7
13	6	10	16	1.96	3.8	0.71	1.6
14	6	5	11	1.98	3.8	0.82	2.0
15	7	4	11	2.37	4.1	0.89	2.2
16	4	6	10	2.12	3.0	0.74	2.0
17	8	6	14	1.88	2.9	0.51	1.85
18	7	6	13	1.99	3.0	0.7	2.0
19	7	6	13	2.88	8.6	1.91	2.4
20	7	5	12	2.3	4.5	1.0	2.05

10.2.1 Bar graph : Dihedral violation statistics for each model [i](#)



The mean(dot),median(x) and the standard deviation are shown in blue with respect to the y axis on the right

10.3 Dihedral-angle violation statistics for the ensemble [i](#)

Violation analysis may find that some restraints are violated in very few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of ensemble.

Number of violated restraints			Fraction of the ensemble	
PSI	PHI	Total	Count ¹	%
10	8	18	1	5.0
4	2	6	2	10.0
0	3	3	3	15.0
1	1	2	4	20.0
2	1	3	5	25.0
0	3	3	6	30.0
0	0	0	7	35.0
0	1	1	8	40.0
0	0	0	9	45.0
0	0	0	10	50.0
0	0	0	11	55.0


Continued on next page...

Continued from previous page...

Number of violated restraints			Fraction of the ensemble	
PSI	PHI	Total	Count ¹	%
2	0	2	12	60.0
2	1	3	13	65.0
0	1	1	14	70.0
0	0	0	15	75.0
1	0	1	16	80.0
1	1	2	17	85.0
0	0	0	18	90.0
0	0	0	19	95.0
1	1	2	20	100.0

¹ Number of models with violations

10.3.1 Bar graph : Dihedral-angle Violation statistics for the ensemble [i](#)




10.4 Most violated dihedral-angle restraints in the ensemble [i](#)

10.4.1 Histogram : Distribution of mean dihedral-angle violations [i](#)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models

in the ensemble



10.4.2 Table: Most violated dihedral-angle restraints [i](#)

The following table provides the mean and the standard deviation of the violations for the 10 worst performing restraints, sorted by number of violated models and the mean violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint.


Key	Atom-1	Atom-2	Atom-3	Atom-4	Models ¹	Mean	SD ²	Median
(1,110)	1:A:59:ILE:N	1:A:59:ILE:CA	1:A:59:ILE:C	1:A:60:PRO:N	20	2.75	0.41	2.8
(1,275)	1:A:143:SER:C	1:A:144:ARG:N	1:A:144:ARG:CA	1:A:144:ARG:C	20	2.68	0.78	2.55
(1,238)	1:A:135:PHE:N	1:A:135:PHE:CA	1:A:135:PHE:C	1:A:136:GLY:N	17	2.79	0.94	2.4
(1,117)	1:A:62:PRO:C	1:A:63:TYR:N	1:A:63:TYR:CA	1:A:63:TYR:C	17	2.24	0.45	2.4
(1,60)	1:A:32:TYR:N	1:A:32:TYR:CA	1:A:32:TYR:C	1:A:33:LYS:N	16	2.2	0.83	2.1
(1,63)	1:A:33:LYS:C	1:A:34:ASP:N	1:A:34:ASP:CA	1:A:34:ASP:C	14	1.58	0.41	1.5
(1,138)	1:A:73:CYS:N	1:A:73:CYS:CA	1:A:73:CYS:C	1:A:74:LEU:N	13	2.19	0.86	1.9
(1,239)	1:A:136:GLY:C	1:A:137:ASP:N	1:A:137:ASP:CA	1:A:137:ASP:C	13	2.06	0.67	1.8
(1,64)	1:A:34:ASP:N	1:A:34:ASP:CA	1:A:34:ASP:C	1:A:35:LEU:N	13	1.7	0.48	1.6
(1,268)	1:A:122:SER:N	1:A:122:SER:CA	1:A:122:SER:C	1:A:123:ASP:N	12	2.44	0.78	2.5

¹ Number of violated models, ²Standard deviation, All angle values are in degree (°)

10.5 All violated dihedral-angle restraints [i](#)

10.5.1 Histogram : Distribution of violations [i](#)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.



10.5.2 Table: All violated dihedral-angle restraints [i](#)

The following table provides the list of violations for the 10 worst performing restraints, sorted by the violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint.

Key	Atom-1	Atom-2	Atom-3	Atom-4	Model ID	Violation (°)
(1,124)	1:A:66:ASN:N	1:A:66:ASN:CA	1:A:66:ASN:C	1:A:67:THR:N	19	8.6
(1,238)	1:A:135:PHE:N	1:A:135:PHE:CA	1:A:135:PHE:C	1:A:136:GLY:N	11	4.7
(1,275)	1:A:143:SER:C	1:A:144:ARG:N	1:A:144:ARG:CA	1:A:144:ARG:C	20	4.5
(1,238)	1:A:135:PHE:N	1:A:135:PHE:CA	1:A:135:PHE:C	1:A:136:GLY:N	6	4.3
(1,125)	1:A:66:ASN:C	1:A:67:THR:N	1:A:67:THR:CA	1:A:67:THR:C	19	4.2
(1,138)	1:A:73:CYS:N	1:A:73:CYS:CA	1:A:73:CYS:C	1:A:74:LEU:N	15	4.1
(1,60)	1:A:32:TYR:N	1:A:32:TYR:CA	1:A:32:TYR:C	1:A:33:LYS:N	12	4.0
(1,275)	1:A:143:SER:C	1:A:144:ARG:N	1:A:144:ARG:CA	1:A:144:ARG:C	4	3.9
(1,238)	1:A:135:PHE:N	1:A:135:PHE:CA	1:A:135:PHE:C	1:A:136:GLY:N	19	3.9
(1,29)	1:A:16:LYS:C	1:A:17:TYR:N	1:A:17:TYR:CA	1:A:17:TYR:C	8	3.8