



wwPDB X-ray Structure Validation Summary Report ⓘ

May 26, 2020 – 11:48 pm BST

PDB ID : 5VPJ
Title : The crystal structure of a thioesterase from *Actinomadura verrucospora*.
Authors : Tan, K.; Joachimiak, G.; Endres, M.; Phillips Jr., G.N.; Joachmiak, A.; Midwest Center for Structural Genomics (MCSG); Enzyme Discovery for Natural Product Biosynthesis (NatPro)
Deposited on : 2017-05-05
Resolution : 2.35 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.









Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1164 (2.36-2.36)
Clashscore	141614	1232 (2.36-2.36)
Ramachandran outliers	138981	1211 (2.36-2.36)
Sidechain outliers	138945	1212 (2.36-2.36)
RSRZ outliers	127900	1150 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	159	 3% 81% 6% 13%
1	B	159	 6% 79% 8% 13%
1	C	159	 13% 79% 8% 13%
1	D	159	 3% 82% 5% 13%
1	E	159	 9% 75% 9% 15%
1	F	159	 4% 82% 5% 13%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	159	 <p>4% 78% 8% 14%</p>
1	H	159	 <p>8% 80% 5% 14%</p>
1	I	159	 <p>3% 77% 9% 14%</p>
1	J	159	 <p>8% 77% 7% 16%</p>
1	K	159	 <p>8% 73% 9% 18%</p>
1	L	159	 <p>3% 76% 11% 13%</p>

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 13045 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Thioesterase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	138	1085	691	182	205	4	3	0	0	0
1	B	138	1079	687	183	202	4	3	0	0	0
1	C	139	1058	677	177	197	4	3	0	0	0
1	D	138	1084	690	182	205	4	3	0	0	0
1	E	135	1049	672	176	194	4	3	0	0	0
1	F	139	1084	690	181	206	4	3	0	0	0
1	G	136	1070	682	178	203	4	3	0	0	0
1	H	136	1065	681	176	201	4	3	0	0	0
1	I	136	1070	684	178	201	4	3	0	0	0
1	J	133	1025	652	174	192	4	3	0	0	0
1	K	131	1022	654	171	190	4	3	0	0	0
1	L	139	1090	693	184	206	4	3	0	0	0

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	H	1	Total	Cl	0	0
			1	1		
2	G	1	Total	Cl	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Cl 1 1	0	0
2	K	2	Total Cl 2 2	0	0

- Molecule 3 is TETRAETHYLENE GLYCOL (three-letter code: PG4) (formula: C₈H₁₈O₅).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	D	1	Total C O 13 8 5	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	32	Total O 32 32	0	0
4	B	22	Total O 22 22	0	0
4	C	25	Total O 25 25	0	0
4	D	38	Total O 38 38	0	0
4	E	7	Total O 7 7	0	0
4	F	18	Total O 18 18	0	0

Continued on next page...



Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	G	34	Total 34	O 34	0	0
4	H	15	Total 15	O 15	0	0
4	I	26	Total 26	O 26	0	0
4	J	8	Total 8	O 8	0	0
4	K	10	Total 10	O 10	0	0
4	L	11	Total 11	O 11	0	0



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Thioesterase





- Molecule 1: Thioesterase




- Molecule 1: Thioesterase



- Molecule 1: Thioesterase




- Molecule 1: Thioesterase




PRO
HIS
GLY
ALA
LEU
ARG
PRO
ARG
ARG
PRO
ARG
ARG
PRO
ARG
ARG

• Molecule 1: Thioesterase




SER ASN ALA MSE R2 T12 A33 A40 L56 F86 Y96 R97 A103 A119 E120 ASP G122 L123 D137 P141 ASP PRO HIS GLY ALA LEU ARG PRO ARG ASP PRO ARG

• Molecule 1: Thioesterase



SER ASN ALA MSE ARG A3 T12 M17 L28 R35 E36 L39 M50 L72 L81 F86 Y98 T99 D100 G101 P102 Q111 A119 ASP GLY L123 P141 PRO HIS GLY ALA LEU ARG PRO ARG SER THR PRO ARG

• Molecule 1: Thioesterase



SER ASN ALA MSE R2 E6 F7 R8 T12 F13 A14 M17 L28 E36 D62 D63 L54 L72 F86 D100 G101 L117 R118 A119 ASP GLY LEU T124 P125 V126 L136 A140 P141 ASP PRO HIS GLY ALA LEU ARG PRO ARG SER THR PRO ARG

ARG

• Molecule 1: Thioesterase



SER ASN ALA MSE ARG A3 T12 L28 R37 F38 L39 M50 L56 C63 L72 L81 F86 T99 P102 G109 L117 R118 ALA GLU ASP L123 D137 P141 D142 PRO HIS GLY ALA LEU ARG PRO ARG ARG THR PRO ARG


• Molecule 1: Thioesterase



SER ASN ALA MSE ARG ALA Y4 T12 L28 R35 E36 A40 T46 V47 A46 H61 D62 ASP L54 S67 F93 D100 G101 P102 A103 R104 Q111 L117 R118 ALA GLU ASP L123 V126 E127 V128 P129 D130 E131 Y139 A140 P141 ASP PRO HIS GLY

ALA
LEU
ARG
PRO
THR
PRO
ARG


• Molecule 1: Thioesterase



SER ASN MSE ARG ALA Y4 F24 L28 R35 E36 L39 V47 A48 R49 M50 D63 L54 F65 L72 I84 D85 F86 H87 R97 V98 T99 D100 R104 Q111 G115 T116 L117 ARG ALA GLU ASP LEU T124 P125 V126 A140 ASP PRO HIS PRO

HIS
GLY
ALA
LEU
ARG
PRO
ARG
SER
GLU
THR
PRO
PRO
ARG

● Molecule 1: Thioesterase

Chain L:  3% 76% 11% 13%

SER ASN ALA M1 R2 L10 M17 L28 R35 E36 L39 A40 T46 M50 H51 D52 L56 E58 L81 F86 Q111 A119 GLU ASP G122 L123 T124 P125 L136 P141 ASP PRO HIS GLY ALA LEU ARG ARG SER GLU THR PRO PRO ARG

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	105.68Å 96.36Å 114.16Å 90.00° 99.16° 90.00°	Depositor
Resolution (Å)	33.68 – 2.35 33.68 – 2.35	Depositor EDS
% Data completeness (in resolution range)	86.8 (33.68-2.35) 86.8 (33.68-2.35)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.15 (at 2.34Å)	Xtrriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.186 , 0.227 0.188 , 0.227	Depositor DCC
R_{free} test set	4168 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	31.2	Xtrriage
Anisotropy	0.080	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 46.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	13045	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.17% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PG4, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.21	0/1106	0.36	0/1500
1	B	0.21	0/1100	0.36	0/1491
1	C	0.21	0/1079	0.38	0/1467
1	D	0.25	0/1105	0.38	0/1499
1	E	0.21	0/1070	0.37	0/1453
1	F	0.21	0/1105	0.36	0/1499
1	G	0.21	0/1091	0.36	0/1480
1	H	0.21	0/1086	0.37	0/1473
1	I	0.21	0/1091	0.36	0/1480
1	J	0.21	0/1044	0.37	0/1416
1	K	0.21	0/1042	0.36	0/1413
1	L	0.23	0/1111	0.37	0/1506
All	All	0.22	0/13030	0.37	0/17677

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1085	0	1037	6	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1079	0	1031	8	0
1	C	1058	0	992	8	0
1	D	1084	0	1038	5	0
1	E	1049	0	1000	8	0
1	F	1084	0	1031	4	0
1	G	1070	0	1024	7	0
1	H	1065	0	1018	5	0
1	I	1070	0	1028	10	0
1	J	1025	0	957	7	0
1	K	1022	0	968	8	0
1	L	1090	0	1045	10	0
2	A	1	0	0	0	0
2	G	1	0	0	0	0
2	H	1	0	0	0	0
2	K	2	0	0	0	0
3	D	13	0	18	1	0
4	A	32	0	0	0	0
4	B	22	0	0	0	0
4	C	25	0	0	0	0
4	D	38	0	0	0	0
4	E	7	0	0	0	0
4	F	18	0	0	0	0
4	G	34	0	0	0	0
4	H	15	0	0	0	0
4	I	26	0	0	0	0
4	J	8	0	0	0	0
4	K	10	0	0	0	0
4	L	11	0	0	0	0
All	All	13045	0	12187	67	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 67 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:39:LEU:HD13	1:E:50:MSE:HE1	1.75	0.68
1:G:35:ARG:NH1	1:G:111:GLN:OE1	2.28	0.67
1:K:28:LEU:HD13	1:L:28:LEU:HG	1.77	0.67
1:G:39:LEU:HD13	1:G:50:MSE:HE1	1.79	0.64
1:I:28:LEU:HD13	1:J:28:LEU:HG	1.78	0.64

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	134/159 (84%)	130 (97%)	3 (2%)	1 (1%)	22	23
1	B	134/159 (84%)	131 (98%)	2 (2%)	1 (1%)	22	23
1	C	135/159 (85%)	128 (95%)	5 (4%)	2 (2%)	10	8
1	D	134/159 (84%)	131 (98%)	2 (2%)	1 (1%)	22	23
1	E	131/159 (82%)	125 (95%)	5 (4%)	1 (1%)	19	20
1	F	135/159 (85%)	131 (97%)	3 (2%)	1 (1%)	22	23
1	G	132/159 (83%)	129 (98%)	2 (2%)	1 (1%)	19	20
1	H	132/159 (83%)	126 (96%)	5 (4%)	1 (1%)	19	20
1	I	132/159 (83%)	126 (96%)	5 (4%)	1 (1%)	19	20
1	J	127/159 (80%)	123 (97%)	4 (3%)	0	100	100
1	K	127/159 (80%)	124 (98%)	2 (2%)	1 (1%)	19	20
1	L	135/159 (85%)	131 (97%)	3 (2%)	1 (1%)	22	23
All	All	1588/1908 (83%)	1535 (97%)	41 (3%)	12 (1%)	19	20

5 of 12 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	86	PHE
1	B	86	PHE
1	I	86	PHE
1	L	86	PHE
1	C	86	PHE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	114/130 (88%)	114 (100%)	0	100	100
1	B	112/130 (86%)	112 (100%)	0	100	100
1	C	106/130 (82%)	106 (100%)	0	100	100
1	D	114/130 (88%)	113 (99%)	1 (1%)	78	87
1	E	108/130 (83%)	107 (99%)	1 (1%)	78	87
1	F	113/130 (87%)	111 (98%)	2 (2%)	59	70
1	G	113/130 (87%)	112 (99%)	1 (1%)	78	87
1	H	112/130 (86%)	110 (98%)	2 (2%)	59	70
1	I	113/130 (87%)	113 (100%)	0	100	100
1	J	104/130 (80%)	104 (100%)	0	100	100
1	K	105/130 (81%)	103 (98%)	2 (2%)	57	68
1	L	114/130 (88%)	111 (97%)	3 (3%)	46	56
All	All	1328/1560 (85%)	1316 (99%)	12 (1%)	78	87

5 of 12 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	12	THR
1	H	17	ASN
1	L	2	ARG
1	G	17	ASN
1	K	115	CYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	K	17	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 5 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	PG4	D	201	-	12,12,12	0.68	0	11,11,11	0.24	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PG4	D	201	-	-	4/10/10/10	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	D	201	PG4	O3-C5-C6-O4
3	D	201	PG4	O4-C7-C8-O5
3	D	201	PG4	O2-C3-C4-O3
3	D	201	PG4	C6-C5-O3-C4

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	201	PG4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	135/159 (84%)	-0.19	4 (2%) 50 61	17, 32, 84, 117	0
1	B	135/159 (84%)	0.39	10 (7%) 14 22	20, 43, 94, 117	0
1	C	136/159 (85%)	0.59	20 (14%) 2 3	17, 45, 109, 150	0
1	D	134/159 (84%)	-0.08	4 (2%) 50 61	18, 32, 78, 110	0
1	E	132/159 (83%)	0.52	15 (11%) 5 7	33, 63, 113, 143	0
1	F	136/159 (85%)	0.08	6 (4%) 34 46	22, 45, 89, 121	0
1	G	133/159 (83%)	0.13	7 (5%) 26 38	22, 34, 68, 136	0
1	H	133/159 (83%)	0.46	13 (9%) 7 12	21, 55, 111, 155	0
1	I	133/159 (83%)	-0.09	4 (3%) 50 61	23, 38, 76, 127	0
1	J	130/159 (81%)	0.53	12 (9%) 9 14	29, 62, 117, 135	0
1	K	128/159 (80%)	0.64	13 (10%) 6 11	35, 68, 123, 136	0
1	L	135/159 (84%)	0.11	4 (2%) 50 61	26, 50, 88, 130	0
All	All	1600/1908 (83%)	0.26	112 (7%) 16 24	17, 47, 107, 155	0

The worst 5 of 112 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	I	123	LEU	6.2
1	H	119	ALA	6.1
1	L	141	PRO	5.7
1	C	126	VAL	5.2
1	H	117	LEU	4.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	PG4	D	201	13/13	0.86	0.20	46,59,64,66	0
2	CL	G	201	1/1	0.88	0.44	85,85,85,85	0
2	CL	K	202	1/1	0.93	0.23	73,73,73,73	0
2	CL	A	201	1/1	0.93	0.30	84,84,84,84	0
2	CL	K	201	1/1	0.96	0.18	62,62,62,62	0
2	CL	H	201	1/1	0.98	0.15	35,35,35,35	1

6.5 Other polymers [i](#)

There are no such residues in this entry.