

wwPDB X-ray Structure Validation Summary Report (i)

Jan 17, 2023 - 05:14 PM EST

| PDB ID | : | 1VQY |
|--------------|---|---|
| Title | : | CRYSTAL STRUCTURE OF a NIPSNAP FAMILY PROTEIN (ATU5224) |
| | | FROM AGROBACTERIUM TUMEFACIENS STR. C58 AT 2.40 A RESO- |
| | | LUTION |
| Authors | : | Joint Center for Structural Genomics (JCSG) |
| Deposited on | | |
| Resolution | : | 2.40 Å(reported) |

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


| MolProbity | : | 4.02b-467 |
|--------------------------------|---|--|
| Mogul | : | 1.8.5 (274361), CSD as541be (2020) |
| Xtriage (Phenix) | : | 1.13 |
| EDS | : | 2.31.2 |
| Percentile statistics | : | 20191225.v01 (using entries in the PDB archive December 25th 2019) |
| Refmac | : | 5.8.0158 |
| CCP4 | : | 7.0.044 (Gargrove) |
| Ideal geometry (proteins) | : | Engh & Huber (2001) |
| Ideal geometry (DNA, RNA) | : | Parkinson et al. (1996) |
| Validation Pipeline (wwPDB-VP) | : | 2.31.2 |

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



| Metric | $\begin{array}{c} \textbf{Whole archive} \\ (\#\textbf{Entries}) \end{array}$ | ${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$ |
|-----------------------|---|---|
| R_{free} | 130704 | 3907 (2.40-2.40) |
| Clashscore | 141614 | 4398 (2.40-2.40) |
| Ramachandran outliers | 138981 | 4318 (2.40-2.40) |
| Sidechain outliers | 138945 | 4319 (2.40-2.40) |
| RSRZ outliers | 127900 | 3811 (2.40-2.40) |

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

| Mol | Chain | Length | Quality of chain | |
|-----|-------|--------|------------------|-------|
| 1 | А | 116 | 78% 12% | 9% |
| 1 | В | 116 | 91% | 7% •• |
| 1 | С | 116 | 82% 9% | 9% |
| 1 | D | 116 | % | 10% |
| 1 | Е | 116 | 76% 14% | 10% |



| Mol | Chain | Length | Quality of chain | | | |
|-----|-------|--------|-------------------|-----|---|----|
| 1 | F | 116 | 78% | 11% | • | 9% |
| 1 | G | 116 | 2% 7 9% | 10% | • | 9% |
| 1 | Н | 116 | 78% | 11% | · | 9% |



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 7283 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

| Mol | Chain | Residues | | A | toms | | | ZeroOcc | AltConf | Trace | |
|-----|-------|----------|-------|-------|------|-----|----|---------|---------|-------|---|
| 1 | 1 1 | 105 | Total | С | Ν | 0 | Se | 0 | 0 | 0 | |
| | А | 105 | 839 | 539 | 146 | 151 | 3 | 0 | 0 | 0 | |
| 1 | В | 114 | Total | С | Ν | Ο | Se | 0 | 0 | 0 | |
| | D | 114 | 924 | 590 | 166 | 165 | 3 | 0 | 0 | 0 | |
| 1 | С | 105 | Total | С | Ν | 0 | Se | 0 | 0 | 0 | |
| 1 | U | 105 | 841 | 540 | 147 | 151 | 3 | 0 | 0 | 0 | |
| 1 | D | 104 | Total | С | Ν | 0 | Se | 0 | 0 | 0 | 0 |
| | D | | 830 | 535 | 145 | 147 | 3 | | 0 | 0 | |
| 1 | Е | 104 | Total | С | Ν | Ο | Se | 0 | 0 | 0 | |
| | Ľ | 104 | 839 | 539 | 148 | 149 | 3 | | | | |
| 1 | F | 105 | Total | С | Ν | Ο | Se | 0 | 0 | 0 | |
| | Г | 105 | 843 | 541 | 149 | 150 | 3 | 0 | 0 | 0 | |
| 1 | С | G | 105 | Total | С | Ν | 0 | Se | 0 | 0 | 0 |
| | G | 105 | 852 | 545 | 151 | 153 | 3 | 0 | 0 | 0 | |
| 1 | Н | 105 | Total | С | Ν | Ο | Se | 0 | 0 | 0 | |
| 1 | 11 | 105 | 830 | 535 | 144 | 148 | 3 | 0 | | U | |

• Molecule 1 is a protein called hypothetical protein AGR_pAT_315.

There are 120 discrepancies between the modelled and reference sequences:

| Chain | Residue | Modelled | Actual | Comment | Reference |
|-------|---------|----------|--------|------------------|------------|
| А | -11 | MSE | - | expression tag | UNP Q8UK99 |
| А | -10 | GLY | - | expression tag | UNP Q8UK99 |
| А | -9 | SER | - | expression tag | UNP Q8UK99 |
| А | -8 | ASP | - | expression tag | UNP Q8UK99 |
| А | -7 | LYS | - | expression tag | UNP Q8UK99 |
| А | -6 | ILE | - | expression tag | UNP Q8UK99 |
| А | -5 | HIS | - | expression tag | UNP Q8UK99 |
| А | -4 | HIS | - | expression tag | UNP Q8UK99 |
| А | -3 | HIS | - | expression tag | UNP Q8UK99 |
| А | -2 | HIS | - | expression tag | UNP Q8UK99 |
| А | -1 | HIS | - | expression tag | UNP Q8UK99 |
| А | 0 | HIS | - | expression tag | UNP Q8UK99 |
| А | 1 | MSE | MET | modified residue | UNP Q8UK99 |



| Chain | Residue | Modelled | Actual | Comment | Reference |
|-------|---------|----------|--------|------------------|------------|
| А | 15 | MSE | MET | modified residue | UNP Q8UK99 |
| А | 54 | MSE | MET | modified residue | UNP Q8UK99 |
| В | -11 | MSE | - | expression tag | UNP Q8UK99 |
| В | -10 | GLY | - | expression tag | UNP Q8UK99 |
| В | -9 | SER | - | expression tag | UNP Q8UK99 |
| В | -8 | ASP | - | expression tag | UNP Q8UK99 |
| В | -7 | LYS | - | expression tag | UNP Q8UK99 |
| В | -6 | ILE | - | expression tag | UNP Q8UK99 |
| В | -5 | HIS | - | expression tag | UNP Q8UK99 |
| В | -4 | HIS | - | expression tag | UNP Q8UK99 |
| В | -3 | HIS | - | expression tag | UNP Q8UK99 |
| В | -2 | HIS | - | expression tag | UNP Q8UK99 |
| В | -1 | HIS | - | expression tag | UNP Q8UK99 |
| В | 0 | HIS | _ | expression tag | UNP Q8UK99 |
| В | 1 | MSE | MET | modified residue | UNP Q8UK99 |
| В | 15 | MSE | MET | modified residue | UNP Q8UK99 |
| В | 54 | MSE | MET | modified residue | UNP Q8UK99 |
| С | -11 | MSE | - | expression tag | UNP Q8UK99 |
| С | -10 | GLY | - | expression tag | UNP Q8UK99 |
| С | -9 | SER | - | expression tag | UNP Q8UK99 |
| С | -8 | ASP | - | expression tag | UNP Q8UK99 |
| С | -7 | LYS | - | expression tag | UNP Q8UK99 |
| С | -6 | ILE | - | expression tag | UNP Q8UK99 |
| С | -5 | HIS | - | expression tag | UNP Q8UK99 |
| С | -4 | HIS | - | expression tag | UNP Q8UK99 |
| С | -3 | HIS | - | expression tag | UNP Q8UK99 |
| С | -2 | HIS | - | expression tag | UNP Q8UK99 |
| С | -1 | HIS | - | expression tag | UNP Q8UK99 |
| С | 0 | HIS | - | expression tag | UNP Q8UK99 |
| С | 1 | MSE | MET | modified residue | UNP Q8UK99 |
| С | 15 | MSE | MET | modified residue | UNP Q8UK99 |
| С | 54 | MSE | MET | modified residue | UNP Q8UK99 |
| D | -11 | MSE | - | expression tag | UNP Q8UK99 |
| D | -10 | GLY | - | expression tag | UNP Q8UK99 |
| D | -9 | SER | - | expression tag | UNP Q8UK99 |
| D | -8 | ASP | - | expression tag | UNP Q8UK99 |
| D | -7 | LYS | - | expression tag | UNP Q8UK99 |
| D | -6 | ILE | - | expression tag | UNP Q8UK99 |
| D | -5 | HIS | - | expression tag | UNP Q8UK99 |
| D | -4 | HIS | - | expression tag | UNP Q8UK99 |
| D | -3 | HIS | - | expression tag | UNP Q8UK99 |
| D | -2 | HIS | _ | expression tag | UNP Q8UK99 |




| Chain | Residue | Modelled | Actual | Comment | Reference |
|-------|---------|----------|--------|------------------|------------|
| D | -1 | HIS | - | expression tag | UNP Q8UK99 |
| D | 0 | HIS | - | expression tag | UNP Q8UK99 |
| D | 1 | MSE | MET | modified residue | UNP Q8UK99 |
| D | 15 | MSE | MET | modified residue | UNP Q8UK99 |
| D | 54 | MSE | MET | modified residue | UNP Q8UK99 |
| Е | -11 | MSE | - | expression tag | UNP Q8UK99 |
| Е | -10 | GLY | _ | expression tag | UNP Q8UK99 |
| Е | -9 | SER | - | expression tag | UNP Q8UK99 |
| Е | -8 | ASP | - | expression tag | UNP Q8UK99 |
| Е | -7 | LYS | - | expression tag | UNP Q8UK99 |
| Е | -6 | ILE | - | expression tag | UNP Q8UK99 |
| Е | -5 | HIS | _ | expression tag | UNP Q8UK99 |
| Е | -4 | HIS | - | expression tag | UNP Q8UK99 |
| Е | -3 | HIS | - | expression tag | UNP Q8UK99 |
| Е | -2 | HIS | - | expression tag | UNP Q8UK99 |
| Е | -1 | HIS | - | expression tag | UNP Q8UK99 |
| Е | 0 | HIS | - | expression tag | UNP Q8UK99 |
| Е | 1 | MSE | MET | modified residue | UNP Q8UK99 |
| Е | 15 | MSE | MET | modified residue | UNP Q8UK99 |
| Е | 54 | MSE | MET | modified residue | UNP Q8UK99 |
| F | -11 | MSE | - | expression tag | UNP Q8UK99 |
| F | -10 | GLY | - | expression tag | UNP Q8UK99 |
| F | -9 | SER | - | expression tag | UNP Q8UK99 |
| F | -8 | ASP | _ | expression tag | UNP Q8UK99 |
| F | -7 | LYS | - | expression tag | UNP Q8UK99 |
| F | -6 | ILE | - | expression tag | UNP Q8UK99 |
| F | -5 | HIS | - | expression tag | UNP Q8UK99 |
| F | -4 | HIS | - | expression tag | UNP Q8UK99 |
| F | -3 | HIS | - | expression tag | UNP Q8UK99 |
| F | -2 | HIS | - | expression tag | UNP Q8UK99 |
| F | -1 | HIS | - | expression tag | UNP Q8UK99 |
| F | 0 | HIS | - | expression tag | UNP Q8UK99 |
| F | 1 | MSE | MET | modified residue | UNP Q8UK99 |
| F | 15 | MSE | MET | modified residue | UNP Q8UK99 |
| F | 54 | MSE | MET | modified residue | UNP Q8UK99 |
| G | -11 | MSE | - | expression tag | UNP Q8UK99 |
| G | -10 | GLY | - | expression tag | UNP Q8UK99 |
| G | -9 | SER | - | expression tag | UNP Q8UK99 |
| G | -8 | ASP | - | expression tag | UNP Q8UK99 |
| G | -7 | LYS | - | expression tag | UNP Q8UK99 |
| G | -6 | ILE | - | expression tag | UNP Q8UK99 |
| G | -5 | HIS | _ | expression tag | UNP Q8UK99 |



| Chain | Residue | Modelled | Actual | Comment | Reference |
|-------|---------|----------|--------|------------------|------------|
| G | -4 | HIS | - | expression tag | UNP Q8UK99 |
| G | -3 | HIS | - | expression tag | UNP Q8UK99 |
| G | -2 | HIS | - | expression tag | UNP Q8UK99 |
| G | -1 | HIS | - | expression tag | UNP Q8UK99 |
| G | 0 | HIS | - | expression tag | UNP Q8UK99 |
| G | 1 | MSE | MET | modified residue | UNP Q8UK99 |
| G | 15 | MSE | MET | modified residue | UNP Q8UK99 |
| G | 54 | MSE | MET | modified residue | UNP Q8UK99 |
| Н | -11 | MSE | - | expression tag | UNP Q8UK99 |
| Н | -10 | GLY | - | expression tag | UNP Q8UK99 |
| Н | -9 | SER | - | expression tag | UNP Q8UK99 |
| Н | -8 | ASP | - | expression tag | UNP Q8UK99 |
| Н | -7 | LYS | - | expression tag | UNP Q8UK99 |
| Н | -6 | ILE | - | expression tag | UNP Q8UK99 |
| Н | -5 | HIS | - | expression tag | UNP Q8UK99 |
| Н | -4 | HIS | - | expression tag | UNP Q8UK99 |
| Н | -3 | HIS | - | expression tag | UNP Q8UK99 |
| Н | -2 | HIS | - | expression tag | UNP Q8UK99 |
| Н | -1 | HIS | - | expression tag | UNP Q8UK99 |
| Н | 0 | HIS | - | expression tag | UNP Q8UK99 |
| Н | 1 | MSE | MET | modified residue | UNP Q8UK99 |
| Н | 15 | MSE | MET | modified residue | UNP Q8UK99 |
| Н | 54 | MSE | MET | modified residue | UNP Q8UK99 |

• Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O_4P).





| Mol | Chain | Residues | Atoms | | | ZeroOcc | AltConf |
|-----|-------|----------|------------|--------|--------|---------|---------|
| 2 | G | 1 | Total 5 | 0 4 | Р 1 | 0 | 0 |


• Molecule 3 is water.

| Mol | Chain | Residues | Atoms | ZeroOcc | AltConf |
|-----|-------|----------|---|---------|---------|
| 3 | А | 51 | Total O 51 51 | 0 | 0 |
| 3 | В | 62 | Total O 62 62 | 0 | 0 |
| 3 | С | 69 | Total O 69 69 | 0 | 0 |
| 3 | D | 63 | Total O 63 63 | 0 | 0 |
| 3 | Е | 58 | Total O 58 58 | 0 | 0 |
| 3 | F | 59 | Total O 59 59 | 0 | 0 |
| 3 | G | 48 | Total O 48 48 | 0 | 0 |
| 3 | Н | 70 | Total O 70 70 | 0 | 0 |



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: hypothetical protein AGR_pAT_315



| Chain F: | 78% | 11% • 9% |
|---|---|----------|
| MSE GLY SER ASP ASP LYS LYS HIS HIS HIS HIS HIS HIS TIS TIS TIS TIS TIS TIS TIS TIS TIS T | R11 R11 15 119 119 731 731 732 732 740 783 863 863 878 863 878 863 878 863 878 863 878 863 878 878 878 878 878 878 878 878 878 87 | |
| • Molecule 1: hypot | hetical protein AGR_pAT_315 | |
| Chain G: | 79% | 10% • 9% |
| MSE GLY SER ASP ASP LLYS LLYS HLS HLS HLS HLS HLS HLS HLS T T C U | R11 N15 L19 L19 138 138 138 849 849 849 849 849 051 N54 050 051 051 051 051 051 051 051 051 051 | |
| • Molecule 1: hypot | hetical protein AGR_pAT_315 | |
| Chain H: | 78% | 11% • 9% |
| MSE GLY SER SER SER ASP ASP HIS HIS HIS HIS HIS 12 V 12 V 12 V 12 V 12 V 12 V 12 V 12 | M15 1119 1119 1119 1119 1119 1119 1104 1104 | |



4 Data and refinement statistics (i)

| Property | Value | Source |
|---|---|-----------|
| Space group | P 21 21 21 | Depositor |
| Cell constants | 68.03Å 110.92Å 129.93Å | Depositor |
| a, b, c, α , β , γ | 90.00° 90.00° 90.00° | Depositor |
| Resolution (Å) | 29.08 - 2.40 | Depositor |
| Resolution (A) | 29.08 - 2.40 | EDS |
| % Data completeness | 97.5 (29.08-2.40) | Depositor |
| (in resolution range) | 97.6 (29.08-2.40) | EDS |
| R _{merge} | (Not available) | Depositor |
| R_{sym} | 0.15 | Depositor |
| $< I/\sigma(I) > 1$ | $3.00 (at 2.39 \text{\AA})$ | Xtriage |
| Refinement program | REFMAC 5.2.0005 | Depositor |
| D D. | 0.186 , 0.244 | Depositor |
| R, R_{free} | 0.197 , 0.253 | DCC |
| R_{free} test set | 1913 reflections (5.00%) | wwPDB-VP |
| Wilson B-factor $(Å^2)$ | 19.5 | Xtriage |
| Anisotropy | 0.392 | Xtriage |
| Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$ | 0.30, 32.4 | EDS |
| L-test for twinning ² | $ < L >=0.47, < L^2>=0.30$ | Xtriage |
| Estimated twinning fraction | No twinning to report. | Xtriage |
| F_o, F_c correlation | 0.92 | EDS |
| Total number of atoms | 7283 | wwPDB-VP |
| Average B, all atoms $(Å^2)$ | 21.0 | wwPDB-VP |

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 8.48% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Chain | Boi | nd lengths | Bo | nd angles |
|-------|-------|------|---------------|------|---------------|
| IVIOI | Unam | RMSZ | # Z > 5 | RMSZ | # Z > 5 |
| 1 | А | 0.60 | 0/852 | 0.66 | 0/1145 |
| 1 | В | 0.54 | 0/942 | 0.68 | 1/1266~(0.1%) |
| 1 | С | 0.61 | 0/854 | 0.70 | 0/1148 |
| 1 | D | 0.60 | 1/844~(0.1%) | 0.71 | 1/1137~(0.1%) |
| 1 | Е | 0.58 | 0/853 | 0.69 | 0/1146 |
| 1 | F | 0.63 | 0/856 | 0.77 | 1/1149~(0.1%) |
| 1 | G | 0.56 | 0/866 | 0.65 | 0/1163 |
| 1 | Н | 0.57 | 1/843~(0.1%) | 0.72 | 2/1134~(0.2%) |
| All | All | 0.59 | 2/6910~(0.0%) | 0.70 | 5/9288~(0.1%) |

All (2) bond length outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Ζ | Observed(Å) | $\mathrm{Ideal}(\mathrm{\AA})$ |
|-----|-------|-----|------|-------|-------|-------------|--------------------------------|
| 1 | D | 15 | MSE | SE-CE | -5.95 | 1.60 | 1.95 |
| 1 | Н | 15 | MSE | SE-CE | -5.61 | 1.62 | 1.95 |

All (5) bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Ζ | $Observed(^{o})$ | $Ideal(^{o})$ |
|-----|-------|-----|------|-----------|------|------------------|---------------|
| 1 | D | 63 | ARG | NE-CZ-NH1 | 6.57 | 123.58 | 120.30 |
| 1 | F | 63 | ARG | NE-CZ-NH1 | 5.55 | 123.07 | 120.30 |
| 1 | В | 63 | ARG | NE-CZ-NH1 | 5.37 | 122.98 | 120.30 |
| 1 | Н | 63 | ARG | NE-CZ-NH1 | 5.34 | 122.97 | 120.30 |
| 1 | Н | 66 | ARG | NE-CZ-NH1 | 5.17 | 122.88 | 120.30 |

There are no chirality outliers.

There are no planarity outliers.



5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 1 | А | 839 | 0 | 846 | 14 | 0 |
| 1 | В | 924 | 0 | 918 | 5 | 0 |
| 1 | С | 841 | 0 | 846 | 14 | 0 |
| 1 | D | 830 | 0 | 842 | 15 | 0 |
| 1 | Е | 839 | 0 | 858 | 17 | 0 |
| 1 | F | 843 | 0 | 858 | 12 | 0 |
| 1 | G | 852 | 0 | 862 | 14 | 0 |
| 1 | Н | 830 | 0 | 834 | 13 | 0 |
| 2 | G | 5 | 0 | 0 | 0 | 0 |
| 3 | А | 51 | 0 | 0 | 0 | 0 |
| 3 | В | 62 | 0 | 0 | 0 | 0 |
| 3 | С | 69 | 0 | 0 | 1 | 0 |
| 3 | D | 63 | 0 | 0 | 1 | 0 |
| 3 | Е | 58 | 0 | 0 | 0 | 0 |
| 3 | F | 59 | 0 | 0 | 1 | 0 |
| 3 | G | 48 | 0 | 0 | 0 | 0 |
| 3 | Н | 70 | 0 | 0 | 1 | 0 |
| All | All | 7283 | 0 | 6864 | 98 | 0 |

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 98 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

| Atom-1 | Atom-2 | Interatomic distance (Å) | Clash overlap (Å) |
|----------------|----------------|-----------------------------|----------------------|
| 1:E:11:ARG:HG3 | 1:E:88:GLU:HG2 | 1.64 | 0.80 |
| 1:H:15:MSE:HE3 | 1:H:19:LEU:HG | 1.66 | 0.78 |
| 1:A:15:MSE:HE1 | 1:A:51:VAL:CG1 | 2.15 | 0.77 |
| 1:G:15:MSE:HE3 | 1:G:19:LEU:HG | 1.70 | 0.73 |
| 1:D:3:VAL:HG11 | 1:D:54:MSE:HE3 | 1.72 | 0.70 |

There are no symmetry-related clashes.



5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Favoured | Allowed | Outliers | Perce | ntiles |
|-----|--------------|---------------|----------------|---------|----------|-------|--------|
| 1 | А | 103/116~(89%) | 101 (98%) | 2(2%) | 0 | 100 | 100 |
| 1 | В | 112/116~(97%) | 110~(98%) | 2(2%) | 0 | 100 | 100 |
| 1 | \mathbf{C} | 103/116~(89%) | 101 (98%) | 2(2%) | 0 | 100 | 100 |
| 1 | D | 102/116~(88%) | 100 (98%) | 2(2%) | 0 | 100 | 100 |
| 1 | Ε | 102/116~(88%) | 99~(97%) | 3~(3%) | 0 | 100 | 100 |
| 1 | F | 103/116~(89%) | $100 \ (97\%)$ | 2(2%) | 1 (1%) | 15 | 23 |
| 1 | G | 103/116~(89%) | 101 (98%) | 2(2%) | 0 | 100 | 100 |
| 1 | Н | 103/116~(89%) | 101 (98%) | 2(2%) | 0 | 100 | 100 |
| All | All | 831/928~(90%) | 813 (98%) | 17~(2%) | 1 (0%) | 51 | 68 |

All (1) Ramachandran outliers are listed below:

| Mol | Chain | \mathbf{Res} | Type |
|-----|-------|----------------|------|
| 1 | F | 1 | MSE |

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Rotameric | Outliers | Percentiles |
|-----|-------|-------------|-----------|----------|-------------|
| 1 | А | 86/97~(89%) | 84 (98%) | 2(2%) | 50 70 |
| 1 | В | 96/97~(99%) | 94 (98%) | 2(2%) | 53 72 |
| 1 | С | 86/97~(89%) | 84 (98%) | 2(2%) | 50 70 |
| 1 | D | 85/97~(88%) | 84 (99%) | 1 (1%) | 71 85 |



| Mol | Chain | Analysed | Rotameric | Outliers | Percentiles |
|-----|-------|---------------|-----------|----------|-------------|
| 1 | Ε | 87/97~(90%) | 85~(98%) | 2(2%) | 50 70 |
| 1 | F | 87/97~(90%) | 86~(99%) | 1 (1%) | 73 87 |
| 1 | G | 89/97~(92%) | 87~(98%) | 2(2%) | 52 71 |
| 1 | Н | 84/97~(87%) | 83~(99%) | 1 (1%) | 71 85 |
| All | All | 700/776~(90%) | 687~(98%) | 13 (2%) | 57 75 |

5 of 13 residues with a non-rotameric side chain are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1 | Ε | 40 | TYR |
| 1 | Ε | 49 | SER |
| 1 | Н | 40 | TYR |
| 1 | G | 49 | SER |
| 1 | G | 88 | GLU |

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1 | D | 53 | HIS |
| 1 | Н | 53 | HIS |

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

1 ligand is modelled in this entry.



In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol Type | Chain | Res | Link | Bond lengths | | | Bond angles | | | | |
|----------|-------|-------|------|--------------|--------|-------|-------------|--------|-------------|----------|---|
| | туре | Chain | nes | | Counts | RMSZ | # Z >2 | Counts | RMSZ | # Z > 2 | |
| | 2 | PO4 | G | 105 | - | 4,4,4 | 0.65 | 0 | $6,\!6,\!6$ | 0.61 | 0 |

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

| Mol | Chain | Analysed | <RSRZ $>$ | # RSRZ > 2 | $\mathbf{OWAB}(\mathbf{\AA}^2)$ | Q<0.9 |
|-----|-------|---------------|-----------|---------------|---------------------------------|-------|
| 1 | А | 102/116~(87%) | -0.07 | 1 (0%) 82 80 | 14, 20, 26, 36 | 0 |
| 1 | В | 111/116~(95%) | 0.11 | 8 (7%) 15 14 | 15, 20, 38, 45 | 0 |
| 1 | С | 102/116~(87%) | -0.33 | 0 100 100 | 15, 20, 26, 38 | 0 |
| 1 | D | 101/116~(87%) | -0.34 | 1 (0%) 82 80 | 15, 20, 25, 33 | 0 |
| 1 | Ε | 101/116~(87%) | -0.14 | 0 100 100 | 15, 20, 26, 30 | 0 |
| 1 | F | 102/116~(87%) | -0.21 | 0 100 100 | 16, 20, 25, 38 | 0 |
| 1 | G | 102/116~(87%) | -0.05 | 2 (1%) 65 63 | 15, 20, 26, 44 | 0 |
| 1 | Н | 102/116~(87%) | -0.32 | 0 100 100 | 14, 20, 25, 38 | 0 |
| All | All | 823/928~(88%) | -0.17 | 12 (1%) 73 72 | 14, 20, 28, 45 | 0 |

The worst 5 of 12 RSRZ outliers are listed below:

| Mol | Chain | Res | Type | RSRZ |
|-----|-------|-----|------|------|
| 1 | В | -3 | HIS | 5.0 |
| 1 | В | -4 | HIS | 4.0 |
| 1 | G | 0 | HIS | 3.8 |
| 1 | G | 61 | ASP | 3.3 |
| 1 | В | -8 | ASP | 3.3 |

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.



6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

| Mol | Type | Chain | Res | Atoms | RSCC | RSR | $\mathbf{B}	ext{-factors}(\mathrm{\AA}^2)$ | Q<0.9 |
|-----|------|-------|-----|-------|------|------|--|-------|
| 2 | PO4 | G | 105 | 5/5 | 0.78 | 0.36 | 44,46,48,61 | 0 |

6.5 Other polymers (i)

There are no such residues in this entry.

