



# wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 16, 2023 – 01:32 PM EDT

PDB ID : 1XK4  
Title : Crystal structure of human calprotectin(S100A8/S100A9)  
Authors : Korndoerfer, I.P.; Brueckner, F.; Skerra, A.  
Deposited on : 2004-09-26  
Resolution : 1.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	93	
1	B	93	
1	E	93	
1	F	93	
1	I	93	
1	J	93	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	C	113	 68% 9% 21%
2	D	113	 73% 7% 19%
2	G	113	 67% 13% 19%
2	H	113	 66% 12% 21%
2	K	113	 73% 8% 19%
2	L	113	 71% 9% 19%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	FLC	F	2008	-	-	X	-

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 9980 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Calgranulin A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	87	711	462	119	128	2	0	0	0
1	B	88	720	467	120	131	2	0	0	0
1	E	89	729	472	121	134	2	0	0	0
1	F	87	711	462	119	128	2	0	0	0
1	I	87	711	462	119	128	2	0	0	0
1	J	88	720	467	120	131	2	0	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	42	SER	CYS	engineered mutation	UNP P05109
B	42	SER	CYS	engineered mutation	UNP P05109
E	42	SER	CYS	engineered mutation	UNP P05109
F	42	SER	CYS	engineered mutation	UNP P05109
I	42	SER	CYS	engineered mutation	UNP P05109
J	42	SER	CYS	engineered mutation	UNP P05109

- Molecule 2 is a protein called Calgranulin B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	C	89	745	470	128	143	4	0	0	0
2	D	92	772	487	134	146	5	0	0	0
2	G	92	772	487	134	146	5	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	89	Total	C	N	O	S	0	0	0
			745	470	128	143	4			
2	K	92	Total	C	N	O	S	0	0	0
			772	487	134	146	5			
2	L	92	Total	C	N	O	S	0	0	0
			772	487	134	146	5			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	3	SER	CYS	engineered mutation	UNP P06702
D	3	SER	CYS	engineered mutation	UNP P06702
G	3	SER	CYS	engineered mutation	UNP P06702
H	3	SER	CYS	engineered mutation	UNP P06702
K	3	SER	CYS	engineered mutation	UNP P06702
L	3	SER	CYS	engineered mutation	UNP P06702

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Ca	0	0
			2	2		
3	B	2	Total	Ca	0	0
			2	2		
3	C	2	Total	Ca	0	0
			2	2		
3	D	2	Total	Ca	0	0
			2	2		
3	E	2	Total	Ca	0	0
			2	2		
3	F	2	Total	Ca	0	0
			2	2		
3	G	2	Total	Ca	0	0
			2	2		
3	H	2	Total	Ca	0	0
			2	2		
3	I	2	Total	Ca	0	0
			2	2		
3	J	2	Total	Ca	0	0
			2	2		
3	K	2	Total	Ca	0	0
			2	2		

*Continued on next page...*


Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	L	2	Total Ca 2 2	0	0

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Cl 1 1	0	0
4	B	1	Total Cl 1 1	0	0
4	F	1	Total Cl 1 1	0	0
4	J	1	Total Cl 1 1	0	0

- Molecule 5 is CITRATE ANION (three-letter code: FLC) (formula: C<sub>6</sub>H<sub>5</sub>O<sub>7</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	F	1	Total C O 13 6 7	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	84	Total O 84 84	0	0

Continued on next page...


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	78	Total 78	O 78	0	0
6	C	127	Total 127	O 127	0	0
6	D	84	Total 84	O 84	0	0
6	E	93	Total 93	O 93	0	0
6	F	86	Total 86	O 86	0	0
6	G	70	Total 70	O 70	0	0
6	H	89	Total 89	O 89	0	0
6	I	76	Total 76	O 76	0	0
6	J	88	Total 88	O 88	0	0
6	K	88	Total 88	O 88	0	0
6	L	96	Total 96	O 96	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Calgranulin A




- Molecule 1: Calgranulin A



- Molecule 1: Calgranulin A



- Molecule 1: Calgranulin A




- Molecule 1: Calgranulin A



- Molecule 1: Calgranulin A







- Molecule 2: Calgranulin B




- Molecule 2: Calgranulin B





- Molecule 2: Calgranulin B



- Molecule 2: Calgranulin B



- Molecule 2: Calgranulin B



- Molecule 2: Calgranulin B



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	48.51Å 78.70Å 177.41Å 90.00° 94.94° 90.00°	Depositor
Resolution (Å)	30.00 – 1.80 176.75 – 1.66	Depositor EDS
% Data completeness (in resolution range)	81.0 (30.00-1.80) 78.9 (176.75-1.66)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	0.00 (at 1.65Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.158 , 0.205 0.160 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	18.7	Xtrriage
Anisotropy	0.737	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 47.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	0.037 for h,-k,-h-l	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	9980	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	27.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.07% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, FLC, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.56	0/725	0.80	1/974 (0.1%)
1	B	0.55	0/734	0.81	2/986 (0.2%)
1	E	0.58	0/743	0.82	1/998 (0.1%)
1	F	0.56	0/725	0.82	1/974 (0.1%)
1	I	0.60	0/725	0.85	1/974 (0.1%)
1	J	0.59	0/734	0.86	2/986 (0.2%)
2	C	0.51	0/757	0.81	2/1014 (0.2%)
2	D	0.54	0/785	0.76	1/1050 (0.1%)
2	G	0.53	0/785	0.84	1/1050 (0.1%)
2	H	0.53	0/757	0.82	2/1014 (0.2%)
2	K	0.54	0/785	0.80	2/1050 (0.2%)
2	L	0.53	0/785	0.83	3/1050 (0.3%)
All	All	0.55	0/9040	0.82	19/12120 (0.2%)

There are no bond length outliers.

The worst 5 of 19 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	G	65	ASP	CB-CG-OD2	8.30	125.77	118.30
2	L	65	ASP	CB-CG-OD2	7.05	124.65	118.30
1	A	52	ASP	CB-CG-OD2	6.54	124.18	118.30
2	H	44	ASP	CB-CG-OD2	6.44	124.09	118.30
2	L	44	ASP	CB-CG-OD2	6.34	124.01	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	711	0	730	10	0
1	B	720	0	736	11	0
1	E	729	0	742	13	0
1	F	711	0	730	14	0
1	I	711	0	730	3	0
1	J	720	0	736	14	0
2	C	745	0	737	10	0
2	D	772	0	766	4	1
2	G	772	0	766	11	1
2	H	745	0	737	11	0
2	K	772	0	766	10	0
2	L	772	0	766	9	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
3	C	2	0	0	0	0
3	D	2	0	0	0	0
3	E	2	0	0	0	0
3	F	2	0	0	0	0
3	G	2	0	0	0	0
3	H	2	0	0	0	0
3	I	2	0	0	0	0
3	J	2	0	0	0	0
3	K	2	0	0	0	0
3	L	2	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	F	1	0	0	0	0
4	J	1	0	0	0	0
5	F	13	0	5	4	0
6	A	84	0	0	4	1
6	B	78	0	0	4	0
6	C	127	0	0	7	0
6	D	84	0	0	3	1
6	E	93	0	0	1	0
6	F	86	0	0	1	0
6	G	70	0	0	3	0
6	H	89	0	0	5	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	I	76	0	0	0	0
6	J	88	0	0	5	0
6	K	88	0	0	9	0
6	L	96	0	0	5	0
All	All	9980	0	8947	113	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 113 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:L:73:GLN:HG2	6:L:4617:HOH:O	1.27	1.27
2:G:13:GLU:HG3	6:G:3521:HOH:O	1.43	1.19
1:A:76:ILE:HG22	6:A:2014:HOH:O	1.47	1.11
2:K:57:LYS:HG2	6:K:3564:HOH:O	1.57	1.04
2:G:42:ARG:NH2	2:G:53:ASN:O	1.90	1.02

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:2041:HOH:O	6:D:4527:HOH:O[1_455]	2.08	0.12
2:D:4:LYS:CB	2:D:64:GLU:OE1[2_747]	2.14	0.06
2:G:4:LYS:CB	2:G:64:GLU:OE1[2_656]	2.14	0.06

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	85/93 (91%)	85 (100%)	0	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	86/93 (92%)	85 (99%)	1 (1%)	0	100	100
1	E	87/93 (94%)	86 (99%)	1 (1%)	0	100	100
1	F	85/93 (91%)	85 (100%)	0	0	100	100
1	I	85/93 (91%)	84 (99%)	1 (1%)	0	100	100
1	J	86/93 (92%)	85 (99%)	1 (1%)	0	100	100
2	C	87/113 (77%)	87 (100%)	0	0	100	100
2	D	90/113 (80%)	89 (99%)	1 (1%)	0	100	100
2	G	90/113 (80%)	90 (100%)	0	0	100	100
2	H	87/113 (77%)	87 (100%)	0	0	100	100
2	K	90/113 (80%)	90 (100%)	0	0	100	100
2	L	90/113 (80%)	90 (100%)	0	0	100	100
All	All	1048/1236 (85%)	1043 (100%)	5 (0%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	77/83 (93%)	75 (97%)	2 (3%)	46	32
1	B	78/83 (94%)	75 (96%)	3 (4%)	33	18
1	E	79/83 (95%)	77 (98%)	2 (2%)	47	34
1	F	77/83 (93%)	74 (96%)	3 (4%)	32	17
1	I	77/83 (93%)	75 (97%)	2 (3%)	46	32
1	J	78/83 (94%)	76 (97%)	2 (3%)	46	32
2	C	84/102 (82%)	81 (96%)	3 (4%)	35	20
2	D	87/102 (85%)	85 (98%)	2 (2%)	50	37
2	G	87/102 (85%)	85 (98%)	2 (2%)	50	37
2	H	84/102 (82%)	82 (98%)	2 (2%)	49	36

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	K	87/102 (85%)	84 (97%)	3 (3%)	37	22
2	L	87/102 (85%)	84 (97%)	3 (3%)	37	22
All	All	982/1110 (88%)	953 (97%)	29 (3%)	41	27

5 of 29 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	31	ARG
2	L	26	LEU
2	H	4	LYS
2	K	26	LEU
2	G	86	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
2	K	28	HIS
2	L	28	HIS
2	D	73	GLN
1	F	44	GLN
2	H	21	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 29 ligands modelled in this entry, 28 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	FLC	F	2008	-	12,12,12	1.09	0	17,17,17	2.16	5 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	FLC	F	2008	-	-	1/16/16/16	-

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	F	2008	FLC	CB-CA-CAC	-5.89	99.54	113.81
5	F	2008	FLC	CB-CG-CGC	-3.59	105.11	113.81
5	F	2008	FLC	OB2-CBC-CB	3.00	118.26	113.05
5	F	2008	FLC	OA1-CAC-CA	-2.48	115.69	122.94
5	F	2008	FLC	OB1-CBC-CB	-2.07	119.31	122.25

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	F	2008	FLC	CBC-CB-CG-CGC

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	F	2008	FLC	4	0



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	87/93 (93%)	-0.32	1 (1%) 80 78	15, 20, 38, 57	0
1	B	88/93 (94%)	-0.31	2 (2%) 60 56	16, 20, 40, 64	0
1	E	89/93 (95%)	-0.31	3 (3%) 45 39	17, 21, 40, 73	0
1	F	87/93 (93%)	-0.39	1 (1%) 80 78	16, 21, 38, 57	0
1	I	87/93 (93%)	-0.38	0 100 100	16, 20, 38, 57	0
1	J	88/93 (94%)	-0.35	1 (1%) 80 78	15, 20, 40, 60	0
2	C	89/113 (78%)	-0.28	0 100 100	16, 23, 39, 53	0
2	D	92/113 (81%)	-0.23	1 (1%) 80 78	18, 24, 41, 63	0
2	G	92/113 (81%)	-0.25	1 (1%) 80 78	20, 25, 41, 63	0
2	H	89/113 (78%)	-0.34	0 100 100	18, 24, 39, 53	0
2	K	92/113 (81%)	-0.31	2 (2%) 62 57	18, 23, 40, 64	0
2	L	92/113 (81%)	-0.27	1 (1%) 80 78	19, 24, 40, 65	0
All	All	1072/1236 (86%)	-0.31	13 (1%) 79 76	15, 23, 40, 73	0

The worst 5 of 13 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	K	95	HIS	3.5
1	E	1	MET	3.5
2	K	4	LYS	3.3
1	B	1	MET	3.3
2	L	95	HIS	3.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	FLC	F	2008	13/13	0.82	0.21	35,51,53,56	0
4	CL	B	2005	1/1	0.84	0.07	74,74,74,74	0
4	CL	A	2004	1/1	0.85	0.07	75,75,75,75	0
4	CL	F	2006	1/1	0.87	0.11	74,74,74,74	0
4	CL	J	2007	1/1	0.89	0.10	66,66,66,66	0
3	CA	J	2522	1/1	0.99	0.04	19,19,19,19	1
3	CA	K	3522	1/1	0.99	0.09	21,21,21,21	1
3	CA	B	2502	1/1	0.99	0.06	24,24,24,24	1
3	CA	C	3502	1/1	0.99	0.08	22,22,22,22	1
3	CA	G	3511	1/1	0.99	0.03	18,18,18,18	0
3	CA	G	3512	1/1	0.99	0.06	21,21,21,21	1
3	CA	H	4512	1/1	0.99	0.07	26,26,26,26	1
3	CA	B	2501	1/1	1.00	0.04	15,15,15,15	0
3	CA	A	1501	1/1	1.00	0.05	16,16,16,16	0
3	CA	H	4511	1/1	1.00	0.03	19,19,19,19	0
3	CA	C	3501	1/1	1.00	0.02	19,19,19,19	0
3	CA	I	1521	1/1	1.00	0.03	15,15,15,15	0
3	CA	I	1522	1/1	1.00	0.04	19,19,19,19	1
3	CA	J	2521	1/1	1.00	0.04	14,14,14,14	0
3	CA	A	1502	1/1	1.00	0.07	22,22,22,22	1
3	CA	K	3521	1/1	1.00	0.03	18,18,18,18	0
3	CA	D	4501	1/1	1.00	0.04	18,18,18,18	0
3	CA	L	4521	1/1	1.00	0.03	19,19,19,19	0
3	CA	L	4522	1/1	1.00	0.15	23,23,23,23	1
3	CA	D	4502	1/1	1.00	0.06	21,21,21,21	1
3	CA	E	1511	1/1	1.00	0.03	15,15,15,15	0
3	CA	E	1512	1/1	1.00	0.06	23,23,23,23	1
3	CA	F	2511	1/1	1.00	0.04	17,17,17,17	0
3	CA	F	2512	1/1	1.00	0.09	21,21,21,21	1

## 6.5 Other polymers [i](#)

There are no such residues in this entry.