



wwPDB X-ray Structure Validation Summary Report ⓘ

May 23, 2020 – 05:55 pm BST

PDB ID : 4XKU
Title : E coli BFR variant Y114F
Authors : Hemmings, A.M.; Le Brun, N.E.; Bradley, J.M.
Deposited on : 2015-01-12
Resolution : 1.78 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.78 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	9185 (1.80-1.76)
Clashscore	141614	10184 (1.80-1.76)
Ramachandran outliers	138981	10051 (1.80-1.76)
Sidechain outliers	138945	10050 (1.80-1.76)
RSRZ outliers	127900	9032 (1.80-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	158	
1	B	158	
1	C	158	
1	D	158	
1	E	158	
1	F	158	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	158	 <p>% 89% 11%</p>
1	H	158	 <p>% 94% 6%</p>
1	I	158	 <p>% 92% 8%</p>
1	J	158	 <p>% 91% 9%</p>
1	K	158	 <p>87% 13%</p>
1	L	158	 <p>89% 11%</p>

2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 19042 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Bacterioferritin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	158	1333	842	226	256	9	0	8	1
1	B	158	1345	853	226	257	9	0	10	1
1	C	158	1330	842	225	256	7	0	7	1
1	D	158	1328	840	223	256	9	0	8	1
1	E	158	1331	842	225	255	9	0	7	1
1	F	158	1337	844	231	254	8	0	8	1
1	G	158	1325	838	224	254	9	0	7	1
1	H	158	1321	833	225	256	7	0	7	1
1	I	158	1337	843	226	260	8	0	9	1
1	J	158	1329	841	228	252	8	0	7	1
1	K	158	1322	836	222	255	9	0	7	1
1	L	158	1342	850	228	255	9	0	10	1

There are 12 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	114	PHE	TYR	engineered mutation	UNP P0ABD3
B	114	PHE	TYR	engineered mutation	UNP P0ABD3
C	114	PHE	TYR	engineered mutation	UNP P0ABD3
D	114	PHE	TYR	engineered mutation	UNP P0ABD3
E	114	PHE	TYR	engineered mutation	UNP P0ABD3

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference					
F	114	PHE	TYR	engineered mutation	UNP P0ABD3					
G	114	PHE	TYR	engineered mutation	UNP P0ABD3					
H	114	PHE	TYR	engineered mutation	UNP P0ABD3					
I	114	PHE	TYR	engineered mutation	UNP P0ABD3					
J	114	PHE	TYR	engineered mutation	UNP P0ABD3					
K	114	PHE	TYR	engineered mutation </tr <tr> <td>L</td> <td>114</td> <td>PHE</td> <td>TYR</td> <td>engineered mutation</td> <td>UNP P0ABD3</td> </tr>	L	114	PHE	TYR	engineered mutation	UNP P0ABD3
L	114	PHE	TYR	engineered mutation	UNP P0ABD3					

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	D	1	Total O S 5 4 1	0	0
2	E	1	Total O S 5 4 1	0	0
2	E	1	Total O S 5 4 1	0	0
2	E	1	Total O S 5 4 1	0	0
2	F	1	Total O S 5 4 1	0	0
2	F	1	Total O S 5 4 1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	G	1	Total O S 5 4 1	0	0
2	H	1	Total O S 5 4 1	0	0
2	H	1	Total O S 5 4 1	0	0
2	H	1	Total O S 5 4 1	0	0
2	K	1	Total O S 5 4 1	0	0
2	L	1	Total O S 5 4 1	0	0
2	L	1	Total O S 5 4 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	234	Total O 234 234	0	0
3	B	243	Total O 243 243	0	0
3	C	234	Total O 234 234	0	0
3	D	231	Total O 231 231	0	0
3	E	271	Total O 271 271	0	0
3	F	273	Total O 273 273	0	0
3	G	255	Total O 255 255	0	0
3	H	259	Total O 259 259	0	0
3	I	241	Total O 241 241	0	0
3	J	260	Total O 260 260	0	0
3	K	243	Total O 243 243	0	0
3	L	243	Total O 243 243	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Bacterioferritin




- Molecule 1: Bacterioferritin




- Molecule 1: Bacterioferritin




- Molecule 1: Bacterioferritin




- Molecule 1: Bacterioferritin




- Molecule 1: Bacterioferritin




- Molecule 1: Bacterioferritin




- Molecule 1: Bacterioferritin





- Molecule 1: Bacterioferritin





- Molecule 1: Bacterioferritin



- Molecule 1: Bacterioferritin



- Molecule 1: Bacterioferritin



4 Data and refinement statistics

Property	Value	Source
Space group	P 42 21 2	Depositor
Cell constants a, b, c, α , β , γ	207.56Å 207.56Å 142.59Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	102.27 – 1.78 102.27 – 1.78	Depositor EDS
% Data completeness (in resolution range)	95.9 (102.27-1.78) 96.2 (102.27-1.78)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.93 (at 1.78Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.8.2_1309)	Depositor
R, R_{free}	0.188 , 0.217 0.188 , 0.217	Depositor DCC
R_{free} test set	14159 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	14.9	Xtrriage
Anisotropy	0.004	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 49.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	19042	wwPDB-VP
Average B, all atoms (Å ²)	11.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 46.20 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.1847e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.32	0/1375	0.45	0/1850
1	B	0.33	0/1396	0.46	0/1878
1	C	0.34	0/1369	0.46	0/1843
1	D	0.33	0/1372	0.47	0/1846
1	E	0.34	0/1372	0.46	0/1844
1	F	0.33	0/1378	0.45	0/1854
1	G	0.34	0/1363	0.47	0/1834
1	H	0.34	0/1363	0.49	0/1837
1	I	0.33	0/1381	0.45	0/1859
1	J	0.33	0/1371	0.46	0/1842
1	K	0.34	0/1362	0.46	0/1833
1	L	0.33	0/1393	0.44	0/1872
All	All	0.33	0/16495	0.46	0/22192

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1333	0	1310	14	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1345	0	1332	21	0
1	C	1330	0	1303	10	0
1	D	1328	0	1306	16	0
1	E	1331	0	1319	13	0
1	F	1337	0	1320	14	0
1	G	1325	0	1304	15	0
1	H	1321	0	1287	8	0
1	I	1337	0	1311	14	0
1	J	1329	0	1317	12	0
1	K	1322	0	1306	16	0
1	L	1342	0	1339	14	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
2	D	5	0	0	0	0
2	E	15	0	0	0	0
2	F	10	0	0	0	0
2	G	5	0	0	0	0
2	H	15	0	0	0	0
2	K	5	0	0	1	0
2	L	10	0	0	0	0
3	A	234	0	0	7	1
3	B	243	0	0	6	0
3	C	234	0	0	2	1
3	D	231	0	0	5	0
3	E	271	0	0	1	2
3	F	273	0	0	8	0
3	G	255	0	0	5	2
3	H	259	0	0	4	0
3	I	241	0	0	2	1
3	J	260	0	0	6	0
3	K	243	0	0	3	1
3	L	243	0	0	3	0
All	All	19042	0	15754	159	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 159 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:I:73:ASP:O	3:I:201:HOH:O	1.97	0.83

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:50[B]:ASP:OD1	3:L:301:HOH:O	2.03	0.76
3:A:530:HOH:O	1:B:31[B]:MET:SD	2.41	0.76
1:G:158:GLY:N	3:G:302:HOH:O	2.22	0.71
1:I:27:LEU:HG	1:I:31[B]:MET:HE2	1.73	0.71

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:490:HOH:O	3:E:469:HOH:O[8_776]	2.14	0.06
3:E:460:HOH:O	3:I:222:HOH:O[8_776]	2.14	0.06
3:C:324:HOH:O	3:G:490:HOH:O[8_776]	2.16	0.04
3:G:476:HOH:O	3:K:493:HOH:O[8_776]	2.18	0.02

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	164/158 (104%)	163 (99%)	1 (1%)	0	100	100
1	B	166/158 (105%)	165 (99%)	1 (1%)	0	100	100
1	C	163/158 (103%)	162 (99%)	1 (1%)	0	100	100
1	D	164/158 (104%)	163 (99%)	1 (1%)	0	100	100
1	E	163/158 (103%)	162 (99%)	1 (1%)	0	100	100
1	F	164/158 (104%)	163 (99%)	1 (1%)	0	100	100
1	G	163/158 (103%)	163 (100%)	0	0	100	100
1	H	163/158 (103%)	162 (99%)	1 (1%)	0	100	100
1	I	165/158 (104%)	165 (100%)	0	0	100	100
1	J	163/158 (103%)	163 (100%)	0	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	K	163/158 (103%)	163 (100%)	0	0	100	100
1	L	166/158 (105%)	165 (99%)	1 (1%)	0	100	100
All	All	1967/1896 (104%)	1959 (100%)	8 (0%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	145/139 (104%)	145 (100%)	0	100	100
1	B	147/139 (106%)	147 (100%)	0	100	100
1	C	143/139 (103%)	143 (100%)	0	100	100
1	D	144/139 (104%)	143 (99%)	1 (1%)	84	79
1	E	145/139 (104%)	144 (99%)	1 (1%)	84	79
1	F	145/139 (104%)	145 (100%)	0	100	100
1	G	143/139 (103%)	142 (99%)	1 (1%)	84	79
1	H	143/139 (103%)	142 (99%)	1 (1%)	84	79
1	I	146/139 (105%)	146 (100%)	0	100	100
1	J	145/139 (104%)	143 (99%)	2 (1%)	67	56
1	K	144/139 (104%)	144 (100%)	0	100	100
1	L	148/139 (106%)	148 (100%)	0	100	100
All	All	1738/1668 (104%)	1732 (100%)	6 (0%)	91	90

5 of 6 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	77	LEU
1	J	154	ILE
1	H	77	LEU
1	E	99	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	J	50	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	E	151	GLN
1	L	17	ASN
1	L	100	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

15 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	B	201	-	4,4,4	0.17	0	6,6,6	0.20	0
2	SO4	A	201	-	4,4,4	0.82	0	6,6,6	0.81	0
2	SO4	D	201	-	4,4,4	0.14	0	6,6,6	0.10	0
2	SO4	K	201	-	4,4,4	0.20	0	6,6,6	0.19	0
2	SO4	F	201	-	4,4,4	0.13	0	6,6,6	0.15	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	SO4	E	201	-	4,4,4	0.11	0	6,6,6	0.21	0
2	SO4	H	203	-	4,4,4	0.21	0	6,6,6	0.17	0
2	SO4	E	202	-	4,4,4	0.11	0	6,6,6	0.23	0
2	SO4	E	203	-	4,4,4	0.12	0	6,6,6	0.21	0
2	SO4	H	201	-	4,4,4	0.18	0	6,6,6	0.20	0
2	SO4	L	201	-	4,4,4	0.21	0	6,6,6	0.21	0
2	SO4	H	202	-	4,4,4	0.13	0	6,6,6	0.19	0
2	SO4	F	202	-	4,4,4	0.19	0	6,6,6	0.19	0
2	SO4	L	202	-	4,4,4	0.16	0	6,6,6	0.13	0
2	SO4	G	201	-	4,4,4	0.17	0	6,6,6	0.21	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	K	201	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	158/158 (100%)	-0.17	2 (1%) 77 77	2, 7, 20, 43	0
1	B	158/158 (100%)	-0.21	1 (0%) 89 89	2, 7, 18, 39	0
1	C	158/158 (100%)	-0.20	1 (0%) 89 89	2, 6, 18, 44	0
1	D	158/158 (100%)	-0.16	1 (0%) 89 89	2, 7, 19, 46	0
1	E	158/158 (100%)	-0.14	1 (0%) 89 89	2, 7, 21, 47	0
1	F	158/158 (100%)	-0.11	2 (1%) 77 77	3, 7, 19, 60	0
1	G	158/158 (100%)	-0.17	2 (1%) 77 77	2, 6, 21, 53	0
1	H	158/158 (100%)	-0.16	2 (1%) 77 77	2, 7, 19, 44	0
1	I	158/158 (100%)	-0.21	1 (0%) 89 89	2, 7, 18, 41	0
1	J	158/158 (100%)	-0.19	1 (0%) 89 89	2, 7, 19, 49	0
1	K	158/158 (100%)	-0.19	0 100 100	2, 7, 19, 50	0
1	L	158/158 (100%)	-0.23	0 100 100	2, 7, 20, 38	0
All	All	1896/1896 (100%)	-0.18	14 (0%) 87 88	2, 7, 20, 60	0

The worst 5 of 14 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	158	GLY	14.9
1	E	158	GLY	11.6
1	H	158	GLY	7.3
1	C	158	GLY	6.0
1	G	158	GLY	4.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	SO4	A	201	5/5	0.93	0.14	33,34,45,53	0
2	SO4	E	203	5/5	0.93	0.18	30,32,36,47	0
2	SO4	E	202	5/5	0.94	0.13	26,30,42,47	0
2	SO4	F	201	5/5	0.95	0.13	26,31,45,48	0
2	SO4	H	203	5/5	0.95	0.19	30,30,36,48	0
2	SO4	H	202	5/5	0.95	0.11	27,28,38,41	0
2	SO4	F	202	5/5	0.95	0.16	19,28,51,55	0
2	SO4	L	201	5/5	0.96	0.15	23,26,41,42	0
2	SO4	K	201	5/5	0.96	0.14	26,29,37,50	0
2	SO4	L	202	5/5	0.96	0.14	30,31,37,44	0
2	SO4	D	201	5/5	0.97	0.10	23,32,40,43	0
2	SO4	B	201	5/5	1.00	0.08	5,5,7,9	0
2	SO4	H	201	5/5	1.00	0.08	6,7,7,8	0
2	SO4	E	201	5/5	1.00	0.09	6,7,8,8	0
2	SO4	G	201	5/5	1.00	0.08	5,6,7,9	0

6.5 Other polymers [i](#)

There are no such residues in this entry.