



Full wwPDB X-ray Structure Validation Report ⓘ

May 14, 2020 – 11:56 am BST

PDB ID : 5XWX
Title : Crystal structure of the four N-terminal immunoglobulin domains of Sidekick-1 protein
Authors : Tang, H.; Dong, Y.; He, Y.
Deposited on : 2017-06-30
Resolution : 1.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1483 (1.56-1.56)
Clashscore	141614	1529 (1.56-1.56)
Ramachandran outliers	138981	1498 (1.56-1.56)
Sidechain outliers	138945	1495 (1.56-1.56)
RSRZ outliers	127900	1465 (1.56-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	458	 3% 74% 8% 18%
1	B	458	 4% 74% 8% 18%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 6499 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protein sidekick-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	376	2931	1854	509	554	14	0	0	0
1	B	376	2931	1854	509	554	14	0	0	0

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
2	A	338	338	338	0	0
2	B	299	299	299	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	63.85Å 118.06Å 122.84Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.52 – 1.55 29.52 – 1.55	Depositor EDS
% Data completeness (in resolution range)	97.1 (29.52-1.55) 97.1 (29.52-1.55)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.14 (at 1.55Å)	Xtrriage
Refinement program	PHENIX 1.9_1690	Depositor
R, R_{free}	0.222 , 0.250 0.227 , 0.252	Depositor DCC
R_{free} test set	1994 reflections (1.52%)	wwPDB-VP
Wilson B-factor (Å ²)	21.4	Xtrriage
Anisotropy	0.237	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 48.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.017 for -h,l,k	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6499	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.20% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.35	0/2999	0.55	0/4083
1	B	0.41	0/2999	0.57	1/4083 (0.0%)
All	All	0.38	0/5998	0.56	1/8166 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	401	LEU	C-N-CD	5.07	139.05	128.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2931	0	2909	43	0
1	B	2931	0	2909	32	0
2	A	338	0	0	12	1
2	B	299	0	0	6	0
All	All	6499	0	5818	74	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (74) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:291:HIS:CD2	1:A:298:ARG:HH12	1.13	1.61
1:A:291:HIS:CD2	1:A:298:ARG:NH1	1.76	1.44
1:A:291:HIS:CG	1:A:298:ARG:NH1	1.95	1.32
1:A:291:HIS:CG	1:A:298:ARG:HH11	1.52	1.26
1:B:392:LYS:O	1:B:394:GLN:OE1	1.56	1.20
1:B:181:CYS:SG	2:B:758:HOH:O	2.09	1.08
1:A:131:ARG:NH2	2:A:502:HOH:O	1.97	0.95
1:A:129:PHE:HB3	1:A:145:LYS:HD2	1.51	0.91
1:B:393:LEU:O	1:B:394:GLN:CD	2.11	0.89
1:A:291:HIS:HD2	1:A:298:ARG:HH12	1.22	0.86
1:B:381:LEU:HD12	1:B:404:GLY:HA2	1.62	0.82
1:B:143:GLN:NE2	2:B:501:HOH:O	2.03	0.82
1:B:393:LEU:C	1:B:394:GLN:OE1	2.18	0.81
1:A:356:GLU:OE2	2:A:503:HOH:O	1.98	0.80
1:A:176:LEU:HD23	2:A:672:HOH:O	1.81	0.80
1:A:181:CYS:SG	2:A:596:HOH:O	2.39	0.80
1:A:264:ASN:ND2	2:A:504:HOH:O	2.05	0.80
1:B:371:CYS:HB3	1:B:404:GLY:O	1.84	0.77
1:A:131:ARG:HG3	1:A:145:LYS:HD3	1.67	0.76
1:A:317:SER:OG	2:A:505:HOH:O	2.07	0.72
1:B:363:VAL:HG12	1:B:364:GLU:HG2	1.72	0.70
1:A:291:HIS:NE2	1:A:298:ARG:NH1	2.40	0.67
1:A:131:ARG:HE	1:A:145:LYS:HE2	1.60	0.66
1:A:356:GLU:HB3	2:A:503:HOH:O	1.97	0.64
1:B:353:ALA:HB3	1:B:372:ARG:HB2	1.80	0.63
1:B:75:LEU:H	1:B:75:LEU:HD12	1.65	0.62
1:B:392:LYS:C	1:B:394:GLN:OE1	2.39	0.61
1:A:125:LEU:N	1:A:125:LEU:HD12	2.16	0.61
1:B:381:LEU:HD12	1:B:404:GLY:CA	2.29	0.60
1:A:167:GLN:HE21	1:A:247:ASP:H	1.48	0.60
1:A:92:GLU:HG3	2:A:510:HOH:O	2.02	0.58
1:A:381:LEU:HD12	1:A:404:GLY:HA2	1.86	0.57
1:B:70:LYS:HE3	2:B:695:HOH:O	2.04	0.57
1:A:124:LYS:HB2	1:A:125:LEU:HD12	1.86	0.57
1:B:235:LYS:NZ	2:B:505:HOH:O	2.40	0.55
1:B:174:ASN:OD1	1:B:208:GLN:NE2	2.39	0.55
1:B:346:ILE:HD12	1:B:377:PRO:HD3	1.89	0.54
1:B:393:LEU:C	1:B:394:GLN:CD	2.63	0.54
1:B:104:ASN:ND2	2:B:507:HOH:O	2.42	0.53
1:A:247:ASP:HB3	1:A:251:HIS:CE1	2.43	0.53
1:A:104:ASN:ND2	2:A:501:HOH:O	1.95	0.53
1:A:294:ARG:HB3	1:A:299:LEU:HD21	1.91	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:371:CYS:HB2	1:B:383:TRP:CZ2	2.44	0.53
1:B:371:CYS:O	1:B:404:GLY:HA3	2.09	0.52
1:B:393:LEU:O	1:B:394:GLN:CG	2.59	0.51
1:A:304:HIS:HE1	1:A:311:THR:HB	1.76	0.51
1:B:161:GLN:CG	1:B:162:ARG:N	2.74	0.51
1:A:167:GLN:NE2	1:A:247:ASP:H	2.09	0.50
1:B:103:ARG:HD3	1:B:130:TYR:CZ	2.47	0.49
1:B:78:ILE:HD13	1:B:230:LYS:HG3	1.95	0.49
1:A:185:GLN:HG3	1:B:90:LEU:HD13	1.94	0.49
1:A:247:ASP:HB3	1:A:251:HIS:HE1	1.77	0.48
1:A:131:ARG:HD3	1:A:142:LEU:HG	1.96	0.47
1:B:64:ASP:HB3	1:B:94:SER:OG	2.15	0.46
1:A:389:PRO:HG2	1:A:392:LYS:HB2	1.97	0.46
1:A:131:ARG:HG3	1:A:145:LYS:CD	2.40	0.46
1:A:291:HIS:CE1	1:A:298:ARG:NH1	2.84	0.46
1:A:81:GLU:HG2	2:A:506:HOH:O	2.15	0.46
1:B:144:ARG:HH21	1:B:431:GLN:HE21	1.65	0.45
1:B:394:GLN:HG2	1:B:394:GLN:O	2.17	0.44
1:A:125:LEU:N	1:A:125:LEU:CD1	2.79	0.44
1:B:401:LEU:HB3	1:B:402:PRO:CD	2.48	0.44
1:A:401:LEU:HA	1:A:402:PRO:HD2	1.81	0.44
1:B:294:ARG:HB3	1:B:299:LEU:HD21	2.00	0.43
1:A:312:ILE:HD13	1:A:343:LEU:HD22	1.99	0.43
1:A:350:TYR:CZ	1:A:374:MET:HB3	2.55	0.42
1:A:291:HIS:CB	1:A:298:ARG:NH1	2.74	0.41
1:A:81:GLU:OE2	2:A:506:HOH:O	2.22	0.41
1:A:420:GLN:HG2	1:A:422:PHE:CE2	2.56	0.41
1:B:149:GLN:CD	2:B:502:HOH:O	2.58	0.41
1:A:81:GLU:HG2	2:A:518:HOH:O	2.21	0.41
1:A:68:TYR:CZ	1:A:92:GLU:HB3	2.55	0.41
1:A:124:LYS:CB	1:A:125:LEU:HD12	2.52	0.40
1:B:394:GLN:N	1:B:394:GLN:OE1	2.53	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:514:HOH:O	2:A:678:HOH:O[2_455]	2.13	0.07

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	374/458 (82%)	365 (98%)	9 (2%)	0	100	100
1	B	374/458 (82%)	366 (98%)	8 (2%)	0	100	100
All	All	748/916 (82%)	731 (98%)	17 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	322/378 (85%)	321 (100%)	1 (0%)	92	85
1	B	322/378 (85%)	321 (100%)	1 (0%)	92	85
All	All	644/756 (85%)	642 (100%)	2 (0%)	92	85

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	362	GLU
1	B	176	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	291	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	304	HIS
1	B	409	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	376/458 (82%)	0.14	15 (3%) 38 44	18, 27, 46, 64	0
1	B	376/458 (82%)	0.22	20 (5%) 26 31	19, 29, 46, 64	0
All	All	752/916 (82%)	0.18	35 (4%) 31 36	18, 28, 46, 64	0

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	250	THR	8.5
1	B	157	MET	7.3
1	B	404	GLY	6.1
1	A	363	VAL	4.4
1	A	158	ASP	4.4
1	B	95	TRP	4.1
1	B	371	CYS	4.0
1	A	157	MET	4.0
1	A	439	THR	3.8
1	B	333	THR	3.7
1	B	158	ASP	3.2
1	A	75	LEU	3.2
1	A	402	PRO	3.1
1	A	403	SER	3.1
1	B	177	PRO	3.0
1	B	179	VAL	3.0
1	B	331	GLY	3.0
1	B	403	SER	2.9
1	A	394	GLN	2.9
1	B	439	THR	2.9
1	A	331	GLY	2.8
1	B	394	GLN	2.7
1	A	105	ASP	2.7
1	B	363	VAL	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	105	ASP	2.6
1	A	271	SER	2.5
1	A	95	TRP	2.5
1	B	372	ARG	2.5
1	B	373	ALA	2.4
1	B	176	LEU	2.4
1	A	159	THR	2.4
1	B	438	VAL	2.3
1	A	251	HIS	2.2
1	B	362	GLU	2.1
1	B	159	THR	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.