



# wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 23, 2023 – 12:45 AM JST

PDB ID : 7XZN  
Title : Formate-tetrahydrofolate ligase from *Peptostreptococcus anaerobius*  
Authors : Fang, C.L.; Zhang, Y.  
Deposited on : 2022-06-03  
Resolution : 2.04 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.04 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1692 (2.04-2.04)
Clashscore	141614	1773 (2.04-2.04)
Ramachandran outliers	138981	1752 (2.04-2.04)
Sidechain outliers	138945	1752 (2.04-2.04)
RSRZ outliers	127900	1672 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	562	 10% 92% 7%
1	B	562	 8% 93% 6%
1	C	562	 8% 90% 9%
1	D	562	 10% 93% 5%

## 2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 16831 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Formate--tetrahydrofolate ligase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	555	4159	2617	717	804	21	0	0	0
1	B	556	4162	2619	715	807	21	0	0	0
1	C	556	4172	2625	719	807	21	0	0	0
1	D	555	4137	2600	714	802	21	0	0	0

There are 16 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	GLY	-	expression tag	UNP A0A379CIH2
A	-2	SER	-	expression tag	UNP A0A379CIH2
A	-1	HIS	-	expression tag	UNP A0A379CIH2
A	0	MET	-	expression tag	UNP A0A379CIH2
B	-3	GLY	-	expression tag	UNP A0A379CIH2
B	-2	SER	-	expression tag	UNP A0A379CIH2
B	-1	HIS	-	expression tag	UNP A0A379CIH2
B	0	MET	-	expression tag	UNP A0A379CIH2
C	-3	GLY	-	expression tag	UNP A0A379CIH2
C	-2	SER	-	expression tag	UNP A0A379CIH2
C	-1	HIS	-	expression tag	UNP A0A379CIH2
C	0	MET	-	expression tag	UNP A0A379CIH2
D	-3	GLY	-	expression tag	UNP A0A379CIH2
D	-2	SER	-	expression tag	UNP A0A379CIH2
D	-1	HIS	-	expression tag	UNP A0A379CIH2
D	0	MET	-	expression tag	UNP A0A379CIH2

- Molecule 2 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0
2	C	1	Total C O 4 2 2	0	0
2	C	1	Total C O 4 2 2	0	0
2	C	1	Total C O 4 2 2	0	0
2	D	1	Total C O 4 2 2	0	0
2	D	1	Total C O 4 2 2	0	0

- Molecule 3 is TRIS(HYDROXYETHYL)AMINOMETHANE (three-letter code: TAM) (formula: C<sub>7</sub>H<sub>17</sub>NO<sub>3</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	11	7	1	3	0	0
3	B	1	11	7	1	3	0	0
3	C	1	11	7	1	3	0	0
3	D	1	11	7	1	3	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	28	28	28	0	0
4	B	36	36	36	0	0
4	C	28	28	28	0	0
4	D	25	25	25	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Formate--tetrahydrofolate ligase




- Molecule 1: Formate--tetrahydrofolate ligase





- Molecule 1: Formate--tetrahydrofolate ligase





● Molecule 1: Formate--tetrahydrofolate ligase



## 4 Data and refinement statistics i

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	215.87Å 116.30Å 107.38Å 90.00° 92.58° 90.00°	Depositor
Resolution (Å)	30.50 – 2.04 54.02 – 2.04	Depositor EDS
% Data completeness (in resolution range)	99.8 (30.50-2.04) 99.9 (54.02-2.04)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.58 (at 2.05Å)	Xtrriage
Refinement program	PHENIX 1.14_3260	Depositor
R, $R_{free}$	0.272 , 0.283 0.285 , 0.296	Depositor DCC
$R_{free}$ test set	2012 reflections (1.20%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	36.0	Xtrriage
Anisotropy	0.408	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 50.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.55$ , $\langle L^2 \rangle = 0.40$	Xtrriage
Estimated twinning fraction	0.016 for -h,-k,l	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	16831	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	54.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 54.13 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.8167e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, TAM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.24	0/4218	0.44	0/5711
1	B	0.24	0/4221	0.43	0/5716
1	C	0.24	0/4232	0.44	0/5731
1	D	0.24	0/4195	0.43	0/5685
All	All	0.24	0/16866	0.43	0/22843

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4159	0	4195	23	0
1	B	4162	0	4190	21	0
1	C	4172	0	4201	29	0
1	D	4137	0	4155	20	0
2	A	16	0	24	2	0
2	B	4	0	6	2	0
2	C	12	0	18	4	0
2	D	8	0	12	1	0
3	A	11	0	17	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	11	0	17	3	0
3	C	11	0	17	3	0
3	D	11	0	17	1	0
4	A	28	0	0	0	0
4	B	36	0	0	1	0
4	C	28	0	0	0	0
4	D	25	0	0	0	0
All	All	16831	0	16869	91	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 91 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:105:ILE:HG22	1:A:106:LYS:H	1.46	0.79
1:D:11:GLN:O	3:D:603:TAM:N	2.25	0.69
1:A:173:ARG:HH21	2:A:603:EDO:H21	1.59	0.68
1:D:333:ARG:NH1	1:D:347:ASN:OD1	2.27	0.67
1:C:333:ARG:NH1	1:C:347:ASN:OD1	2.32	0.62

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	553/562 (98%)	532 (96%)	21 (4%)	0	<a href="#">100</a> <a href="#">100</a>
1	B	554/562 (99%)	535 (97%)	19 (3%)	0	<a href="#">100</a> <a href="#">100</a>
1	C	554/562 (99%)	532 (96%)	21 (4%)	1 (0%)	47 39
1	D	553/562 (98%)	531 (96%)	22 (4%)	0	<a href="#">100</a> <a href="#">100</a>

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	2214/2248 (98%)	2130 (96%)	83 (4%)	1 (0%)	100 100

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	507	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	434/447 (97%)	431 (99%)	3 (1%)	84 84
1	B	434/447 (97%)	432 (100%)	2 (0%)	88 89
1	C	436/447 (98%)	432 (99%)	4 (1%)	78 79
1	D	430/447 (96%)	427 (99%)	3 (1%)	84 84
All	All	1734/1788 (97%)	1722 (99%)	12 (1%)	84 84

5 of 12 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	127	PHE
1	C	431	GLU
1	D	127	PHE
1	D	44	TYR
1	B	44	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

14 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	TAM	A	605	-	7,10,10	0.51	0	9,12,12	5.43	6 (66%)
3	TAM	B	602	-	7,10,10	0.46	0	9,12,12	1.06	1 (11%)
2	EDO	A	604	-	3,3,3	0.45	0	2,2,2	0.39	0
2	EDO	D	601	-	3,3,3	0.45	0	2,2,2	0.35	0
3	TAM	D	603	-	7,10,10	0.53	0	9,12,12	0.58	0
2	EDO	C	602	-	3,3,3	0.45	0	2,2,2	0.30	0
2	EDO	A	601	-	3,3,3	0.45	0	2,2,2	0.26	0
2	EDO	A	603	-	3,3,3	0.46	0	2,2,2	0.30	0
2	EDO	B	601	-	3,3,3	0.46	0	2,2,2	0.28	0
2	EDO	C	603	-	3,3,3	0.42	0	2,2,2	0.44	0
2	EDO	D	602	-	3,3,3	0.46	0	2,2,2	0.31	0
2	EDO	A	602	-	3,3,3	0.46	0	2,2,2	0.33	0
2	EDO	C	601	-	3,3,3	0.46	0	2,2,2	0.34	0
3	TAM	C	604	-	7,10,10	0.45	0	9,12,12	0.70	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	TAM	A	605	-	-	7/12/12/12	-
3	TAM	B	602	-	-	11/12/12/12	-
2	EDO	A	604	-	-	0/1/1/1	-
2	EDO	D	601	-	-	0/1/1/1	-
3	TAM	D	603	-	-	7/12/12/12	-
2	EDO	C	602	-	-	1/1/1/1	-
2	EDO	A	601	-	-	1/1/1/1	-
2	EDO	A	603	-	-	0/1/1/1	-
2	EDO	B	601	-	-	0/1/1/1	-
2	EDO	C	603	-	-	0/1/1/1	-
2	EDO	D	602	-	-	1/1/1/1	-
2	EDO	A	602	-	-	0/1/1/1	-
2	EDO	C	601	-	-	0/1/1/1	-
3	TAM	C	604	-	-	6/12/12/12	-

There are no bond length outliers.

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	605	TAM	C1-C-N	-8.95	82.58	108.09
3	A	605	TAM	C2-C-N	-8.05	85.16	108.09
3	A	605	TAM	C3-C-N	-6.39	89.89	108.09
3	A	605	TAM	C2-C-C1	5.73	120.62	110.50
3	A	605	TAM	C3-C-C1	4.95	119.23	110.50

There are no chirality outliers.

5 of 34 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	605	TAM	C3-C-C1-C4
3	A	605	TAM	N-C-C1-C4
3	A	605	TAM	C1-C-C2-C5
3	A	605	TAM	N-C-C2-C5
3	A	605	TAM	C1-C-C3-C6

There are no ring outliers.

10 monomers are involved in 16 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	602	TAM	3	0
2	D	601	EDO	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	603	TAM	1	0
2	C	602	EDO	1	0
2	A	601	EDO	1	0
2	A	603	EDO	1	0
2	B	601	EDO	2	0
2	C	603	EDO	2	0
2	C	601	EDO	1	0
3	C	604	TAM	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	555/562 (98%)	0.89	56 (10%) <b>7</b> <b>7</b>	31, 54, 88, 102	0
1	B	556/562 (98%)	0.84	44 (7%) <b>12</b> <b>13</b>	32, 52, 81, 96	0
1	C	556/562 (98%)	0.80	46 (8%) <b>11</b> <b>11</b>	31, 50, 76, 104	0
1	D	555/562 (98%)	0.82	57 (10%) <b>6</b> <b>6</b>	31, 50, 78, 106	0
All	All	2222/2248 (98%)	0.84	203 (9%) <b>9</b> <b>9</b>	31, 52, 82, 106	0

The worst 5 of 203 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	68	ALA	10.9
1	A	430	GLU	7.1
1	B	394	VAL	5.7
1	A	68	ALA	5.6
1	A	433	GLU	5.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	TAM	B	602	11/11	0.38	0.33	61,70,76,77	0
3	TAM	D	603	11/11	0.54	0.35	69,74,81,81	0
2	EDO	A	604	4/4	0.55	0.35	56,60,61,63	0
3	TAM	C	604	11/11	0.63	0.30	57,71,75,81	0
2	EDO	A	601	4/4	0.67	0.19	50,52,55,57	0
2	EDO	C	603	4/4	0.68	0.22	55,56,59,60	0
2	EDO	D	602	4/4	0.71	0.22	43,57,60,61	0
2	EDO	C	602	4/4	0.72	0.28	33,42,51,53	0
2	EDO	A	602	4/4	0.74	0.21	47,51,51,52	0
2	EDO	C	601	4/4	0.76	0.26	33,38,45,54	0
2	EDO	B	601	4/4	0.77	0.23	35,48,52,58	0
3	TAM	A	605	11/11	0.78	0.21	75,78,82,83	0
2	EDO	D	601	4/4	0.86	0.25	39,40,50,55	0
2	EDO	A	603	4/4	0.89	0.20	35,47,51,56	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.