



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 15, 2020 – 01:11 pm BST

PDB ID : 2YHE  
Title : Structure determination of the stereoselective inverting sec-alkylsulfatase Pisa1 from *Pseudomonas* sp.  
Authors : Kepplinger, B.; Faber, K.; Macheroux, P.; Schober, M.; Knaus, T.; Wagner, U.G.  
Deposited on : 2011-04-29  
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	668	 81% 13% • •
1	B	668	 80% 14% • 5%
1	C	668	 77% 16% • 5%
1	D	668	 78% 15% • 5%
1	E	668	 77% 16% • 5%
1	F	668	 79% 15% • 5%

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 31033 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SEC-ALKYL SULFATASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	639	5044	3196	888	941	19	0	10	0
1	B	634	4959	3142	863	935	19	0	4	0
1	C	634	4948	3133	863	933	19	0	2	0
1	D	634	4961	3145	867	930	19	0	4	0
1	E	634	4959	3144	863	933	19	0	5	0
1	F	634	4954	3137	865	933	19	0	3	0

There are 54 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	107	ARG	HIS	conflict	UNP F8KAY7
A	661	LEU	-	expression tag	UNP F8KAY7
A	662	GLU	-	expression tag	UNP F8KAY7
A	663	HIS	-	expression tag	UNP F8KAY7
A	664	HIS	-	expression tag	UNP F8KAY7
A	665	HIS	-	expression tag	UNP F8KAY7
A	666	HIS	-	expression tag	UNP F8KAY7
A	667	HIS	-	expression tag	UNP F8KAY7
A	668	HIS	-	expression tag	UNP F8KAY7
B	107	ARG	HIS	conflict	UNP F8KAY7
B	661	LEU	-	expression tag	UNP F8KAY7
B	662	GLU	-	expression tag	UNP F8KAY7
B	663	HIS	-	expression tag	UNP F8KAY7
B	664	HIS	-	expression tag	UNP F8KAY7
B	665	HIS	-	expression tag	UNP F8KAY7
B	666	HIS	-	expression tag	UNP F8KAY7
B	667	HIS	-	expression tag	UNP F8KAY7

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	668	HIS	-	expression tag	UNP F8KAY7
C	107	ARG	HIS	conflict	UNP F8KAY7
C	661	LEU	-	expression tag	UNP F8KAY7
C	662	GLU	-	expression tag	UNP F8KAY7
C	663	HIS	-	expression tag	UNP F8KAY7
C	664	HIS	-	expression tag	UNP F8KAY7
C	665	HIS	-	expression tag	UNP F8KAY7
C	666	HIS	-	expression tag	UNP F8KAY7
C	667	HIS	-	expression tag	UNP F8KAY7
C	668	HIS	-	expression tag	UNP F8KAY7
D	107	ARG	HIS	conflict	UNP F8KAY7
D	661	LEU	-	expression tag	UNP F8KAY7
D	662	GLU	-	expression tag	UNP F8KAY7
D	663	HIS	-	expression tag	UNP F8KAY7
D	664	HIS	-	expression tag	UNP F8KAY7
D	665	HIS	-	expression tag	UNP F8KAY7
D	666	HIS	-	expression tag	UNP F8KAY7
D	667	HIS	-	expression tag	UNP F8KAY7
D	668	HIS	-	expression tag	UNP F8KAY7
E	107	ARG	HIS	conflict	UNP F8KAY7
E	661	LEU	-	expression tag	UNP F8KAY7
E	662	GLU	-	expression tag	UNP F8KAY7
E	663	HIS	-	expression tag	UNP F8KAY7
E	664	HIS	-	expression tag	UNP F8KAY7
E	665	HIS	-	expression tag	UNP F8KAY7
E	666	HIS	-	expression tag	UNP F8KAY7
E	667	HIS	-	expression tag	UNP F8KAY7
E	668	HIS	-	expression tag	UNP F8KAY7
F	107	ARG	HIS	conflict	UNP F8KAY7
F	661	LEU	-	expression tag	UNP F8KAY7
F	662	GLU	-	expression tag	UNP F8KAY7
F	663	HIS	-	expression tag	UNP F8KAY7
F	664	HIS	-	expression tag	UNP F8KAY7
F	665	HIS	-	expression tag	UNP F8KAY7
F	666	HIS	-	expression tag	UNP F8KAY7
F	667	HIS	-	expression tag	UNP F8KAY7
F	668	HIS	-	expression tag	UNP F8KAY7

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).


*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
-----	-------	----------	-------	---------	---------

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	D	2	Total Zn 2 2	0	0
2	E	2	Total Zn 2 2	0	0
2	B	2	Total Zn 2 2	0	0
2	C	2	Total Zn 2 2	0	0
2	A	2	Total Zn 2 2	0	0
2	F	2	Total Zn 2 2	0	0

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0
3	C	1	Total O S 5 4 1	0	0
3	D	1	Total O S 5 4 1	0	0

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	E	1	Total	O	S	0	0
			5	4	1		
3	F	1	Total	O	S	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	257	Total	O	0	0
			257	257		
4	B	225	Total	O	0	0
			225	225		
4	C	181	Total	O	0	0
			181	181		
4	D	178	Total	O	0	0
			178	178		
4	E	166	Total	O	0	0
			166	166		
4	F	159	Total	O	0	0
			159	159		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: SEC-ALKYL SULFATASE




#### • Molecule 1: SEC-ALKYL SULFATASE





#### • Molecule 1: SEC-ALKYL SULFATASE







• Molecule 1: SEC-ALKYL SULFATASE





• Molecule 1: SEC-ALKYL SULFATASE







● Molecule 1: SEC-ALKYL SULFATASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	74.73Å 200.59Å 245.33Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.91 – 2.70 19.90 – 2.70	Depositor EDS
% Data completeness (in resolution range)	100.0 (19.91-2.70) 100.0 (19.90-2.70)	Depositor EDS
$R_{merge}$	0.17	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.47 (at 2.71Å)	Xtrriage
Refinement program	REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.183 , 0.230 0.178 , 0.225	Depositor DCC
$R_{free}$ test set	5090 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	41.1	Xtrriage
Anisotropy	0.046	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.26 , 48.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	31033	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	47.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.97% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.50	0/5184	0.73	2/7031 (0.0%)
1	B	0.50	0/5076	0.72	2/6885 (0.0%)
1	C	0.50	0/5059	0.73	2/6862 (0.0%)
1	D	0.51	0/5078	0.72	1/6887 (0.0%)
1	E	0.51	0/5079	0.72	0/6891
1	F	0.50	0/5068	0.72	1/6874 (0.0%)
All	All	0.50	0/30544	0.72	8/41430 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1
1	D	0	1
All	All	0	2

There are no bond length outliers.

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	539	GLY	C-N-CA	10.21	147.24	121.70
1	D	662	GLU	C-N-CA	8.46	142.84	121.70
1	A	664	HIS	C-N-CA	7.62	140.75	121.70
1	C	396	ILE	CA-CB-CG2	5.41	121.73	110.90
1	C	225	MET	CG-SD-CE	-5.32	91.69	100.20

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	353	GLN	Sidechain
1	D	662	GLU	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5044	0	5016	57	0
1	B	4959	0	4918	52	0
1	C	4948	0	4901	55	0
1	D	4961	0	4934	67	0
1	E	4959	0	4923	78	0
1	F	4954	0	4911	62	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
2	C	2	0	0	0	0
2	D	2	0	0	0	0
2	E	2	0	0	0	0
2	F	2	0	0	0	0
3	A	5	0	0	0	0
3	B	5	0	0	0	0
3	C	5	0	0	0	0
3	D	5	0	0	0	0
3	E	5	0	0	0	0
3	F	5	0	0	0	0
4	A	257	0	0	10	0
4	B	225	0	0	6	0
4	C	181	0	0	11	0
4	D	178	0	0	7	0
4	E	166	0	0	8	0
4	F	159	0	0	3	0
All	All	31033	0	29603	355	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 355 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:40:ILE:CD1	1:E:40:ILE:CG1	1.74	1.58
1:E:552:PHE:CE1	1:E:618:ALA:HB2	1.77	1.19
1:D:661:LEU:HD23	1:D:661:LEU:H	1.00	1.15
1:D:67:LEU:HD12	4:D:2011:HOH:O	1.46	1.15
1:E:661:LEU:H	1:E:661:LEU:HD23	0.98	1.13

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	647/668 (97%)	626 (97%)	18 (3%)	3 (0%)	29	54
1	B	636/668 (95%)	615 (97%)	19 (3%)	2 (0%)	41	66
1	C	634/668 (95%)	613 (97%)	19 (3%)	2 (0%)	41	66
1	D	636/668 (95%)	614 (96%)	20 (3%)	2 (0%)	41	66
1	E	637/668 (95%)	618 (97%)	17 (3%)	2 (0%)	41	66
1	F	635/668 (95%)	613 (96%)	18 (3%)	4 (1%)	25	50
All	All	3825/4008 (95%)	3699 (97%)	111 (3%)	15 (0%)	34	60

5 of 15 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	57	ARG
1	A	665	HIS
1	B	57	ARG
1	C	57	ARG
1	D	56	ASP

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	527/539 (98%)	485 (92%)	42 (8%)	12	27
1	B	516/539 (96%)	476 (92%)	40 (8%)	12	29
1	C	514/539 (95%)	464 (90%)	50 (10%)	8	19
1	D	516/539 (96%)	472 (92%)	44 (8%)	10	24
1	E	516/539 (96%)	470 (91%)	46 (9%)	9	22
1	F	515/539 (96%)	471 (92%)	44 (8%)	10	24
All	All	3104/3234 (96%)	2838 (91%)	266 (9%)	10	24

5 of 266 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	609	LEU
1	D	327	VAL
1	F	399[B]	GLU
1	C	616	GLN
1	D	90[A]	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	213	HIS
1	F	311	ASN
1	E	616	GLN
1	B	259	ASN
1	E	311	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 18 ligands modelled in this entry, 12 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	SO4	C	801	-	4,4,4	0.38	0	6,6,6	0.35	0
3	SO4	B	801	-	4,4,4	0.28	0	6,6,6	0.45	0
3	SO4	D	801	-	4,4,4	0.74	0	6,6,6	0.33	0
3	SO4	F	801	-	4,4,4	0.35	0	6,6,6	0.29	0
3	SO4	A	801	-	4,4,4	0.42	0	6,6,6	0.21	0
3	SO4	E	801	-	4,4,4	0.54	0	6,6,6	0.28	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	639/668 (95%)	-0.60	5 (0%) 86 87	21, 41, 78, 135	0
1	B	634/668 (94%)	-0.51	13 (2%) 63 65	25, 42, 86, 133	0
1	C	634/668 (94%)	-0.28	11 (1%) 70 72	25, 49, 85, 120	0
1	D	634/668 (94%)	-0.50	10 (1%) 72 74	25, 42, 76, 127	0
1	E	634/668 (94%)	-0.44	18 (2%) 53 54	24, 42, 80, 135	0
1	F	634/668 (94%)	-0.50	10 (1%) 72 74	22, 44, 76, 134	0
All	All	3809/4008 (95%)	-0.47	67 (1%) 68 70	21, 43, 82, 135	0

The worst 5 of 67 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	54	PHE	6.7
1	F	55	ALA	5.5
1	B	54	PHE	5.4
1	A	54	PHE	5.1
1	C	54	PHE	4.9

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	ZN	C	700	1/1	0.98	0.03	46,46,46,46	0
2	ZN	B	701	1/1	0.99	0.07	40,40,40,40	0
3	SO4	B	801	5/5	0.99	0.07	36,38,41,42	0
3	SO4	C	801	5/5	0.99	0.11	45,49,50,52	0
3	SO4	E	801	5/5	0.99	0.09	30,34,35,36	0
2	ZN	D	701	1/1	0.99	0.07	34,34,34,34	0
2	ZN	B	700	1/1	0.99	0.05	44,44,44,44	0
2	ZN	A	700	1/1	0.99	0.04	39,39,39,39	0
2	ZN	F	700	1/1	0.99	0.03	43,43,43,43	0
2	ZN	A	701	1/1	0.99	0.07	38,38,38,38	0
2	ZN	E	700	1/1	0.99	0.03	43,43,43,43	0
2	ZN	C	701	1/1	0.99	0.06	45,45,45,45	0
2	ZN	E	701	1/1	0.99	0.06	37,37,37,37	0
3	SO4	D	801	5/5	0.99	0.08	28,31,34,37	0
2	ZN	D	700	1/1	1.00	0.05	45,45,45,45	0
2	ZN	F	701	1/1	1.00	0.09	46,46,46,46	0
3	SO4	F	801	5/5	1.00	0.06	30,32,34,35	0
3	SO4	A	801	5/5	1.00	0.10	34,35,39,40	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.