



# Full wwPDB X-ray Structure Validation Report i

Dec 21, 2020 – 11:03 am GMT

PDB ID : 6YHL  
Title : Crystal structure of CNFy from Yersinia pseudotuberculosis - N-terminal fragment comprising residues 1-704  
Authors : Lukat, P.; Gazdag, E.M.; Heidler, T.V.; Blankenfeldt, W.  
Deposited on : 2020-03-30  
Resolution : 3.28 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the i symbol.

---

The following versions of software and data (see references ①) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.16  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.16


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## X-RAY DIFFRACTION


The reported resolution of this entry is 3.28 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1177 (3.32-3.24)
Clashscore	141614	1044 (3.30-3.26)
Ramachandran outliers	138981	1026 (3.30-3.26)
Sidechain outliers	138945	1025 (3.30-3.26)
RSRZ outliers	127900	1141 (3.32-3.24)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



## 2 Entry composition [\(i\)](#)

There is only 1 type of molecule in this entry. The entry contains 21444 atoms, of which 10422 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Cytotoxic necrotizing factor.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	695	10675	3524	5186	874	1081	10	0	0	0
1	B	698	10769	3546	5236	885	1091	11	0	0	0


### 3 Residue-property plots [\(i\)](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Cytotoxic necrotizing factor



- Molecule 1: Cytotoxic necrotizing factor



## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	94.42Å 99.90Å 131.36Å 90.00° 95.86° 90.00°	Depositor
Resolution (Å)	54.68 – 3.28 54.68 – 3.28	Depositor EDS
% Data completeness (in resolution range)	99.9 (54.68-3.28) 100.0 (54.68-3.28)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) >$ <sup>1</sup>	1.88 (at 3.26Å)	Xtriage
Refinement program	PHENIX 1.13_2998	Depositor
$R$ , $R_{free}$	0.240 , 0.265 0.242 , 0.267	Depositor DCC
$R_{free}$ test set	1889 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	98.3	Xtriage
Anisotropy	0.240	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 47.0	EDS
L-test for twinning <sup>2</sup>	$<  L  > = 0.49$ , $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	21444	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	103.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.23% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $< |L| >$ ,  $< L^2 >$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [\(i\)](#)

### 5.1 Standard geometry [\(i\)](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.26	0/5620	0.44	0/7655
1	B	0.28	0/5665	0.45	1/7715 (0.0%)
All	All	0.27	0/11285	0.45	1/15370 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	B	351	VAL	CG1-CB-CG2	5.01	118.92	110.90

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5489	5186	5185	85	0
1	B	5533	5236	5235	65	0
All	All	11022	10422	10420	143	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (143) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:456:ILE:HD11	1:A:467:LEU:HD11	1.48	0.95
1:A:63:LYS:NZ	1:B:473:GLU:OE2	2.14	0.80
1:B:115:ASP:OD1	1:B:117:THR:HG22	1.81	0.79
1:B:623:GLU:OE1	1:B:626:PHE:N	2.16	0.78
1:A:115:ASP:OD1	1:A:117:THR:HG22	1.84	0.78
1:B:451:ASN:ND2	1:B:451:ASN:O	2.17	0.77
1:B:153:GLU:OE2	1:B:504:ARG:NH1	2.17	0.77
1:A:153:GLU:OE2	1:A:504:ARG:NH1	2.20	0.75
1:A:333:TYR:O	1:A:336:ASN:ND2	2.20	0.74
1:A:445:GLU:N	1:A:445:GLU:OE1	2.21	0.73
1:A:15:GLU:O	1:A:19:ASN:ND2	2.22	0.73
1:B:445:GLU:OE1	1:B:445:GLU:N	2.22	0.72
1:A:456:ILE:HD11	1:A:467:LEU:CD1	2.18	0.72
1:B:456:ILE:HD11	1:B:467:LEU:HD11	1.71	0.72
1:B:623:GLU:OE1	1:B:625:ASN:N	2.23	0.71
1:A:214:MET:HB3	1:A:219:THR:HG23	1.78	0.65
1:B:6:GLN:HB3	1:B:348:PHE:HE1	1.60	0.65
1:A:77:GLU:OE2	1:A:92:ARG:NH2	2.29	0.65
1:A:56:THR:CG2	1:A:76:ILE:HD11	2.26	0.65
1:B:38:THR:O	1:B:39:LEU:HD12	1.97	0.64
1:B:622:LEU:HD22	1:B:683:ALA:HA	1.80	0.64
1:A:292:MET:O	1:A:295:ASN:ND2	2.30	0.64
1:A:137:SER:OG	1:A:320:ASP:OD1	2.16	0.63
1:B:190:ASN:ND2	1:B:195:ILE:O	2.31	0.63
1:A:614:THR:HG21	1:A:618:SER:O	1.99	0.63
1:A:666:ILE:HG22	1:A:679:ILE:HD11	1.81	0.61
1:A:346:PRO:HG2	1:A:350:ILE:HD11	1.83	0.61
1:B:456:ILE:HD11	1:B:467:LEU:CD1	2.30	0.61
1:A:227:THR:HG23	1:A:366:SER:O	2.01	0.61
1:A:349:ASP:O	1:A:350:ILE:HD12	2.01	0.60
1:B:294:SER:OG	1:B:635:LEU:O	2.17	0.60
1:A:56:THR:HG23	1:A:76:ILE:HD11	1.85	0.59
1:A:20:PHE:CG	1:A:338:LEU:HD21	2.38	0.58
1:B:77:GLU:OE2	1:B:92:ARG:NH2	2.35	0.58
1:B:651:SER:OG	1:B:693:ARG:NH1	2.34	0.57
1:B:71:LEU:HD23	1:B:112:ASP:O	2.06	0.56
1:B:491:GLU:OE2	1:B:518:ARG:NH2	2.38	0.56
1:A:287:ASP:O	1:A:288:THR:OG1	2.16	0.56
1:A:666:ILE:HG22	1:A:679:ILE:CD1	2.36	0.56
1:B:438:LEU:HD12	1:B:439:GLU:N	2.21	0.56
1:B:188:ILE:O	1:B:197:ARG:NH2	2.40	0.55
1:A:350:ILE:O	1:A:351:VAL:HG22	2.06	0.55

Continued on next page...

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:94:ASP:HB2	1:A:95:ILE:HD12	1.89	0.54
1:A:622:LEU:HG	1:A:623:GLU:HG2	1.89	0.54
1:A:535:ILE:HD11	1:A:610:ALA:CB	2.38	0.53
1:A:651:SER:OG	1:A:693:ARG:NH1	2.36	0.53
1:A:98:GLU:O	1:A:101:SER:OG	2.25	0.53
1:B:614:THR:HG21	1:B:618:SER:O	2.09	0.53
1:B:2:LYS:HE2	1:B:495:GLU:CB	2.38	0.53
1:B:434:LEU:HD21	1:B:477:ILE:HD12	1.90	0.53
1:A:349:ASP:C	1:A:350:ILE:HD12	2.29	0.53
1:A:540:ILE:HD13	1:A:692:ASP:HB3	1.92	0.52
1:B:451:ASN:OD1	1:B:514:ILE:N	2.34	0.51
1:B:280:GLN:HB3	1:B:286:VAL:HG23	1.92	0.51
1:A:543:PRO:HD2	1:A:546:GLU:HG3	1.91	0.51
1:B:668:ASN:HB2	1:B:677:TRP:CE3	2.46	0.50
1:B:127:TYR:CE2	1:B:131:LEU:HD11	2.47	0.50
1:A:190:ASN:ND2	1:A:195:ILE:O	2.44	0.50
1:A:216:ASP:N	1:A:216:ASP:OD1	2.45	0.50
1:A:399:GLY:HA2	1:A:403:ALA:HB3	1.94	0.49
1:A:429:LEU:HD11	1:B:59:ARG:HH22	1.76	0.49
1:A:197:ARG:HB3	1:A:298:ILE:HD12	1.94	0.49
1:A:540:ILE:CD1	1:A:692:ASP:HB3	2.43	0.49
1:A:668:ASN:HB2	1:A:677:TRP:CE3	2.47	0.49
1:B:603:GLU:HA	1:B:655:PHE:CD1	2.48	0.49
1:B:399:GLY:HA2	1:B:403:ALA:HB3	1.93	0.49
1:B:622:LEU:HG	1:B:623:GLU:CG	2.43	0.49
1:A:227:THR:N	1:A:366:SER:O	2.41	0.49
1:A:56:THR:HG23	1:A:76:ILE:CD1	2.43	0.49
1:B:216:ASP:OD1	1:B:216:ASP:N	2.46	0.49
1:A:209:ASP:HB2	1:A:224:PRO:HG2	1.96	0.48
1:B:568:ASP:OD1	1:B:569:SER:N	2.46	0.48
1:B:211:PHE:O	1:B:221:LEU:HD12	2.14	0.48
1:A:196:LEU:HD13	1:A:198:TYR:OH	2.14	0.47
1:B:456:ILE:CD1	1:B:467:LEU:HD11	2.43	0.47
1:A:258:LYS:O	1:A:314:ARG:NH2	2.39	0.47
1:A:281:ILE:HG13	1:A:286:VAL:HG21	1.95	0.47
1:A:223:ILE:O	1:A:223:ILE:HG23	2.15	0.47
1:B:4:GLN:HG3	1:B:496:THR:HG21	1.96	0.47
1:B:451:ASN:OD1	1:B:513:LYS:HA	2.15	0.46
1:A:211:PHE:O	1:A:221:LEU:HD12	2.16	0.46
1:A:603:GLU:HA	1:A:655:PHE:CD1	2.51	0.46
1:B:223:ILE:O	1:B:223:ILE:HG23	2.15	0.46

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:56:THR:HG22	1:A:76:ILE:HD11	1.97	0.46
1:B:544:TYR:CD1	1:B:553:LEU:HD12	2.51	0.46
1:A:451:ASN:OD1	1:A:451:ASN:N	2.49	0.46
1:A:203:TYR:HD2	1:A:277:MET:HE3	1.80	0.46
1:A:349:ASP:O	1:A:350:ILE:HB	2.15	0.46
1:A:608:ASN:ND2	1:A:617:GLY:O	2.49	0.46
1:A:543:PRO:HD3	1:A:695:PHE:HD1	1.81	0.46
1:A:137:SER:N	1:A:323:ILE:HG21	2.31	0.45
1:A:300:GLU:OE1	1:A:602:THR:HG21	2.16	0.45
1:A:535:ILE:HD11	1:A:610:ALA:HB1	1.98	0.45
1:B:376:ILE:O	1:B:385:ARG:NE	2.49	0.45
1:B:612:TYR:O	1:B:614:THR:HG23	2.16	0.45
1:B:37:THR:O	1:B:38:THR:OG1	2.33	0.45
1:B:623:GLU:HG2	1:B:626:PHE:HB2	1.97	0.45
1:B:436:ASP:OD1	1:B:436:ASP:N	2.50	0.45
1:B:550:SER:HB2	1:B:554:SER:HB3	1.98	0.45
1:B:619:ASN:OD1	1:B:620:LYS:N	2.50	0.45
1:B:557:ASN:OD1	1:B:557:ASN:N	2.51	0.44
1:A:456:ILE:HG22	1:A:457:ASN:N	2.32	0.44
1:A:197:ARG:NH2	1:A:300:GLU:O	2.50	0.44
1:A:6:GLN:HB3	1:A:348:PHE:HE1	1.82	0.44
1:A:568:ASP:OD1	1:A:569:SER:N	2.51	0.44
1:A:580:ILE:HG22	1:A:677:TRP:CZ2	2.52	0.44
1:A:115:ASP:OD2	1:B:503:ARG:NH1	2.46	0.44
1:A:281:ILE:HG12	1:A:286:VAL:HB	2.00	0.44
1:B:660:LYS:HB3	1:B:686:GLU:HB3	1.99	0.44
1:A:178:LYS:NZ	1:A:182:GLU:OE2	2.39	0.43
1:B:94:ASP:HB2	1:B:95:ILE:HD12	1.98	0.43
1:A:30:TYR:HB2	1:A:139:LEU:HD21	2.00	0.43
1:A:436:ASP:N	1:A:436:ASP:OD1	2.52	0.43
1:A:4:GLN:O	1:A:7:HIS:N	2.51	0.43
1:B:137:SER:N	1:B:323:ILE:HG21	2.33	0.43
1:B:262:LEU:HD21	1:B:336:ASN:OD1	2.18	0.43
1:A:279:GLU:O	1:A:283:SER:OG	2.23	0.43
1:A:428:ILE:HG12	1:A:433:ASN:HD21	1.84	0.43
1:B:38:THR:O	1:B:38:THR:OG1	2.36	0.43
1:A:557:ASN:OD1	1:A:557:ASN:N	2.52	0.43
1:A:71:LEU:HD23	1:A:112:ASP:O	2.18	0.43
1:A:35:PHE:O	1:A:40:PRO:HA	2.19	0.42
1:A:404:ILE:HB	1:A:405:PRO:HD3	2.00	0.42
1:A:275:ASN:O	1:A:279:GLU:HG3	2.19	0.42

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:117:THR:HG21	1:B:503:ARG:HG2	2.00	0.42
1:B:428:ILE:HD11	1:B:437:PHE:CE1	2.54	0.42
1:B:623:GLU:HG2	1:B:626:PHE:CB	2.49	0.42
1:A:198:TYR:C	1:A:298:ILE:HD11	2.40	0.42
1:B:456:ILE:HG22	1:B:457:ASN:N	2.34	0.42
1:A:627:ILE:HG23	1:A:628:GLU:H	1.85	0.42
1:A:503:ARG:HG2	1:B:117:THR:HG21	2.02	0.42
1:B:508:THR:HG21	1:B:522:LEU:HB2	2.02	0.42
1:B:521:GLY:O	1:B:697:PHE:CD1	2.73	0.41
1:A:249:ASP:OD1	1:A:251:ASN:N	2.53	0.41
1:A:124:PRO:O	1:A:128:LEU:HG	2.21	0.41
1:A:41:TRP:HB2	1:B:555:PHE:HE1	1.86	0.41
1:A:612:TYR:O	1:A:614:THR:HG23	2.21	0.41
1:A:503:ARG:NH1	1:B:115:ASP:OD2	2.54	0.41
1:B:210:MET:SD	1:B:223:ILE:HD12	2.62	0.40
1:B:558:ASP:HB2	1:B:652:ARG:HH21	1.86	0.40
1:A:352:LEU:HB2	1:A:355:VAL:HG21	2.04	0.40
1:A:376:ILE:O	1:A:385:ARG:NE	2.54	0.40
1:B:10:PHE:CD1	1:B:350:ILE:HD13	2.56	0.40

There are no symmetry-related clashes.

### 5.3 Torsion angles

#### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	693/704 (98%)	657 (95%)	31 (4%)	5 (1%)	22 56
1	B	696/704 (99%)	664 (95%)	30 (4%)	2 (0%)	41 72
All	All	1389/1408 (99%)	1321 (95%)	61 (4%)	7 (0%)	29 62

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	351	VAL
1	A	4	GLN
1	A	351	VAL
1	A	350	ILE
1	B	39	LEU
1	A	551	VAL
1	A	527	PRO

### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	592/632 (94%)	577 (98%)	15 (2%)	47 72
1	B	600/632 (95%)	589 (98%)	11 (2%)	59 78
All	All	1192/1264 (94%)	1166 (98%)	26 (2%)	52 74

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	163	TYR
1	A	197	ARG
1	A	203	TYR
1	A	230	PHE
1	A	258	LYS
1	A	309	PHE
1	A	351	VAL
1	A	419	ASN
1	A	420	ASN
1	A	433	ASN
1	A	451	ASN
1	A	504	ARG
1	A	551	VAL
1	A	633	PHE
1	A	673	TYR
1	B	203	TYR
1	B	258	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	298	ILE
1	B	351	VAL
1	B	420	ASN
1	B	430	ASN
1	B	504	ARG
1	B	553	LEU
1	B	670	ASN
1	B	671	VAL
1	B	673	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [\(i\)](#)

There are no ligands in this entry.

### 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [\(i\)](#)

### 6.1 Protein, DNA and RNA chains [\(i\)](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	695/704 (98%)	0.25	10 (1%) 75 74	54, 100, 138, 159	0
1	B	698/704 (99%)	0.13	2 (0%) 94 94	51, 86, 122, 148	0
All	All	1393/1408 (98%)	0.19	12 (0%) 84 84	51, 94, 133, 159	0

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	526	GLN	3.8
1	A	221	LEU	3.1
1	A	689	ALA	3.0
1	A	688	LEU	2.9
1	A	550	SER	2.7
1	B	551	VAL	2.7
1	A	687	VAL	2.5
1	A	231	ILE	2.5
1	A	571	LEU	2.3
1	A	213	ALA	2.2
1	A	176	ILE	2.1
1	B	432	ASN	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [\(i\)](#)

There are no ligands in this entry.

## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.