

Full wwPDB X-ray Structure Validation Report (i)

May 13, 2024 – 06:22 PM JST

PDB ID : 8YHR

Title : DHODH in complex with furocoumavirin

Authors: Hara, K.; Okumura, H.; Nakahara, M.; Sato, M.; Hashimoto, H.; Osada, H.;

Watanabe, K.

Deposited on : 2024-02-28

Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.36.2buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36.2

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY\ DIFFRACTION$

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},{\rm resolution\ range}(\mathring{\rm A})) \end{array}$
R_{free}	130704	4298 (1.70-1.70)
Clashscore	141614	4695 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)
RSRZ outliers	127900	4222 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	٨	207	3%		
1	А	387	87%	•	9%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	ACT	A	409	_	-	X	-



2 Entry composition (i)

There are 8 unique types of molecules in this entry. The entry contains 5925 atoms, of which 2841 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Dihydroorotate dehydrogenase (quinone), mitochondrial.


Mol	Chain	Residues			Atoms	S			ZeroOcc	AltConf	Trace
1	Δ	353	Total	С	Н	N	О	S	0	0	0
1	11	000	5462	1697	2762	498	501	4		U	U

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	9	MET	-	initiating methionine	UNP Q02127
A	10	GLY	-	expression tag	UNP Q02127
A	11	SER	-	expression tag	UNP Q02127
A	12	SER	-	expression tag	UNP Q02127
A	13	HIS	-	expression tag	UNP Q02127
A	14	HIS	-	expression tag	UNP Q02127
A	15	HIS	-	expression tag	UNP Q02127
A	16	HIS	-	expression tag	UNP Q02127
A	17	HIS	-	expression tag	UNP Q02127
A	18	HIS	-	expression tag	UNP Q02127
A	19	SER	-	expression tag	UNP Q02127
A	20	SER	-	expression tag	UNP Q02127
A	21	GLY	-	expression tag	UNP Q02127
A	22	LEU	-	expression tag	UNP Q02127
A	23	VAL	-	expression tag	UNP Q02127
A	24	PRO	-	expression tag	UNP Q02127
A	25	ARG	-	expression tag	UNP Q02127
A	26	GLY	-	expression tag	UNP Q02127
A	27	SER	-	expression tag	UNP Q02127
A	28	HIS	-	expression tag	UNP Q02127


• Molecule 2 is OROTIC ACID (three-letter code: ORO) (formula: $C_5H_4N_2O_4$) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues		Ato	oms	}	ZeroOcc	AltConf	
9	Λ	1	Total	С	Н	N	О	0	0
2	A	1	14	5	3	2	4	0	0


 \bullet Molecule 3 is FLAVIN MONONUCLEOTIDE (three-letter code: FMN) (formula: $C_{17}H_{21}N_4O_9P)$ (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues		A	tom	ıs			ZeroOcc	AltConf
3	A	1	Total 50	C 17	H 19	N 4	O 9	P 1	0	0


• Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).





Mol	Chain	Residues	A	ton	ns		ZeroOcc	AltConf
4	A	1	Total 14		H 8		0	0
4	A	1	Total 14				0	0
4	A	1	Total 14	C 3	H 8	O 3	0	0

 \bullet Molecule 5 is ACETATE ION (three-letter code: ACT) (formula: $\mathrm{C_2H_3O_2}).$



Mol	Chain	Residues	A	tor	ns		ZeroOcc	AltConf
5	Λ	1	Total	С	Н	О	0	0
)	A	1	7	2	3	2		


Continued on next page...



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C H O 7 2 3 2	0	0
5	A	1	Total C H O 7 2 3 2	0	0
5	A	1	Total C H O 7 2 3 2	0	0
5	A	1	Total C H O 7 2 3 2	0	0


• Molecule 6 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Ato	oms		ZeroOcc	AltConf
6	A	1	Total 5	O 4	S 1	0	0

• Molecule 7 is 4-methyl-8-[(S)-oxidanyl(phenyl)methyl]-9-phenyl-furo[2,3-h]chromen-2 -one (three-letter code: A1LYU) (formula: $C_{25}H_{18}O_4$) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	A	A ton	ns		ZeroOcc	AltConf
7	Λ	1	Total	С	Н	О	0	0
'	A	1	47	25	18	4		U

• Molecule 8 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	270	Total O 270 270	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Dihydroorotate dehydrogenase (quinone), mitochondrial





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants	90.63Å 90.63Å 122.18Å	Donositon
a, b, c, α , β , γ	90.00° 90.00° 120.00°	Depositor
Resolution (Å)	36.40 - 1.70	Depositor
rtesolution (A)	48.21 - 1.70	EDS
% Data completeness	98.6 (36.40-1.70)	Depositor
(in resolution range)	96.5 (48.21-1.70)	EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.03 (at 1.70Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
P. P.	0.146 , 0.165	Depositor
R, R_{free}	0.149 , 0.166	DCC
R_{free} test set	1990 reflections (3.14%)	wwPDB-VP
Wilson B-factor (Å ²)	17.9	Xtriage
Anisotropy	0.183	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.45, 53.7	EDS
L-test for twinning ²	$< L > = 0.50, < L^2> = 0.33$	Xtriage
Estimated twinning fraction	0.018 for -h,-k,l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	5925	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.44% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ACT, GOL, ORO, SO4, FMN, A1LYU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

1	rai	Chain	Bond	lengths	Bond angles		
101	MIOI		RMSZ	# Z > 5	RMSZ	# Z > 5	
	1	A	0.43	0/2744	0.62	0/3707	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2700	2762	2761	11	0
2	A	11	3	3	0	0
3	A	31	19	19	0	0
4	A	18	24	24	1	0
5	A	20	15	15	2	0
6	A	5	0	0	0	0
7	A	29	18	0	0	0
8	A	270	0	0	5	1
All	All	3084	2841	2822	12	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (12) close contacts within the same asymmetric unit are listed below, sorted by their clash



magnitude.

Atom-1	Atom-2	$\begin{array}{c} \text{Interatomic} \\ \text{distance (Å)} \end{array}$	Clash overlap (Å)
1:A:50:ASP:OD1	8:A:501:HOH:O	2.06	0.74
1:A:34:GLU:OE2	1:A:371:ARG:NH2	2.24	0.61
1:A:265:GLU:OE1	8:A:502:HOH:O	2.16	0.59
1:A:56:ARG:NE	8:A:503:HOH:O	2.36	0.57
1:A:199:ARG:NH1	1:A:239:GLU:OE2	2.40	0.55
1:A:149:ASN:HD21	5:A:409:ACT:H1	1.71	0.55
1:A:79:GLU:OE2	1:A:81:ARG:NH1	2.44	0.50
1:A:245:ARG:HG2	1:A:248:ARG:NH1	2.33	0.44
1:A:32:GLY:H	4:A:405:GOL:C2	2.32	0.43
5:A:409:ACT:OXT	8:A:503:HOH:O	2.21	0.43
1:A:281:VAL:HA	1:A:282:THR:HA	1.94	0.40
1:A:81:ARG:NH2	8:A:515:HOH:O	2.54	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	1200111 2		$egin{array}{c} { m Clash} \\ { m overlap} \ ({ m \AA}) \end{array}$
8:A:506:HOH:O	8:A:513:HOH:O[5_677]	2.04	0.16

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	349/387 (90%)	338 (97%)	11 (3%)	0	100 100	

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar



resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	284/313 (91%)	282 (99%)	2 (1%)	84 77	

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	33	ASP
1	A	355	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



N / - 1	Т	Clasica	Das	T :1-	В	ond leng	gths	В	ond ang	gles
Mol	Type	Chain	Res	Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	GOL	A	405	-	5,5,5	0.49	0	5,5,5	0.29	0
4	GOL	A	403	-	5,5,5	0.58	0	5,5,5	0.52	0
4	GOL	A	404	-	5,5,5	0.51	0	5,5,5	0.21	0
7	A1LYU	A	412	-	28,33,33	2.21	12 (42%)	35,48,48	1.90	10 (28%)
5	ACT	A	409	-	3,3,3	0.83	0	3,3,3	0.50	0
5	ACT	A	406	-	3,3,3	1.01	0	3,3,3	1.00	0
6	SO4	A	411	-	4,4,4	0.14	0	6,6,6	0.18	0
2	ORO	A	401	-	9,11,11	1.42	2 (22%)	8,15,15	2.12	2 (25%)
5	ACT	A	407	-	3,3,3	0.77	0	3,3,3	0.87	0
5	ACT	A	408	-	3,3,3	0.87	0	3,3,3	0.79	0
3	FMN	A	402	-	33,33,33	1.03	2 (6%)	48,50,50	1.26	8 (16%)
5	ACT	A	410	-	3,3,3	0.83	0	3,3,3	0.80	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	405	-	-	2/4/4/4	-
4	GOL	A	403	-	-	2/4/4/4	-
4	GOL	A	404	-	-	4/4/4/4	-
7	A1LYU	A	412	-	-	0/8/12/12	0/5/5/5
2	ORO	A	401	-	-	4/4/4/4	0/1/1/1
3	FMN	A	402	-	-	4/18/18/18	0/3/3/3

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	$\operatorname{Ideal}(\text{\AA})$
7	A	412	A1LYU	O02-CE3	4.37	1.45	1.38
7	A	412	A1LYU	C03-C01	4.32	1.40	1.35
3	A	402	FMN	C4A-N5	4.02	1.38	1.30
7	A	412	A1LYU	CG-C05	3.61	1.54	1.49
7	A	412	A1LYU	CG-CD2	3.46	1.48	1.42
7	A	412	A1LYU	C12-C11	-3.27	1.46	1.52
7	A	412	A1LYU	CZ3-C01	3.24	1.50	1.45
2	A	401	ORO	O72-C7	-2.72	1.22	1.30
7	A	412	A1LYU	CH2-CZ3	2.62	1.44	1.39
7	A	412	A1LYU	C15-C13	2.51	1.44	1.38
7	A	412	A1LYU	C03-C04	2.49	1.49	1.44

Continued on next page...



 $Continued\ from\ previous\ page...$

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\text{\AA})$	$\operatorname{Ideal}(ext{\AA})$
3	A	402	FMN	C10-N1	2.16	1.37	1.33
7	A	412	A1LYU	C10-C08	2.12	1.43	1.38
7	A	412	A1LYU	C02-C01	2.05	1.54	1.50
7	A	412	A1LYU	C06-C05	2.03	1.43	1.39
2	A	401	ORO	O4-C4	-2.02	1.19	1.24

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
7	A	412	A1LYU	C06-C05-CG	-5.11	113.03	120.77
7	A	412	A1LYU	C04-C03-C01	-4.96	118.81	123.12
2	A	401	ORO	C5-C4-N3	-4.71	118.58	124.08
2	A	401	ORO	C6-C5-C4	3.37	118.91	116.73
7	A	412	A1LYU	CH2-CZ3-C01	-3.08	119.93	124.27
3	A	402	FMN	C4-C4A-N5	3.00	122.50	118.23
3	A	402	FMN	C4A-C10-N10	2.78	120.55	116.48
7	A	412	A1LYU	C07-C05-CG	2.75	124.93	120.77
3	A	402	FMN	C10-C4A-N5	-2.67	119.19	124.86
7	A	412	A1LYU	C14-C12-C13	2.64	121.58	118.29
3	A	402	FMN	C4A-C4-N3	2.56	119.68	113.19
7	A	412	A1LYU	CZ2-CE2-CD2	2.54	122.05	117.84
3	A	402	FMN	C9A-C5A-N5	-2.46	119.76	122.43
3	A	402	FMN	C10-N1-C2	2.42	121.75	116.90
7	A	412	A1LYU	O02-C04-C03	2.32	120.15	117.17
7	A	412	A1LYU	CG-CD2-CE2	-2.29	107.31	110.52
3	A	402	FMN	C4-N3-C2	-2.22	121.54	125.64
3	A	402	FMN	C4A-C10-N1	-2.09	119.89	124.73
7	A	412	A1LYU	CH2-CZ3-CE3	2.06	120.88	118.42
7	A	412	A1LYU	C07-C05-C06	2.05	121.68	117.59

There are no chirality outliers.

All (16) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401	ORO	N1-C6-C7-O72
2	A	401	ORO	C5-C6-C7-O71
2	A	401	ORO	C5-C6-C7-O72
4	A	403	GOL	O1-C1-C2-C3
4	A	404	GOL	C1-C2-C3-O3
2	A	401	ORO	N1-C6-C7-O71
4	A	404	GOL	O1-C1-C2-C3
4	A	405	GOL	C1-C2-C3-O3

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	A	403	GOL	O1-C1-C2-O2
4	A	404	GOL	O2-C2-C3-O3
4	A	405	GOL	O2-C2-C3-O3
3	A	402	FMN	C2'-C3'-C4'-O4'
4	A	404	GOL	O1-C1-C2-O2
3	A	402	FMN	O3'-C3'-C4'-O4'
3	A	402	FMN	C4'-C5'-O5'-P
3	A	402	FMN	O3'-C3'-C4'-C5'


There are no ring outliers.

2 monomers are involved in 3 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	405	GOL	1	0
5	A	409	ACT	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	# RSRZ > 2		$OWAB(A^2)$	Q < 0.9	
1	A	353/387 (91%)	-0.21	12 (3%)	45	50	12, 19, 42, 86	0

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	71	ARG	7.9
1	A	69	ARG	7.1
1	A	188	VAL	5.4
1	A	73	GLN	4.6
1	A	72	PHE	4.2
1	A	70	ALA	4.0
1	A	66	LEU	3.5
1	A	68	PRO	3.4
1	A	227	ALA	2.9
1	A	225	GLY	2.8
1	A	31	THR	2.3
1	A	212	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,




median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
4	GOL	A	405	6/6	0.68	0.34	54,73,82,89	0
4	GOL	A	403	6/6	0.89	0.10	32,50,58,60	0
5	ACT	A	408	4/4	0.90	0.08	40,48,56,57	0
4	GOL	A	404	6/6	0.91	0.13	37,54,69,82	0
5	ACT	A	409	4/4	0.91	0.10	28,33,59,59	0
5	ACT	A	407	4/4	0.93	0.10	27,32,37,38	0
5	ACT	A	410	4/4	0.94	0.11	40,51,61,61	0
5	ACT	A	406	4/4	0.97	0.09	27,30,32,32	0
7	A1LYU	A	412	29/29	0.97	0.07	13,19,24,28	0
2	ORO	A	401	11/11	0.98	0.06	11,14,17,18	0
6	SO4	A	411	5/5	0.99	0.04	30,31,40,43	0
3	FMN	A	402	31/31	0.99	0.08	10,12,16,17	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.














6.5 Other polymers (i)

There are no such residues in this entry.

