



# wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 9, 2020 – 10:23 AM BST

PDB ID : 1Y08  
Title : Structure of the C-terminal domain of human thrombospondin-2  
Authors : Carlson, C.B.; Bernstein, D.A.; Annis, D.S.; Misenheimer, T.M.; Hannah, B.A.; Mosher, D.F.; Keck, J.L.  
Deposited on : 2005-01-26  
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.13.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	634	
2	B	3	
3	C	2	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	B	1	-	-	-	X
2	NAG	B	2	-	-	-	X
2	MAN	B	3	X	-	-	-
3	NAG	C	2	-	-	-	X

## 2 Entry composition i

There are 6 unique types of molecules in this entry. The entry contains 5085 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called thrombospondin-2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	621	4844	2949	837	1017	36	5	0	0	0

There are 17 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	548	ALA	-	cloning artifact	UNP P35442
A	549	ASP	-	cloning artifact	UNP P35442
A	550	PRO	-	cloning artifact	UNP P35442
A	675	MSE	MET	modified residue	UNP P35442
A	964	MSE	MET	modified residue	UNP P35442
A	1041	MSE	MET	modified residue	UNP P35442
A	1134	MSE	MET	modified residue	UNP P35442
A	1159	MSE	MET	modified residue	UNP P35442
A	1173	ALA	-	expression tag	UNP P35442
A	1174	ALA	-	expression tag	UNP P35442
A	1175	GLY	-	expression tag	UNP P35442
A	1176	HIS	-	expression tag	UNP P35442
A	1177	HIS	-	expression tag	UNP P35442
A	1178	HIS	-	expression tag	UNP P35442
A	1179	HIS	-	expression tag	UNP P35442
A	1180	HIS	-	expression tag	UNP P35442
A	1181	HIS	-	expression tag	UNP P35442

- Molecule 2 is an oligosaccharide called alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	3	39	22	2	15	0	0	0

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	C	2	28	16	2	10	0	0	0

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
4	A	1	14	8	1	5	0	0
4	A	1	14	8	1	5	0	0

- Molecule 5 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	30	Total 30	Ca 30	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	116	Total 116	O 116	0	0







## 4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	93.44Å 121.59Å 155.33Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.60 26.86 – 2.60	Depositor EDS
% Data completeness (in resolution range)	95.0 (20.00-2.60) 94.9 (26.86-2.60)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.08	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.59 (at 2.60Å)	Xtrriage
Refinement program	REFMAC 5.1.24	Depositor
R, $R_{free}$	0.218 , 0.284 0.216 , 0.275	Depositor DCC
$R_{free}$ test set	2575 reflections (9.85%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	41.8	Xtrriage
Anisotropy	0.803	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 67.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	5085	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	68.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.52% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, NAG, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.57	0/4962	0.87	27/6755 (0.4%)

There are no bond length outliers.

The worst 5 of 27 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	696	ASP	CB-CG-OD2	6.86	124.47	118.30
1	A	866	ASP	CB-CG-OD2	6.72	124.35	118.30
1	A	738	ASP	CB-CG-OD2	6.51	124.16	118.30
1	A	797	ASP	CB-CG-OD2	5.89	123.60	118.30
1	A	718	ASP	CB-CG-OD2	5.88	123.59	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4844	0	4248	87	0
2	B	39	0	34	0	0
3	C	28	0	25	0	0
4	A	28	0	26	5	0
5	A	30	0	0	0	0
6	A	116	0	0	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	5085	0	4333	87	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 87 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:597:VAL:HB	1:A:598:PRO:CD	1.90	1.02
1:A:597:VAL:HB	1:A:598:PRO:HD3	1.03	0.99
1:A:597:VAL:CB	1:A:598:PRO:HD3	1.93	0.97
1:A:719:ASN:H	1:A:728:GLN:HE22	1.25	0.83
1:A:608:PRO:O	1:A:609:ARG:HB2	1.79	0.81

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	619/634 (98%)	550 (89%)	57 (9%)	12 (2%)	<b>8</b> <b>15</b>

5 of 12 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	597	VAL
1	A	598	PRO
1	A	609	ARG
1	A	716	ILE
1	A	584	ASN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	540/544 (99%)	483 (89%)	57 (11%)	<b>6</b> <b>12</b>

5 of 57 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	750	THR
1	A	824	ARG
1	A	1131	LYS
1	A	759	LEU
1	A	811	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	842	HIS
1	A	883	ASN
1	A	990	GLN
1	A	834	HIS
1	A	911	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

5 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	B	1	1,2	14,14,15	0.71	1 (7%)	17,19,21	1.26	1 (5%)
2	NAG	B	2	2	14,14,15	0.63	0	17,19,21	0.96	1 (5%)
2	MAN	B	3	2	11,11,12	0.65	0	15,15,17	2.05	3 (20%)
3	NAG	C	1	1,3	14,14,15	0.71	0	17,19,21	1.17	1 (5%)
3	NAG	C	2	3	14,14,15	0.50	0	17,19,21	0.71	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	1,2	-	5/6/23/26	0/1/1/1
2	NAG	B	2	2	-	4/6/23/26	0/1/1/1
2	MAN	B	3	2	1/1/4/5	0/2/19/22	0/1/1/1
3	NAG	C	1	1,3	-	4/6/23/26	0/1/1/1
3	NAG	C	2	3	-	4/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1	NAG	C1-C2	2.05	1.55	1.52

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	3	MAN	C1-O5-C5	5.10	119.10	112.19
2	B	3	MAN	C1-C2-C3	4.78	115.54	109.67
2	B	1	NAG	C1-O5-C5	4.04	117.67	112.19
2	B	3	MAN	O5-C1-C2	3.16	115.65	110.77
3	C	1	NAG	C4-C3-C2	3.08	115.53	111.02

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	B	3	MAN	C1


5 of 17 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	C	1	NAG	C8-C7-N2-C2
2	B	2	NAG	C8-C7-N2-C2
2	B	2	NAG	O7-C7-N2-C2
2	B	1	NAG	C8-C7-N2-C2
2	B	1	NAG	O7-C7-N2-C2

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





## 5.6 Ligand geometry [i](#)

Of 32 ligands modelled in this entry, 30 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	NAG	A	6	1	14,14,15	0.67	0	17,19,21	0.95	0
4	NAG	A	7	1	14,14,15	0.59	0	17,19,21	1.19	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	6	1	-	4/6/23/26	0/1/1/1
4	NAG	A	7	1	-	3/6/23/26	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	7	NAG	C1-O5-C5	3.68	117.18	112.19
4	A	7	NAG	O5-C1-C2	-2.04	108.07	111.29

There are no chirality outliers.

5 of 7 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	7	NAG	C8-C7-N2-C2
4	A	7	NAG	O7-C7-N2-C2
4	A	6	NAG	C8-C7-N2-C2
4	A	6	NAG	O7-C7-N2-C2
4	A	6	NAG	C4-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	6	NAG	5	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	616/634 (97%)	1.23	104 (16%) <b>1</b> <b>1</b>	40, 61, 122, 158	0

The worst 5 of 104 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	583	GLY	12.9
1	A	973	THR	11.0
1	A	972	THR	10.6
1	A	582	LEU	10.0
1	A	641	THR	8.7

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.


### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	NAG	B	2	14/15	0.40	0.88	123,127,131,132	0
2	MAN	B	3	11/12	0.62	0.37	133,136,137,138	0
2	NAG	B	1	14/15	0.64	0.43	114,119,121,122	0
3	NAG	C	2	14/15	0.70	0.52	115,123,126,127	0
3	NAG	C	1	14/15	0.80	0.27	80,89,98,107	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	NAG	A	7	14/15	0.67	0.34	140,146,149,149	0
4	NAG	A	6	14/15	0.75	0.33	103,117,128,133	0
5	CA	A	1184	1/1	0.79	0.13	74,74,74,74	0
5	CA	A	1202	1/1	0.81	0.10	71,71,71,71	0
5	CA	A	1188	1/1	0.86	0.09	55,55,55,55	0
5	CA	A	1187	1/1	0.88	0.08	70,70,70,70	0
5	CA	A	1199	1/1	0.90	0.09	59,59,59,59	0
5	CA	A	1197	1/1	0.90	0.09	63,63,63,63	0
5	CA	A	1201	1/1	0.91	0.10	70,70,70,70	0
5	CA	A	1203	1/1	0.91	0.06	81,81,81,81	0
5	CA	A	1194	1/1	0.92	0.10	55,55,55,55	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	CA	A	1207	1/1	0.93	0.07	65,65,65,65	0
5	CA	A	1208	1/1	0.93	0.09	58,58,58,58	0
5	CA	A	1210	1/1	0.93	0.13	51,51,51,51	0
5	CA	A	1206	1/1	0.93	0.11	62,62,62,62	0
5	CA	A	1205	1/1	0.94	0.15	72,72,72,72	0
5	CA	A	1211	1/1	0.94	0.06	116,116,116,116	0
5	CA	A	1182	1/1	0.95	0.07	46,46,46,46	0
5	CA	A	1186	1/1	0.95	0.13	58,58,58,58	0
5	CA	A	1200	1/1	0.96	0.05	62,62,62,62	0
5	CA	A	1209	1/1	0.96	0.13	45,45,45,45	0
5	CA	A	1192	1/1	0.97	0.10	55,55,55,55	0
5	CA	A	1183	1/1	0.97	0.08	50,50,50,50	0
5	CA	A	1189	1/1	0.97	0.07	61,61,61,61	0
5	CA	A	1204	1/1	0.97	0.13	70,70,70,70	0
5	CA	A	1193	1/1	0.98	0.07	57,57,57,57	0
5	CA	A	1198	1/1	0.98	0.10	50,50,50,50	0
5	CA	A	1196	1/1	0.98	0.11	52,52,52,52	0
5	CA	A	1191	1/1	0.98	0.07	53,53,53,53	0
5	CA	A	1190	1/1	0.99	0.11	63,63,63,63	0
5	CA	A	1185	1/1	0.99	0.12	57,57,57,57	0
5	CA	A	1195	1/1	0.99	0.11	53,53,53,53	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.