



# wwPDB X-ray Structure Validation Summary Report i

May 16, 2020 – 10:02 am BST

PDB ID : 4YO3  
Title : Enteroaggregative Escherichia Coli TssA N-terminal fragment  
Authors : Durand, E.; Zoued, A.; Spinelli, S.; Douzi, B.; Brunet, Y.R.; Bebeacua, C.; Legrand, P.; Journet, L.; Mignot, T.; Cambillau, C.; Cascales, E.  
Deposited on : 2015-03-11  
Resolution : 3.37 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>  
with specific help available everywhere you see the i symbol.

---

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## X-RAY DIFFRACTION

The reported resolution of this entry is 3.37 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1691 (3.46-3.30)
Clashscore	141614	1762 (3.46-3.30)
Ramachandran outliers	138981	1732 (3.46-3.30)
Sidechain outliers	138945	1731 (3.46-3.30)
RSRZ outliers	127900	1635 (3.46-3.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



Continued on next page...

Continued from previous page...



## 2 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 14496 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called TssA.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	157	Total	C 1208	N 774	O 210	S 219	Se 1	0	0	0
1	B	157	Total	C 1208	N 774	O 210	S 219	Se 1	0	0	0
1	C	157	Total	C 1208	N 774	O 210	S 219	Se 1	0	0	0
1	D	157	Total	C 1208	N 774	O 210	S 219	Se 1	0	0	0
1	E	157	Total	C 1208	N 774	O 210	S 219	Se 1	0	0	0
1	F	157	Total	C 1208	N 774	O 210	S 219	Se 1	0	0	0
1	G	157	Total	C 1208	N 774	O 210	S 219	Se 1	0	0	0
1	H	157	Total	C 1208	N 774	O 210	S 219	Se 1	0	0	0
1	I	157	Total	C 1208	N 774	O 210	S 219	Se 1	0	0	0
1	J	157	Total	C 1208	N 774	O 210	S 219	Se 1	0	0	0
1	K	157	Total	C 1208	N 774	O 210	S 219	Se 1	0	0	0
1	L	157	Total	C 1208	N 774	O 210	S 219	Se 1	0	0	0


### 3 Residue-property plots [\(i\)](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: TssA




- Molecule 1: TssA




- Molecule 1: TssA




- Molecule 1: TssA




- Molecule 1: TssA




- Molecule 1: TssA




- Molecule 1: TssA



- Molecule 1: TssA




- Molecule 1: TssA






- Molecule 1: TssA



- Molecule 1: TssA



- Molecule 1: TssA



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	134.13Å    178.05Å    106.76Å 90.00°    90.00°    90.00°	Depositor
Resolution (Å)	28.39 – 3.37 28.16 – 3.37	Depositor EDS
% Data completeness (in resolution range)	100.0 (28.39-3.37) 100.0 (28.16-3.37)	Depositor EDS
$R_{merge}$	0.15	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) >$ <sup>1</sup>	3.37 (at 3.39Å)	Xtriage
Refinement program	BUSTER 2.10.1	Depositor
$R$ , $R_{free}$	0.250 , 0.261 0.277 , 0.284	Depositor DCC
$R_{free}$ test set	1839 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	94.3	Xtriage
Anisotropy	0.584	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.27 , 69.0	EDS
L-test for twinning <sup>2</sup>	$<  L  > = 0.47$ , $< L^2 > = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	14496	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	111.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.48% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $< |L| >$ ,  $< L^2 >$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.52	0/1232	0.75	1/1673 (0.1%)
1	B	0.52	0/1232	0.72	0/1673
1	C	0.49	0/1232	0.72	0/1673
1	D	0.49	0/1232	0.72	0/1673
1	E	0.48	0/1232	0.70	1/1673 (0.1%)
1	F	0.47	0/1232	0.70	0/1673
1	G	0.50	0/1232	0.72	0/1673
1	H	0.47	0/1232	0.69	0/1673
1	I	0.47	0/1232	0.71	0/1673
1	J	0.47	0/1232	0.69	0/1673
1	K	0.46	0/1232	0.69	0/1673
1	L	0.54	0/1232	0.74	0/1673
All	All	0.49	0/14784	0.71	2/20076 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
1	C	0	1
1	D	0	1
1	E	0	1
1	F	0	1
1	G	0	1
1	H	0	1
1	I	0	2
1	J	0	1
1	K	0	1
1	L	0	1
All	All	0	13

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	312	SER	C-N-CA	5.09	134.42	121.70
1	E	312	SER	C-N-CA	5.02	134.24	121.70

There are no chirality outliers.

5 of 13 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	266	PRO	Peptide
1	B	266	PRO	Peptide
1	C	266	PRO	Peptide
1	D	266	PRO	Peptide
1	E	266	PRO	Peptide

## 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1208	0	1151	14	0
1	B	1208	0	1151	17	0
1	C	1208	0	1151	16	0
1	D	1208	0	1151	13	0
1	E	1208	0	1151	16	0
1	F	1208	0	1151	12	0
1	G	1208	0	1151	13	0
1	H	1208	0	1151	13	0
1	I	1208	0	1151	12	0
1	J	1208	0	1151	11	0
1	K	1208	0	1151	12	0
1	L	1208	0	1151	18	0
All	All	14496	0	13812	152	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 152 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:306:MSE:SE	1:B:306:MSE:CE	2.18	1.41
1:L:236:LEU:HD23	1:L:253:LEU:HD11	1.49	0.95
1:L:236:LEU:CD2	1:L:253:LEU:HD11	2.23	0.67
1:C:370:SER:O	1:C:374:GLN:HG2	1.95	0.66
1:G:311:ALA:C	1:G:313:HIS:H	1.99	0.65

There are no symmetry-related clashes.

### 5.3 Torsion angles [\(i\)](#)

#### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	155/157 (99%)	140 (90%)	10 (6%)	5 (3%)	4 24
1	B	155/157 (99%)	141 (91%)	9 (6%)	5 (3%)	4 24
1	C	155/157 (99%)	140 (90%)	10 (6%)	5 (3%)	4 24
1	D	155/157 (99%)	141 (91%)	9 (6%)	5 (3%)	4 24
1	E	155/157 (99%)	142 (92%)	8 (5%)	5 (3%)	4 24
1	F	155/157 (99%)	141 (91%)	9 (6%)	5 (3%)	4 24
1	G	155/157 (99%)	140 (90%)	9 (6%)	6 (4%)	3 20
1	H	155/157 (99%)	141 (91%)	9 (6%)	5 (3%)	4 24
1	I	155/157 (99%)	141 (91%)	8 (5%)	6 (4%)	3 20
1	J	155/157 (99%)	141 (91%)	9 (6%)	5 (3%)	4 24
1	K	155/157 (99%)	141 (91%)	9 (6%)	5 (3%)	4 24
1	L	155/157 (99%)	138 (89%)	13 (8%)	4 (3%)	5 28
All	All	1860/1884 (99%)	1687 (91%)	112 (6%)	61 (3%)	4 24

5 of 61 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	266	PRO

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	267	PRO
1	B	223	ILE
1	B	266	PRO
1	B	267	PRO

### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	118/129 (92%)	99 (84%)	19 (16%)	2 10
1	B	118/129 (92%)	100 (85%)	18 (15%)	2 12
1	C	118/129 (92%)	101 (86%)	17 (14%)	3 14
1	D	118/129 (92%)	102 (86%)	16 (14%)	3 16
1	E	118/129 (92%)	104 (88%)	14 (12%)	5 20
1	F	118/129 (92%)	103 (87%)	15 (13%)	4 18
1	G	118/129 (92%)	102 (86%)	16 (14%)	3 16
1	H	118/129 (92%)	102 (86%)	16 (14%)	3 16
1	I	118/129 (92%)	103 (87%)	15 (13%)	4 18
1	J	118/129 (92%)	103 (87%)	15 (13%)	4 18
1	K	118/129 (92%)	102 (86%)	16 (14%)	3 16
1	L	118/129 (92%)	105 (89%)	13 (11%)	6 24
All	All	1416/1548 (92%)	1226 (87%)	190 (13%)	4 16

5 of 190 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	245	GLU
1	G	305	ASP
1	K	361	MSE
1	F	273	ARG
1	F	347	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 23 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	264	GLN
1	G	243	GLN
1	K	339	GLN
1	F	339	GLN
1	G	313	HIS

### 5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [\(i\)](#)

There are no ligands in this entry.

### 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [\(i\)](#)

### 6.1 Protein, DNA and RNA chains [\(i\)](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	153/157 (97%)	0.04	4 (2%) 56 59	54, 76, 109, 162	0
1	B	153/157 (97%)	0.05	7 (4%) 32 36	57, 81, 116, 175	0
1	C	153/157 (97%)	0.12	6 (3%) 39 43	63, 102, 138, 171	0
1	D	153/157 (97%)	0.16	7 (4%) 32 36	65, 98, 130, 168	0
1	E	153/157 (97%)	0.09	2 (1%) 77 81	57, 95, 135, 170	0
1	F	153/157 (97%)	0.13	4 (2%) 56 59	69, 106, 145, 160	0
1	G	153/157 (97%)	0.08	6 (3%) 39 43	66, 89, 119, 176	0
1	H	153/157 (97%)	0.14	9 (5%) 22 26	74, 109, 149, 167	0
1	I	153/157 (97%)	0.59	16 (10%) 6 8	76, 134, 169, 181	0
1	J	153/157 (97%)	0.36	9 (5%) 22 26	93, 125, 159, 178	0
1	K	153/157 (97%)	0.41	12 (7%) 13 16	92, 122, 157, 178	0
1	L	153/157 (97%)	2.90	75 (49%) 0 0	121, 184, 211, 221	0
All	All	1836/1884 (97%)	0.42	157 (8%) 10 13	54, 106, 182, 221	0

The worst 5 of 157 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	L	221	SER	13.7
1	L	264	GLN	13.3
1	L	266	PRO	12.5
1	L	291	TYR	11.0
1	L	222	ALA	10.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [\(i\)](#)

There are no ligands in this entry.

### 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.