

wwPDB X-ray Structure Validation Summary Report (i)

Nov 12, 2024 – 02:18 PM JST

PDB ID : 8YR5

Title: Crystal structure of E. coli phosphatidylserine synthase in apo state

Authors: Kim, J.; Lee, E.; Cho, G.

Deposited on : 2024-03-20

Resolution : 2.83 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS: 3.0

buster-report : 1.1.7 (2018)

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)

CCP4 : 9.0.003 (Gargrove)

Density-Fitness : 1.0.11

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.39

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 2.83 Å.


Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Menic	(# Entries)	$(\# ext{Entries}, ext{ resolution range}(ext{Å}))$
R_{free}	164625	1367 (2.86-2.82)
Clashscore	180529	1455 (2.86-2.82)
Ramachandran outliers	177936	1422 (2.86-2.82)
Sidechain outliers	177891	1423 (2.86-2.82)
RSRZ outliers	164620	1368 (2.86-2.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	461	87%	7% • 5%
1	В	461	87%	8% • 5%
1	С	461	86%	8% • 5%
1	D	461	88%	8% •
1	Е	461	88%	8% • •
1	F	461	86%	8% • 5%





Continued from previous page...

Mol	Chain	Length	Quality of chain		
1		4.0.1	9%		
1	G	461	86%	8%	5%
1	Н	461	86%	8%	5%
-	_	101	⁹ 6		
1	1	461	87%	8%	• •
1	J	461	3%		
1	J	461	86%	9%	• •
1	K	461	86%	8%	5%
1	11	401	3%	0 %	<u></u>
1	L	461	87%	8%	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SO4	A	501	-	-	X	-
2	SO4	A	502	-	-	X	-
2	SO4	В	501	-	-	X	-
2	SO4	Е	501	-	-	X	-
2	SO4	G	502	-	-	X	-



2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 43895 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called CDP-diacylglycerol--serine O-phosphatidyltransferase.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	A	437	Total	С	N	О	S	0	0	0
1	A	437	3628	2309	658	650	11	U	0	
1	В	439	Total	С	N	О	S	0	0	0
1	Б	439	3643	2319	662	651	11	U	U	
1	С	440	Total	С	N	О	S	0	0	0
1		440	3651	2323	664	653	11	U	U	
1	D	446	Total	С	N	О	S	0	0	0
1	D	440	3685	2343	671	660	11		0	0
1	Е	445	Total	С	N	О	S	0	0	0
1	E	440	3676	2337	669	659	11		U	
1	F	437	Total	С	N	О	S	0	0	0
1	Г	457	3628	2309	658	650	11			U
1	G	437	Total	С	N	О	S	0	0	0
1	G	401	3623	2306	655	651	11	0	0	
1	Н	436	Total	С	N	О	S	0	0	0
1	11	450	3620	2305	656	648	11	0	0	
1	I	444	Total	С	N	О	S	0	0	0
1	1	444	3665	2331	665	658	11	U	U	
1	J	443	Total	С	N	О	S	0	0	0
1	9	440	3675	2340	668	656	11	0	0	
1	K	437	Total	С	N	О	S	0	0	0
1	17	401	3623	2306	655	651	11	0	U	U
1	L	441	Total	С	N	О	S	0	0	0
1	П	441	3658	2329	663	655	11	U	U	

There are 132 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual Comment		Reference
A	-9	MET	-	initiating methionine	UNP P23830
A	-8	GLY	-	expression tag	UNP P23830
A	-7	SER	-	expression tag	UNP P23830
A	-6	HIS	-	expression tag	UNP P23830
A	-5	HIS	-	expression tag	UNP P23830



 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled	Actual	Comment	Reference
A	-4	HIS	-	expression tag	UNP P23830
A	-3	HIS	-	expression tag	UNP P23830
A	-2	HIS	-	expression tag	UNP P23830
A	-1	HIS	-	expression tag	UNP P23830
A	0	GLY	-	expression tag	UNP P23830
A	1	SER	-	expression tag	UNP P23830
В	-9	MET	-	initiating methionine	UNP P23830
В	-8	GLY	-	expression tag	UNP P23830
В	-7	SER	-	expression tag	UNP P23830
В	-6	HIS	-	expression tag	UNP P23830
В	-5	HIS	-	expression tag	UNP P23830
В	-4	HIS	-	expression tag	UNP P23830
В	-3	HIS	-	expression tag	UNP P23830
В	-2	HIS	-	expression tag	UNP P23830
В	-1	HIS	-	expression tag	UNP P23830
В	0	GLY	-	expression tag	UNP P23830
В	1	SER	-	expression tag	UNP P23830
С	-9	MET	-	initiating methionine	UNP P23830
С	-8	GLY	-	expression tag	UNP P23830
С	-7	SER	-	expression tag	UNP P23830
С	-6	HIS	-	expression tag	UNP P23830
С	-5	HIS	-	expression tag	UNP P23830
С	-4	HIS	-	expression tag	UNP P23830
С	-3	HIS	-	expression tag	UNP P23830
С	-2	HIS	-	expression tag	UNP P23830
С	-1	HIS	-	expression tag	UNP P23830
С	0	GLY	-	expression tag	UNP P23830
С	1	SER	-	expression tag	UNP P23830
D	-9	MET	-	initiating methionine	UNP P23830
D	-8	GLY	-	expression tag	UNP P23830
D	-7	SER	-	expression tag	UNP P23830
D	-6	HIS	-	expression tag	UNP P23830
D	-5	HIS	-	expression tag	UNP P23830
D	-4	HIS	-	expression tag	UNP P23830
D	-3	HIS	-	expression tag	UNP P23830
D	-2	HIS	-	expression tag	UNP P23830
D	-1	HIS	-	expression tag	UNP P23830
D	0	GLY	-	expression tag	UNP P23830
D	1	SER		expression tag	UNP P23830
Е	-9	MET	-	initiating methionine	UNP P23830
Е	-8	GLY	-	expression tag	UNP P23830
Е	-7	SER	-	expression tag	UNP P23830



Continued from previous page...

Chain	Residue	Modelled Modelled	Actual	Comment	Reference
Е	-6	HIS	-	expression tag	UNP P23830
Е	-5	HIS	-	expression tag	UNP P23830
Е	-4	HIS	-	expression tag	UNP P23830
Е	-3	HIS	-	expression tag	UNP P23830
Е	-2	HIS	-	expression tag	UNP P23830
Е	-1	HIS	-	expression tag	UNP P23830
Е	0	GLY	-	expression tag	UNP P23830
Е	1	SER	-	expression tag	UNP P23830
F	-9	MET	-	initiating methionine	UNP P23830
F	-8	GLY	-	expression tag	UNP P23830
F	-7	SER	-	expression tag	UNP P23830
F	-6	HIS	-	expression tag	UNP P23830
F	-5	HIS	-	expression tag	UNP P23830
F	-4	HIS	-	expression tag	UNP P23830
F	-3	HIS	-	expression tag	UNP P23830
F	-2	HIS	-	expression tag	UNP P23830
F	-1	HIS	_	expression tag	UNP P23830
F	0	GLY	-	expression tag	UNP P23830
F	1	SER	-	expression tag	UNP P23830
G	-9	MET	_	initiating methionine	UNP P23830
G	-8	GLY	-	expression tag	UNP P23830
G	-7	SER	-	expression tag	UNP P23830
G	-6	HIS	-	expression tag	UNP P23830
G	-5	HIS	-	expression tag	UNP P23830
G	-4	HIS	-	expression tag	UNP P23830
G	-3	HIS	-	expression tag	UNP P23830
G	-2	HIS	-	expression tag	UNP P23830
G	-1	HIS	-	expression tag	UNP P23830
G	0	GLY	-	expression tag	UNP P23830
G	1	SER	-	expression tag	UNP P23830
H	-9	MET	-	initiating methionine	UNP P23830
H	-8	GLY	-	expression tag	UNP P23830
H	-7	SER	-	expression tag	UNP P23830
H	-6	HIS	-	expression tag	UNP P23830
H	-5	HIS	-	expression tag	UNP P23830
H	-4	HIS	-	expression tag	UNP P23830
H	-3	HIS	-	expression tag	UNP P23830
H	-2	HIS	-	expression tag	UNP P23830
Н	-1	HIS	-	expression tag	UNP P23830
H	0	GLY	-	expression tag	UNP P23830
H	1	SER	-	expression tag	UNP P23830
I	-9	MET	-	initiating methionine	UNP P23830



Continued from previous page...


Chain	Residue	Modelled Modelled	Actual	Comment	Reference
I	-8	GLY	-	expression tag	UNP P23830
I	-7	SER	-	expression tag	UNP P23830
I	-6	HIS	-	expression tag	UNP P23830
I	-5	HIS	-	expression tag	UNP P23830
I	-4	HIS	-	expression tag	UNP P23830
I	-3	HIS	-	expression tag	UNP P23830
I	-2	HIS	-	expression tag	UNP P23830
I	-1	HIS	-	expression tag	UNP P23830
I	0	GLY	-	expression tag	UNP P23830
I	1	SER	-	expression tag	UNP P23830
J	-9	MET	-	initiating methionine	UNP P23830
J	-8	GLY	-	expression tag	UNP P23830
J	-7	SER	-	expression tag	UNP P23830
J	-6	HIS	-	expression tag	UNP P23830
J	-5	HIS	-	expression tag	UNP P23830
J	-4	HIS	-	expression tag	UNP P23830
J	-3	HIS	-	expression tag	UNP P23830
J	-2	HIS	-	expression tag	UNP P23830
J	-1	HIS	-	expression tag	UNP P23830
J	0	GLY	-	expression tag	UNP P23830
J	1	SER	-	expression tag	UNP P23830
K	-9	MET	-	initiating methionine	UNP P23830
K	-8	GLY	-	expression tag	UNP P23830
K	-7	SER	-	expression tag	UNP P23830
K	-6	HIS	-	expression tag	UNP P23830
K	-5	HIS	-	expression tag	UNP P23830
K	-4	HIS	-	expression tag	UNP P23830
K	-3	HIS	-	expression tag	UNP P23830
K	-2	HIS	-	expression tag	UNP P23830
K	-1	HIS	-	expression tag	UNP P23830
K	0	GLY	ı	expression tag	UNP P23830
K	1	SER	-	expression tag	UNP P23830
L	-9	MET	ı	initiating methionine	UNP P23830
L	-8	GLY	-	expression tag	UNP P23830
L	-7	SER	-	expression tag	UNP P23830
L	-6	HIS	-	expression tag	UNP P23830
L	-5	HIS		expression tag	UNP P23830
L	-4	HIS	-	expression tag	UNP P23830
L	-3	HIS	-	expression tag	UNP P23830
L	-2	HIS	-	expression tag	UNP P23830
L	-1	HIS	-	expression tag	UNP P23830
L	0	GLY	-	expression tag	UNP P23830



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
L	1	SER	-	expression tag	UNP P23830

• Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O_4S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Ato	oms		ZeroOcc	AltConf
2	A	1	Total	О	S	0	0
	Λ	1	5	4	1	U	U
2	A	1	Total	Ο	S	0	0
	11	1	5	4	1	· ·	Ŭ
2	В	1	Total	Ο	S	0	0
	Б	1	5	4	1	· ·	-
2	В	1	Total	Ο	S	0	0
	D	1	5	4	1	Ŭ	Ü
2	С	1	Total	Ο	S	0	0
			5	4	1	Ü	Ü
2	$^{\rm C}$	1	Total	O	\mathbf{S}	0	0
_		_	5	4	1	Ů	Ů
2	D	1	Total	O	S	0	0
_	_	_	5	4	1	,	Ů
2	D	1	Total	O	S	0	0
_	_	_	5	4	1	, and the second	Ů
2	E	1	Total	O	S	0	0
	_	_	5	4	1		
2	E	1	Total	O	S	0	0
			5	4	1		



Continued from previous page...


Mol	Chain	Residues	Ato	oms		ZeroOcc	AltConf
2	F	1	Total	O 4	S	0	0
0	П	1	5 Total	4 O	1 S	0	0
2	F	1	5	4	1	0	0
2	G	1	Total	O	S	0	0
_			5	4	1	Ů	
2	G	1	Total 5	O 4	S 1	0	0
			Total	0	S		
2	G	1	5	4	1	0	0
2	Н	1	Total	О	S	0	0
	11	1	5	4	1	U	U
2	Н	1	Total	O	S	0	0
			5	4	1		_
2	I	1	Total 5	O 4	S 1	0	0
			Total	O	S		
2	I	1	5	4	1	0	0
2	J	1	Total	О	S	0	0
	J	1	5	4	1	U	U
2	K	1	Total	O	S	0	0
			5	4	1		_
2	K	1	Total 5	O 4	S 1	0	0
<u> </u>	_	_	Total	O	S		
2	L	1	5	4	1	0	0
2	L	1	Total	О	S	0	0
	П	1	5	4	1		U




3 Residue-property plots (i)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: CDP-diacylglycerol--serine O-phosphatidyltransferase











4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1	Depositor
Cell constants	94.04Å 94.10Å 194.75Å	Donositon
a, b, c, α , β , γ	$88.24^{\circ} 87.70^{\circ} 60.07^{\circ}$	Depositor
Resolution (Å)	48.64 - 2.83	Depositor
Resolution (A)	48.64 - 2.83	EDS
% Data completeness	98.2 (48.64-2.83)	Depositor
(in resolution range)	98.2 (48.64-2.83)	EDS
R_{merge}	0.19	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.43 (at 2.81Å)	Xtriage
Refinement program	REFMAC 5.8.0425	Depositor
R, R_{free}	0.242 , 0.285	Depositor
	0.242 , 0.280	DCC
R_{free} test set	6358 reflections $(4.70%)$	wwPDB-VP
Wilson B-factor (Å ²)	66.7	Xtriage
Anisotropy	0.021	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	$0.32 \; , 42.7$	EDS
L-test for twinning ²	$< L > = 0.47, < L^2> = 0.29$	Xtriage
	$0.014~\mathrm{for}~\mathrm{k,-h+k,l}$	
	0.014 for h-k,h,l	
	0.014 for -h+k,-h,l	
	0.014 for -k,h-k,l	
	0.023 for h,h-k,-l	
Estimated twinning fraction	0.083 for -h+k,k,-l	Xtriage
	0.019 for -h,-k,l	
	0.016 for k,h,-l	
	0.019 for -k,-h,-l	
	0.023 for -h,-h+k,-l	
	0.024 for h-k,-k,-l	
F_o, F_c correlation	0.92	EDS
Total number of atoms	43895	wwPDB-VP
Average B, all atoms (Å ²)	71.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.53% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Mol Chain		lengths	Bond angles		
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.36	0/3705	0.76	7/5014 (0.1%)	
1	В	0.36	0/3720	0.76	0/5033	
1	С	0.37	0/3728	0.78	6/5044 (0.1%)	
1	D	0.36	0/3763	0.76	4/5092 (0.1%)	
1	Е	0.36	0/3754	0.76	4/5081 (0.1%)	
1	F	0.35	0/3705	0.74	6/5014~(0.1%)	
1	G	0.35	0/3700	0.74	1/5008~(0.0%)	
1	Н	0.36	0/3697	0.76	5/5003 (0.1%)	
1	I	0.36	0/3743	0.78	4/5067 (0.1%)	
1	J	0.35	0/3753	0.77	3/5076 (0.1%)	
1	K	0.35	0/3700	0.75	3/5008 (0.1%)	
1	L	0.34	0/3736	0.72	1/5054~(0.0%)	
All	All	0.35	0/44704	0.76	44/60494 (0.1%)	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	D	0	1
1	Ε	0	4
1	Н	0	1
1	I	0	1
1	J	0	2
1	K	0	1
1	L	0	2
All	All	0	14

There are no bond length outliers.



The	worst	5	$\circ f$	44	hond	angle	outliers	are	listed	below.
1110	WOISU	U	OI	77	Dona	angic	Outilities	arc	nsucu	DCIOW.

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$Ideal(^{o})$
1	J	419	ARG	CG-CD-NE	10.70	134.28	111.80
1	A	254	LYS	CA-CB-CG	7.79	130.53	113.40
1	D	437	ARG	CB-CG-CD	6.94	129.64	111.60
1	Е	7	ARG	CG-CD-NE	6.64	125.75	111.80
1	A	329	ARG	NE-CZ-NH2	-6.42	117.09	120.30

There are no chirality outliers.

5 of 14 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	381	ARG	Sidechain
1	A	7	ARG	Sidechain
1	D	96	ARG	Sidechain
1	Е	7	ARG	Sidechain
1	Е	78	ARG	Sidechain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3628	0	3653	22	0
1	В	3643	0	3674	27	1
1	С	3651	0	3680	34	2
1	D	3685	0	3717	29	0
1	Е	3676	0	3704	35	0
1	F	3628	0	3653	37	0
1	G	3623	0	3645	22	0
1	Н	3620	0	3647	34	0
1	I	3665	0	3691	38	2
1	J	3675	0	3705	30	1
1	K	3623	0	3645	22	0
1	L	3658	0	3683	24	0
2	A	10	0	0	5	0
2	В	10	0	0	3	0
2	С	10	0	0	1	0
2	D	10	0	0	1	0



$\alpha \cdots 1$	c		
Continued	trom	nrevious	ทด.ดค

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	Е	10	0	0	3	0
2	F	10	0	0	2	0
2	G	15	0	0	2	0
2	Н	10	0	0	2	0
2	I	10	0	0	0	0
2	J	5	0	0	0	0
2	K	10	0	0	1	0
2	Ĺ	10	0	0	1	0
All	All	43895	0	44097	282	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 282 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:D:12:GLN:NE2	1:F:204:ASN:O	1.72	1.22
1:C:224:ASP:OD2	1:E:213:ASN:CB	1.94	1.15
1:C:224:ASP:OD2	1:E:213:ASN:HB3	1.48	1.11
1:F:16:GLN:CD	1:L:204:ASN:HD21	1.54	1.09
1:H:213:ASN:CB	1:I:224:ASP:OD2	2.03	1.07

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ (\rm \mathring{A}) \end{array}$	$\begin{array}{c} \text{Clash} \\ \text{overlap } (\text{\AA}) \end{array}$
1:C:175:ARG:NH2	1:I:26:ASP:OD2[1_556]	1.86	0.34
1:C:26:ASP:OD2	1:I:175:ARG:NH2[1_556]	1.92	0.28
1:B:164:ASP:OD2	1:J:111:ARG:CD[1_565]	2.07	0.13

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	433/461 (94%)	413 (95%)	20 (5%)	0	100	100
1	В	435/461 (94%)	415 (95%)	20 (5%)	0	100	100
1	C	436/461 (95%)	414 (95%)	22 (5%)	0	100	100
1	D	444/461 (96%)	424 (96%)	20 (4%)	0	100	100
1	E	443/461 (96%)	420 (95%)	23 (5%)	0	100	100
1	F	433/461 (94%)	413 (95%)	20 (5%)	0	100	100
1	G	433/461 (94%)	413 (95%)	20 (5%)	0	100	100
1	Н	432/461 (94%)	412 (95%)	20 (5%)	0	100	100
1	I	442/461 (96%)	420 (95%)	22 (5%)	0	100	100
1	J	439/461 (95%)	417 (95%)	22 (5%)	0	100	100
1	K	433/461 (94%)	413 (95%)	20 (5%)	0	100	100
1	L	437/461 (95%)	417 (95%)	20 (5%)	0	100	100
All	All	5240/5532 (95%)	4991 (95%)	249 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric Outliers		Percentiles		
1	A	395/412~(96%)	390 (99%)	5 (1%)	65 83		
1	В	$396/412\ (96\%)$	388 (98%)	8 (2%)	50 74		
1	С	$397/412\ (96\%)$	390 (98%)	7 (2%)	54 76		
1	D	$399/412\ (97\%)$	392 (98%)	7 (2%)	54 76		
1	E	$398/412\ (97\%)$	390 (98%)	8 (2%)	50 74		
1	F	$395/412\ (96\%)$	387 (98%)	8 (2%)	50 74		
1	G	$395/412\ (96\%)$	386 (98%)	9 (2%)	45 70		
1	Н	394/412 (96%)	386 (98%)	8 (2%)	50 74		
1	I	397/412 (96%)	389 (98%)	8 (2%)	50 74		



Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	J	399/412 (97%)	387 (97%)	12 (3%)	36 61		
1	K	395/412 (96%)	386 (98%)	9 (2%)	45 70		
1	L	398/412 (97%)	389 (98%)	9 (2%)	45 70		
All	All	4758/4944 (96%)	4660 (98%)	98 (2%)	48 72		

5 of 98 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	Н	359	LYS
1	J	8	ASN
1	Н	449	ARG
1	I	334	ARG
1	J	282	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 33 such sidechains are listed below:

Mol	Chain	Res	Type
1	K	343	GLN
1	L	138	HIS
1	L	343	GLN
1	Е	343	GLN
1	Е	138	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.



5.6 Ligand geometry (i)

24 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Type		Chain	Res	Link	Bond lengths			Bond angles		
IVIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	SO4	L	502	-	4,4,4	0.34	0	6,6,6	0.08	0
2	SO4	D	502	-	4,4,4	0.35	0	6,6,6	0.19	0
2	SO4	Е	501	-	4,4,4	0.48	0	6,6,6	0.43	0
2	SO4	A	501	-	4,4,4	0.44	0	6,6,6	0.34	0
2	SO4	G	503	-	4,4,4	0.33	0	6,6,6	0.13	0
2	SO4	В	501	-	4,4,4	0.37	0	6,6,6	0.30	0
2	SO4	L	501	-	4,4,4	0.40	0	6,6,6	0.15	0
2	SO4	I	501	-	4,4,4	0.44	0	6,6,6	0.16	0
2	SO4	С	502	ı	4,4,4	0.33	0	6,6,6	0.13	0
2	SO4	D	501	-	4,4,4	0.37	0	6,6,6	0.29	0
2	SO4	С	501	-	4,4,4	0.38	0	6,6,6	0.24	0
2	SO4	F	502	-	4,4,4	0.34	0	6,6,6	0.11	0
2	SO4	K	501	-	4,4,4	0.38	0	6,6,6	0.18	0
2	SO4	I	502	ı	4,4,4	0.32	0	6,6,6	0.10	0
2	SO4	В	502	-	4,4,4	0.32	0	6,6,6	0.11	0
2	SO4	G	502	-	4,4,4	0.41	0	6,6,6	0.24	0
2	SO4	K	502	ı	4,4,4	0.34	0	6,6,6	0.08	0
2	SO4	A	502	-	4,4,4	0.34	0	6,6,6	0.16	0
2	SO4	F	501	-	4,4,4	0.28	0	6,6,6	0.47	0
2	SO4	Н	501	-	4,4,4	0.50	0	6,6,6	0.21	0
2	SO4	Е	502	-	4,4,4	0.35	0	6,6,6	0.10	0
2	SO4	G	501	-	4,4,4	0.40	0	6,6,6	0.14	0
2	SO4	J	501	-	4,4,4	0.35	0	6,6,6	0.11	0
2	SO4	Н	502	-	4,4,4	0.35	0	6,6,6	0.15	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.




14 monomers are involved in 21 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	Е	501	SO4	2	0
2	A	501	SO4	3	0
2	В	501	SO4	3	0
2	L	501	SO4	1	0
2	D	501	SO4	1	0
2	С	501	SO4	1	0
2	F	502	SO4	1	0
2	K	501	SO4	1	0
2	G	502	SO4	2	0
2	A	502	SO4	2	0
2	F	501	SO4	1	0
2	Н	501	SO4	1	0
2	Е	502	SO4	1	0
2	Н	502	SO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.























































































































5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	# RSRZ > 2		$OWAB(\AA^2)$	Q<0.9	
1	A	437/461 (94%)	-0.05	2 (0%)	87	86	29, 63, 103, 136	0
1	В	439/461 (95%)	0.02	7 (1%)	70	67	33, 59, 100, 122	0
1	С	440/461 (95%)	0.06	6 (1%)	73	69	32, 63, 106, 133	0
1	D	446/461 (96%)	0.10	4 (0%)	81	79	36, 66, 107, 158	0
1	E	445/461 (96%)	0.12	10 (2%)	62	59	34, 65, 108, 147	0
1	F	437/461 (94%)	0.16	8 (1%)	67	64	30, 71, 117, 155	0
1	G	437/461 (94%)	0.12	4 (0%)	81	79	36, 67, 111, 147	0
1	Н	436/461 (94%)	0.07	2 (0%)	87	86	36, 65, 107, 135	0
1	I	444/461 (96%)	0.15	6 (1%)	73	69	34, 64, 104, 134	0
1	J	443/461 (96%)	0.36	14 (3%)	50	45	38, 74, 119, 158	0
1	K	437/461 (94%)	0.32	12 (2%)	56	52	41, 73, 120, 188	0
1	L	441/461 (95%)	0.44	16 (3%)	46	40	45, 85, 126, 160	0
All	All	5282/5532 (95%)	0.15	91 (1%)	69	65	29, 68, 114, 188	0

The worst 5 of 91 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	J	85	VAL	4.6
1	D	451	LEU	4.4
1	L	208	SER	4.2
1	I	451	LEU	4.1
1	J	53	ILE	4.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}(\mathring{\mathbf{A}}^2)$	Q < 0.9
2	SO4	F	502	5/5	0.67	0.13	101,102,110,117	0
2	SO4	Ε	502	5/5	0.69	0.13	106,114,122,128	0
2	SO4	J	501	5/5	0.71	0.12	119,128,130,131	0
2	SO4	L	502	5/5	0.71	0.10	106,127,133,140	0
2	SO4	I	502	5/5	0.72	0.12	91,93,105,106	0
2	SO4	D	502	5/5	0.77	0.11	72,98,106,116	0
2	SO4	K	502	5/5	0.77	0.11	109,114,119,135	0
2	SO4	A	502	5/5	0.77	0.13	90,102,112,115	0
2	SO4	С	502	5/5	0.78	0.10	85,89,105,112	0
2	SO4	В	502	5/5	0.78	0.10	89,93,96,105	0
2	SO4	Н	502	5/5	0.79	0.11	77,98,108,110	0
2	SO4	G	503	5/5	0.84	0.10	79,91,98,108	0
2	SO4	Ε	501	5/5	0.94	0.08	32,44,49,51	0
2	SO4	K	501	5/5	0.94	0.06	49,62,64,71	0
2	SO4	L	501	5/5	0.95	0.08	50,55,58,61	0
2	SO4	D	501	5/5	0.95	0.11	49,50,53,55	0
2	SO4	G	502	5/5	0.96	0.06	41,53,56,58	0
2	SO4	F	501	5/5	0.97	0.07	33,33,39,39	0
2	SO4	В	501	5/5	0.97	0.06	37,42,50,56	0
2	SO4	G	501	5/5	0.97	0.07	42,47,51,56	0
2	SO4	A	501	5/5	0.98	0.06	34,36,41,48	0
2	SO4	I	501	5/5	0.98	0.05	33,41,43,52	0
2	SO4	С	501	5/5	0.98	0.05	33,35,40,49	0
2	SO4	Н	501	5/5	0.98	0.05	38,42,45,49	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around SO4 F 502:













Electron density around SO4 E 502:





















Electron density around SO4 I 502:












Electron density around SO4 D 502:

 $2mF_o$ -DF_c (at 0.7 rmsd) in gray mF_o -DF_c (at 3 rmsd) in purple (negative) and green (positive)


























Electron density around SO4 H 502: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${ m mF}_o{ m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)




Electron density around SO4 G 503:














Electron density around SO4 K 501: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_{o}\text{-}\mathrm{DF}_{c}$ (at 3 rmsd) in purple (negative) and green (positive)




Electron density around SO4 L 501: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_{o}\text{-}\mathrm{DF}_{c}$ (at 3 rmsd) in purple (negative) and green (positive)
















Electron density around SO4 B 501:












Electron density around SO4 G 501:












Electron density around SO4 A 501:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around SO4 I 501:













Electron density around SO4 C 501:











6.5 Other polymers (i)

There are no such residues in this entry.

