



# wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 17, 2023 – 11:06 PM EDT

PDB ID : 4Z90  
Title : ELIC bound with the anesthetic isoflurane in the resting state  
Authors : Chen, Q.; Kinde, M.; Arjunan, P.; Cohen, A.; Xu, Y.; Tang, P.  
Deposited on : 2015-04-09  
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2092 (3.00-3.00)
Clashscore	141614	2416 (3.00-3.00)
Ramachandran outliers	138981	2333 (3.00-3.00)
Sidechain outliers	138945	2336 (3.00-3.00)
RSRZ outliers	127900	1990 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	322	 3% 77% 18% . .
1	B	322	 3% 73% 22% . .
1	C	322	 2% 74% 21% . .
1	D	322	 4% 75% 19% . .
1	E	322	 4% 75% 20% . .

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	322	
1	G	322	
1	H	322	
1	I	322	
1	J	322	

## 2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 25850 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Gamma-aminobutyric-acid receptor subunit beta-1.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	309	2527	1645	424	452	6	0	0	0
1	B	309	2527	1645	424	452	6	0	0	0
1	C	309	2527	1645	424	452	6	0	0	0
1	D	309	2527	1645	424	452	6	0	0	0
1	E	309	2527	1645	424	452	6	0	0	0
1	F	309	2527	1645	424	452	6	0	0	0
1	G	309	2527	1645	424	452	6	0	0	0
1	H	309	2527	1645	424	452	6	0	0	0
1	I	309	2527	1645	424	452	6	0	0	0
1	J	309	2527	1645	424	452	6	0	0	0

- Molecule 2 is (2R)-2-chloro-2-(difluoromethoxy)-1,1,1-trifluoroethane (three-letter code: 4LE) (formula: C<sub>3</sub>H<sub>2</sub>ClF<sub>5</sub>O).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	Cl	F	O		
2	A	1	Total	C	Cl	F	O	0	0
			10	3	1	5	1		
2	E	1	Total	C	Cl	F	O	0	0
			10	3	1	5	1		
2	F	1	Total	C	Cl	F	O	0	0
			10	3	1	5	1		
2	F	1	Total	C	Cl	F	O	0	0
			10	3	1	5	1		

- Molecule 3 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula:  $C_6H_{13}NO_4S$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
3	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
3	B	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
3	C	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
3	D	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
3	F	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
3	G	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
3	H	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
3	I	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
3	J	1	Total	C	N	O	S	0	0
			12	6	1	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	33	Total	O	0	0
			33	33		
4	B	45	Total	O	0	0
			45	45		
4	C	58	Total	O	0	0
			58	58		
4	D	30	Total	O	0	0
			30	30		
4	E	48	Total	O	0	0
			48	48		
4	F	33	Total	O	0	0
			33	33		
4	G	55	Total	O	0	0
			55	55		
4	H	42	Total	O	0	0
			42	42		
4	I	43	Total	O	0	0
			43	43		
4	J	33	Total	O	0	0
			33	33		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Gamma-aminobutyric-acid receptor subunit beta-1




- Molecule 1: Gamma-aminobutyric-acid receptor subunit beta-1





- Molecule 1: Gamma-aminobutyric-acid receptor subunit beta-1









- Molecule 1: Gamma-aminobutyric-acid receptor subunit beta-1




- Molecule 1: Gamma-aminobutyric-acid receptor subunit beta-1




- Molecule 1: Gamma-aminobutyric-acid receptor subunit beta-1




- Molecule 1: Gamma-aminobutyric-acid receptor subunit beta-1









- Molecule 1: Gamma-aminobutyric-acid receptor subunit beta-1




- Molecule 1: Gamma-aminobutyric-acid receptor subunit beta-1



- Molecule 1: Gamma-aminobutyric-acid receptor subunit beta-1





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	105.85Å 267.08Å 111.13Å 90.00° 106.94° 90.00°	Depositor
Resolution (Å)	36.97 – 3.00 39.63 – 3.00	Depositor EDS
% Data completeness (in resolution range)	99.8 (36.97-3.00) 99.8 (39.63-3.00)	Depositor EDS
$R_{merge}$	0.25	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.75 (at 3.01Å)	Xtrriage
Refinement program	PHENIX 1.9_1692	Depositor
R, $R_{free}$	0.209 , 0.243 0.210 , 0.241	Depositor DCC
$R_{free}$ test set	5886 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	92.7	Xtrriage
Anisotropy	0.343	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 61.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.024 for l,-k,h	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	25850	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	97.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.10% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MES, 4LE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.61	0/2595	0.60	0/3536
1	B	0.73	0/2595	0.68	0/3536
1	C	0.71	0/2595	0.65	0/3536
1	D	0.81	0/2595	0.68	0/3536
1	E	0.66	0/2595	0.62	0/3536
1	F	0.59	0/2595	0.61	0/3536
1	G	0.74	0/2595	0.66	1/3536 (0.0%)
1	H	0.64	0/2595	0.65	1/3536 (0.0%)
1	I	0.68	0/2595	0.63	0/3536
1	J	0.55	0/2595	0.58	0/3536
All	All	0.68	0/25950	0.64	2/35360 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	H	128	LEU	CA-CB-CG	6.07	129.27	115.30
1	G	240	LEU	CA-CB-CG	5.30	127.50	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2527	0	2503	43	0
1	B	2527	0	2503	60	0
1	C	2527	0	2503	58	0
1	D	2527	0	2503	56	1
1	E	2527	0	2503	45	0
1	F	2527	0	2503	53	0
1	G	2527	0	2503	49	1
1	H	2527	0	2503	68	0
1	I	2527	0	2503	59	0
1	J	2527	0	2503	68	0
2	A	10	0	1	0	0
2	E	10	0	1	0	0
2	F	20	0	2	2	0
3	A	24	0	24	0	0
3	B	12	0	12	0	0
3	C	12	0	12	1	0
3	D	12	0	12	0	0
3	F	12	0	12	0	0
3	G	12	0	12	1	0
3	H	12	0	12	1	0
3	I	12	0	12	0	0
3	J	12	0	12	1	0
4	A	33	0	0	8	0
4	B	45	0	0	9	0
4	C	58	0	0	11	0
4	D	30	0	0	8	0
4	E	48	0	0	10	0
4	F	33	0	0	5	0
4	G	55	0	0	12	0
4	H	42	0	0	5	0
4	I	43	0	0	8	0
4	J	33	0	0	5	0
All	All	25850	0	25154	511	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 511 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:123:ARG:CG	1:D:198:VAL:HG12	1.79	1.11
1:B:79:ILE:HD11	1:B:131:GLU:HB3	1.24	1.09

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:123:ARG:HG2	1:D:198:VAL:CG1	1.82	1.07
1:I:123:ARG:CG	1:I:198:VAL:HG12	1.83	1.07
1:A:123:ARG:HG2	1:A:198:VAL:CG1	1.85	1.06

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:13:ASP:OD2	1:G:174:ARG:NH2[2_556]	2.16	0.04

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	307/322 (95%)	283 (92%)	22 (7%)	2 (1%)	22	60
1	B	307/322 (95%)	282 (92%)	23 (8%)	2 (1%)	22	60
1	C	307/322 (95%)	282 (92%)	24 (8%)	1 (0%)	41	76
1	D	307/322 (95%)	284 (92%)	22 (7%)	1 (0%)	41	76
1	E	307/322 (95%)	281 (92%)	24 (8%)	2 (1%)	22	60
1	F	307/322 (95%)	283 (92%)	22 (7%)	2 (1%)	22	60
1	G	307/322 (95%)	282 (92%)	23 (8%)	2 (1%)	22	60
1	H	307/322 (95%)	282 (92%)	24 (8%)	1 (0%)	41	76
1	I	307/322 (95%)	286 (93%)	19 (6%)	2 (1%)	22	60
1	J	307/322 (95%)	282 (92%)	24 (8%)	1 (0%)	41	76
All	All	3070/3220 (95%)	2827 (92%)	227 (7%)	16 (0%)	29	68

5 of 16 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	I	182	GLN
1	C	182	GLN
1	E	182	GLN
1	F	182	GLN
1	H	182	GLN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	277/284 (98%)	275 (99%)	2 (1%)	84	94
1	B	277/284 (98%)	274 (99%)	3 (1%)	73	90
1	C	277/284 (98%)	275 (99%)	2 (1%)	84	94
1	D	277/284 (98%)	273 (99%)	4 (1%)	67	88
1	E	277/284 (98%)	270 (98%)	7 (2%)	47	79
1	F	277/284 (98%)	275 (99%)	2 (1%)	84	94
1	G	277/284 (98%)	276 (100%)	1 (0%)	91	97
1	H	277/284 (98%)	271 (98%)	6 (2%)	52	81
1	I	277/284 (98%)	274 (99%)	3 (1%)	73	90
1	J	277/284 (98%)	275 (99%)	2 (1%)	84	94
All	All	2770/2840 (98%)	2738 (99%)	32 (1%)	71	90

5 of 32 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	I	302	LEU
1	I	308	PHE
1	E	29	LEU
1	E	14	VAL
1	J	247	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 28 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	251	ASN
1	J	251	ASN
1	F	251	ASN
1	I	169	HIS
1	F	233	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

14 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	MES	C	401	-	12,12,12	2.26	1 (8%)	14,16,16	2.68	7 (50%)
3	MES	D	401	-	12,12,12	2.13	1 (8%)	14,16,16	2.58	6 (42%)
3	MES	I	401	-	12,12,12	2.20	1 (8%)	14,16,16	2.17	6 (42%)
3	MES	B	401	-	12,12,12	2.37	1 (8%)	14,16,16	2.25	5 (35%)
3	MES	J	401	-	12,12,12	2.25	1 (8%)	14,16,16	2.28	7 (50%)
3	MES	F	403	-	12,12,12	2.21	1 (8%)	14,16,16	2.54	7 (50%)
2	4LE	F	401	-	6,9,9	0.38	0	7,13,13	1.43	1 (14%)
3	MES	A	402	-	12,12,12	2.33	1 (8%)	14,16,16	2.53	7 (50%)



Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	MES	A	403	-	12,12,12	2.32	1 (8%)	14,16,16	1.97	6 (42%)
3	MES	G	401	-	12,12,12	2.26	1 (8%)	14,16,16	2.30	7 (50%)
2	4LE	F	402	-	6,9,9	0.50	0	7,13,13	1.35	0
2	4LE	A	401	-	6,9,9	0.67	0	7,13,13	1.73	2 (28%)
2	4LE	E	401	-	6,9,9	0.39	0	7,13,13	1.46	2 (28%)
3	MES	H	401	-	12,12,12	2.28	1 (8%)	14,16,16	2.58	7 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MES	C	401	-	-	5/6/14/14	0/1/1/1
3	MES	D	401	-	-	1/6/14/14	0/1/1/1
3	MES	I	401	-	-	2/6/14/14	0/1/1/1
3	MES	B	401	-	-	4/6/14/14	0/1/1/1
3	MES	J	401	-	-	5/6/14/14	0/1/1/1
3	MES	F	403	-	-	1/6/14/14	0/1/1/1
2	4LE	F	401	-	-	5/8/10/10	-
3	MES	A	402	-	-	3/6/14/14	0/1/1/1
3	MES	A	403	-	-	3/6/14/14	0/1/1/1
3	MES	G	401	-	-	3/6/14/14	0/1/1/1
2	4LE	F	402	-	-	4/8/10/10	-
2	4LE	A	401	-	-	3/8/10/10	-
2	4LE	E	401	-	-	3/8/10/10	-
3	MES	H	401	-	-	4/6/14/14	0/1/1/1

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	401	MES	C8-S	-7.99	1.66	1.77
3	A	403	MES	C8-S	-7.81	1.66	1.77
3	A	402	MES	C8-S	-7.80	1.66	1.77
3	H	401	MES	C8-S	-7.64	1.66	1.77
3	C	401	MES	C8-S	-7.63	1.66	1.77

The worst 5 of 70 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	402	MES	C5-N4-C3	5.33	120.84	108.83
3	H	401	MES	C5-N4-C3	5.26	120.67	108.83
3	C	401	MES	C5-N4-C3	5.16	120.45	108.83
3	F	403	MES	C5-N4-C3	5.10	120.31	108.83
3	D	401	MES	C5-N4-C3	5.09	120.28	108.83

There are no chirality outliers.

5 of 46 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401	4LE	F01-C01-C02-O01
2	A	401	4LE	F02-C01-C02-O01
2	A	401	4LE	F03-C01-C02-O01
2	E	401	4LE	F01-C01-C02-O01
2	E	401	4LE	F02-C01-C02-O01

There are no ring outliers.

6 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	401	MES	1	0
3	J	401	MES	1	0
2	F	401	4LE	1	0
3	G	401	MES	1	0
2	F	402	4LE	1	0
3	H	401	MES	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	309/322 (95%)	-0.22	9 (2%) 51 23	62, 93, 151, 241	0
1	B	309/322 (95%)	-0.26	10 (3%) 47 20	55, 80, 175, 227	0
1	C	309/322 (95%)	-0.26	7 (2%) 60 31	53, 83, 179, 228	0
1	D	309/322 (95%)	-0.11	13 (4%) 36 14	51, 78, 171, 249	0
1	E	309/322 (95%)	-0.14	12 (3%) 39 15	58, 89, 169, 236	0
1	F	309/322 (95%)	-0.23	12 (3%) 39 15	59, 98, 166, 264	0
1	G	309/322 (95%)	-0.25	8 (2%) 56 27	54, 81, 149, 231	0
1	H	309/322 (95%)	-0.11	17 (5%) 25 9	59, 88, 175, 256	0
1	I	309/322 (95%)	-0.16	14 (4%) 33 12	58, 86, 175, 236	0
1	J	309/322 (95%)	-0.05	15 (4%) 29 11	63, 98, 190, 244	0
All	All	3090/3220 (95%)	-0.18	117 (3%) 40 16	51, 88, 174, 264	0

The worst 5 of 117 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	180	SER	9.8
1	D	181	VAL	8.2
1	D	179	SER	7.9
1	I	179	SER	6.6
1	B	290	GLY	6.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	MES	C	401	12/12	0.80	0.19	109,123,138,142	0
3	MES	I	401	12/12	0.81	0.26	125,131,157,160	0
3	MES	J	401	12/12	0.83	0.21	146,154,166,168	0
3	MES	F	403	12/12	0.84	0.15	117,127,146,150	0
3	MES	A	402	12/12	0.85	0.18	106,120,174,174	0
3	MES	A	403	12/12	0.85	0.15	127,135,140,142	0
3	MES	D	401	12/12	0.86	0.22	115,125,128,128	0
3	MES	H	401	12/12	0.87	0.17	116,122,150,150	0
3	MES	B	401	12/12	0.93	0.09	108,116,132,136	0
3	MES	G	401	12/12	0.94	0.21	108,117,138,140	0
2	4LE	E	401	10/10	0.95	0.42	109,119,146,148	0
2	4LE	A	401	10/10	0.97	0.30	116,119,131,140	0
2	4LE	F	402	10/10	0.97	0.30	109,117,122,133	0
2	4LE	F	401	10/10	0.98	0.36	119,120,127,146	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.