



# Full wwPDB X-ray Structure Validation Report ⓘ

Jul 25, 2022 – 04:14 pm BST

PDB ID : 7ZCQ  
Title : Nitrite-bound MSOX movie series dataset 25 (20 MGy) of the copper nitrite reductase from Bradyrhizobium sp. ORS 375 (two-domain) - NO/water intermediate  
Authors : Rose, S.L.; Baba, S.; Okumura, H.; Antonyuk, S.V.; Sasaki, D.; Tosha, T.; Kumasaka, T.; Eady, R.R.; Yamamoto, M.; Hasnain, S.S.  
Deposited on : 2022-03-28  
Resolution : 1.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at <http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 2.29  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0267  
CCP4 : 7.1.010 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.29


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*




The reported resolution of this entry is 1.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1509 (1.38-1.34)
Clashscore	141614	1551 (1.38-1.34)
Ramachandran outliers	138981	1530 (1.38-1.34)
Sidechain outliers	138945	1530 (1.38-1.34)
RSRZ outliers	127900	1487 (1.38-1.34)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	350	 90% 7% . .
2	E	2	 100%
2	F	2	 100%

## 2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 3531 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Copper-containing nitrite reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	341	2824	1819	482	506	17	1	40	0

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	initiating methionine	UNP H0SLX7
A	342	GLU	-	expression tag	UNP H0SLX7
A	343	ASN	-	expression tag	UNP H0SLX7
A	344	LEU	-	expression tag	UNP H0SLX7
A	345	TYR	-	expression tag	UNP H0SLX7
A	346	PHE	-	expression tag	UNP H0SLX7
A	347	GLN	-	expression tag	UNP H0SLX7
A	348	GLY	-	expression tag	UNP H0SLX7
A	349	GLY	-	expression tag	UNP H0SLX7
A	350	SER	-	expression tag	UNP H0SLX7

- Molecule 2 is an oligosaccharide called beta-D-fructofuranose-(2-1)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
			Total	C	O			
2	E	2	23	12	11	0	2	0
2	F	2	23	12	11	0	2	0

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	1
3	A	1	Total O S 5 4 1	0	1
3	A	1	Total O S 5 4 1	0	1
3	A	1	Total O S 5 4 1	0	1
3	A	1	Total O S 10 8 2	0	1
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	1
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	1

- Molecule 4 is COPPER (II) ION (three-letter code: CU) (formula: Cu) (labeled as "Ligand of Interest" by depositor).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	Cu	0	0
			2	2		

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	1
			6	3	3		
5	A	1	Total	C	O	2	0
			6	3	3		
5	A	1	Total	C	O	0	1
			12	6	6		

- Molecule 6 is NITRIC OXIDE (three-letter code: NO) (formula: NO) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	N	O		
6	A	1	2	1	1	0	1

- Molecule 7 is NITRITE ION (three-letter code: NO<sub>2</sub>) (formula: NO<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	N	O		
7	A	1	3	1	2	0	0
7	A	1	3	1	2	0	1
7	A	1	3	1	2	0	0


- Molecule 8 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	531	Total O 564 564	0	117


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Copper-containing nitrite reductase



- Molecule 2: beta-D-fructofuranose-(2-1)-alpha-D-glucopyranose



- Molecule 2: beta-D-fructofuranose-(2-1)-alpha-D-glucopyranose





## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 3	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	107.04Å 107.04Å 107.04Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	75.69 – 1.35 75.69 – 1.35	Depositor EDS
% Data completeness (in resolution range)	96.6 (75.69-1.35) 96.6 (75.69-1.35)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.00 (at 1.35Å)	Xtrriage
Refinement program	REFMAC 5.8.0267	Depositor
R, $R_{free}$	0.131 , 0.157 0.132 , 0.158	Depositor DCC
$R_{free}$ test set	4412 reflections (5.10%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	16.4	Xtrriage
Anisotropy	0.000	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	(Not available) , (Not available)	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.036 for l,-k,h	Xtrriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	3531	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	22.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.46% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NO2, GOL, GLC, CU, NO, FRU, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.89	3/3006 (0.1%)	0.99	8/4080 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	270[B]	GLU	CD-OE1	12.02	1.38	1.25
1	A	202	GLU	CD-OE1	5.92	1.32	1.25
1	A	318	GLU	CD-OE2	5.15	1.31	1.25

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	154[A]	ARG	CG-CD-NE	-8.61	93.71	111.80
1	A	154[B]	ARG	CG-CD-NE	-8.61	93.71	111.80
1	A	117	ARG	NE-CZ-NH1	-8.03	116.29	120.30
1	A	279[A]	ARG	CB-CA-C	6.12	122.64	110.40
1	A	279[B]	ARG	CB-CA-C	6.12	122.64	110.40
1	A	154[A]	ARG	NE-CZ-NH1	-5.85	117.37	120.30
1	A	154[B]	ARG	NE-CZ-NH1	-5.85	117.37	120.30
1	A	92	ASP	CB-CG-OD1	5.20	122.98	118.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	299	VAL	Peptide
1	A	301	HIS	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2824	0	2863	19	0
2	E	23	0	21	0	0
2	F	23	0	21	5	0
3	A	60	0	0	0	0
4	A	2	0	0	0	0
5	A	24	0	29	0	0
6	A	2	0	0	1	0
7	A	9	0	0	0	0
8	A	564	0	0	4	0
All	All	3531	0	2934	24	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:F:1[A]:GLC:H3	2:F:2[A]:FRU:H11	1.46	0.95
1:A:154[B]:ARG:HD2	8:A:797:HOH:O	1.83	0.77
2:F:1[A]:GLC:C3	2:F:2[A]:FRU:H11	2.20	0.68
1:A:252:ILE:HD12	1:A:297:ALA:HB3	1.82	0.61
1:A:87[C]:MET:CE	1:A:140:HIS:CE1	2.84	0.59
1:A:92:ASP:OD2	6:A:410[A]:NO:O	2.21	0.58
1:A:28:GLN:O	1:A:154[B]:ARG:NH1	2.32	0.57
2:F:1[A]:GLC:C4	2:F:2[A]:FRU:O2	2.49	0.50
1:A:87[C]:MET:HE3	1:A:140:HIS:NE2	2.27	0.49
1:A:28:GLN:OE1	1:A:31:LYS:HE3	2.13	0.49
1:A:339:SER:HB3	1:A:341[A]:ASN:HD21	1.78	0.48

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:87[C]:MET:CE	1:A:140:HIS:NE2	2.76	0.48
2:F:2[A]:FRU:H12	2:F:2[A]:FRU:H5	1.36	0.47
1:A:53[A]:PHE:CZ	1:A:178:LEU:HD22	2.51	0.46
1:A:341[A]:ASN:HB2	8:A:541:HOH:O	2.16	0.45
1:A:207[B]:LYS:HD3	8:A:920:HOH:O	2.16	0.45
1:A:87[C]:MET:HE3	1:A:140:HIS:CE1	2.52	0.44
1:A:341[B]:ASN:HD22	1:A:341[B]:ASN:HA	1.63	0.43
2:F:1[A]:GLC:H3	2:F:2[A]:FRU:C1	2.34	0.42
1:A:53[A]:PHE:HD2	8:A:676:HOH:O	2.01	0.42
1:A:67[B]:MET:HB2	1:A:149[B]:VAL:HG22	2.01	0.42
1:A:28:GLN:OE1	1:A:31:LYS:CE	2.68	0.41
1:A:169[A]:ARG:HD3	1:A:171:TYR:OH	2.20	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	377/350 (108%)	369 (98%)	8 (2%)	0	<a href="#">100</a> <a href="#">100</a>

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	312/284 (110%)	307 (98%)	5 (2%)	62 30

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	53[A]	PHE
1	A	53[B]	PHE
1	A	248	ARG
1	A	261[A]	GLU
1	A	261[B]	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	GLC	E	1[A]	2	11,11,12	2.36	5 (45%)	15,15,17	3.22	9 (60%)
2	FRU	E	2[A]	2	11,12,12	1.38	1 (9%)	10,18,18	1.95	1 (10%)
2	GLC	F	1[A]	2	11,11,12	2.25	4 (36%)	15,15,17	3.91	9 (60%)
2	FRU	F	2[A]	2	11,12,12	3.06	4 (36%)	10,18,18	4.30	6 (60%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	E	1[A]	2	-	0/2/19/22	0/1/1/1
2	FRU	E	2[A]	2	-	0/5/24/24	0/1/1/1
2	GLC	F	1[A]	2	-	1/2/19/22	0/1/1/1
2	FRU	F	2[A]	2	-	5/5/24/24	0/1/1/1

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	2[A]	FRU	O3-C3	7.97	1.58	1.42
2	F	1[A]	GLC	C4-C5	5.49	1.64	1.53
2	E	1[A]	GLC	O3-C3	5.05	1.54	1.43
2	F	2[A]	FRU	O4-C4	4.26	1.53	1.43
2	E	2[A]	FRU	O2-C2	3.48	1.46	1.40
2	E	1[A]	GLC	C2-C3	3.41	1.57	1.52
2	F	2[A]	FRU	C4-C3	2.83	1.64	1.52
2	F	1[A]	GLC	O3-C3	2.77	1.49	1.43
2	E	1[A]	GLC	C4-C3	2.50	1.58	1.52
2	F	1[A]	GLC	O4-C4	2.35	1.48	1.43
2	F	1[A]	GLC	C4-C3	2.14	1.57	1.52
2	E	1[A]	GLC	O2-C2	2.01	1.47	1.43
2	F	2[A]	FRU	O2-C2	-2.00	1.37	1.40
2	E	1[A]	GLC	O5-C5	2.00	1.47	1.43

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	1[A]	GLC	C1-C2-C3	8.82	120.50	109.67
2	F	1[A]	GLC	C3-C4-C5	7.51	123.64	110.24
2	F	2[A]	FRU	O3-C3-C4	6.63	136.24	113.32
2	F	2[A]	FRU	O4-C4-C3	6.61	131.94	112.15
2	F	2[A]	FRU	O4-C4-C5	-6.44	92.44	111.05
2	F	1[A]	GLC	O3-C3-C2	6.12	121.70	109.99
2	F	1[A]	GLC	O4-C4-C3	-5.38	97.90	110.35
2	F	1[A]	GLC	O5-C5-C6	-5.37	98.78	107.20
2	F	1[A]	GLC	C1-C2-C3	-5.10	103.39	109.67
2	F	2[A]	FRU	O5-C5-C6	4.81	122.23	108.85
2	E	2[A]	FRU	O2-C2-O5	4.68	118.53	109.50
2	F	2[A]	FRU	O2-C2-O5	4.48	118.16	109.50

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	1[A]	GLC	C1-O5-C5	-3.60	107.32	112.19
2	F	1[A]	GLC	O6-C6-C5	-3.57	99.05	111.29
2	E	1[A]	GLC	O3-C3-C4	3.32	118.04	110.35
2	E	1[A]	GLC	C3-C4-C5	3.32	116.16	110.24
2	F	1[A]	GLC	C1-O5-C5	-3.29	107.73	112.19
2	F	1[A]	GLC	C6-C5-C4	3.04	120.13	113.00
2	E	1[A]	GLC	O5-C5-C4	2.87	117.81	110.83
2	E	1[A]	GLC	O4-C4-C5	-2.79	102.38	109.30
2	E	1[A]	GLC	C2-C3-C4	-2.66	106.29	110.89
2	F	2[A]	FRU	O1-C1-C2	2.66	117.51	111.86
2	F	1[A]	GLC	O2-C2-C3	2.63	115.40	110.14
2	E	1[A]	GLC	O2-C2-C1	-2.16	104.73	109.15
2	E	1[A]	GLC	O4-C4-C3	2.15	115.33	110.35

There are no chirality outliers.

All (6) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	F	2[A]	FRU	O1-C1-C2-C3
2	F	2[A]	FRU	O1-C1-C2-O2
2	F	2[A]	FRU	O1-C1-C2-O5
2	F	2[A]	FRU	C4-C5-C6-O6
2	F	1[A]	GLC	O5-C5-C6-O6
2	F	2[A]	FRU	O5-C5-C6-O6

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	F	1[A]	GLC	4	0
2	F	2[A]	FRU	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





## 5.6 Ligand geometry

Of 22 ligands modelled in this entry, 2 are monoatomic - leaving 20 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	SO4	A	416	-	4,4,4	0.14	0	6,6,6	0.18	0
6	NO	A	410[A]	4	0,1,1	-	-	-	-	-
3	SO4	A	402	-	4,4,4	0.58	0	6,6,6	0.32	0
7	NO2	A	418	-	1,2,2	2.62	1 (100%)	0,1,1	-	-
3	SO4	A	401	-	4,4,4	0.37	0	6,6,6	0.49	0
3	SO4	A	411[A]	-	4,4,4	0.32	0	6,6,6	0.21	0
5	GOL	A	408	-	5,5,5	3.08	1 (20%)	5,5,5	4.31	2 (40%)
7	NO2	A	420	-	1,2,2	1.38	0	0,1,1	-	-
5	GOL	A	409[A]	-	5,5,5	0.20	0	5,5,5	0.41	0
3	SO4	A	417[B]	-	4,4,4	0.13	0	6,6,6	0.14	0
5	GOL	A	409[B]	-	5,5,5	0.28	0	5,5,5	0.43	0
5	GOL	A	407[A]	-	5,5,5	0.42	0	5,5,5	0.61	0
3	SO4	A	403[A]	-	4,4,4	0.35	0	6,6,6	0.21	0
3	SO4	A	415[A]	-	4,4,4	0.54	0	6,6,6	0.51	0
3	SO4	A	413[A]	-	4,4,4	0.31	0	6,6,6	0.28	0
3	SO4	A	413[B]	-	4,4,4	0.30	0	6,6,6	0.13	0
7	NO2	A	419[A]	-	1,2,2	1.38	0	0,1,1	-	-
3	SO4	A	412[A]	-	4,4,4	0.35	0	6,6,6	0.40	0
3	SO4	A	404[A]	-	4,4,4	0.29	0	6,6,6	0.38	0
3	SO4	A	414	-	4,4,4	0.51	0	6,6,6	0.23	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	A	409[B]	-	-	3/4/4/4	-
5	GOL	A	407[A]	-	-	4/4/4/4	-
5	GOL	A	408	-	-	1/4/4/4	-
5	GOL	A	409[A]	-	-	0/4/4/4	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	408	GOL	C3-C2	-6.82	1.23	1.51
7	A	418	NO2	O1-N	2.62	1.35	1.22

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	408	GOL	O2-C2-C3	8.06	144.63	109.12
5	A	408	GOL	C3-C2-C1	-4.85	92.85	111.70

There are no chirality outliers.

All (8) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	A	407[A]	GOL	O1-C1-C2-O2
5	A	407[A]	GOL	O1-C1-C2-C3
5	A	407[A]	GOL	C1-C2-C3-O3
5	A	408	GOL	O1-C1-C2-C3
5	A	409[B]	GOL	C1-C2-C3-O3
5	A	407[A]	GOL	O2-C2-C3-O3
5	A	409[B]	GOL	O2-C2-C3-O3
5	A	409[B]	GOL	O1-C1-C2-C3

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	410[A]	NO	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	341/350 (97%)	-0.29	3 (0%) 84 87	12, 16, 34, 52	17 (4%)

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	2	ASP	8.2
1	A	338	GLY	2.6
1	A	339	SER	2.6

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	GLC	F	1[A]	11/12	0.73	0.23	26,43,48,51	11
2	FRU	E	2[A]	12/12	0.85	0.25	38,46,51,52	12
2	FRU	F	2[A]	12/12	0.86	0.15	21,39,44,46	12
2	GLC	E	1[A]	11/12	0.92	0.14	19,33,43,44	11

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around Chain E:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	SO4	A	416	5/5	0.79	0.27	51,52,56,59	5
5	GOL	A	407[A]	6/6	0.81	0.27	37,44,47,48	6
5	GOL	A	409[A]	6/6	0.87	0.11	18,19,22,22	6
5	GOL	A	409[B]	6/6	0.87	0.11	32,38,40,40	6
3	SO4	A	417[B]	5/5	0.88	0.23	35,42,50,50	5
5	GOL	A	408	6/6	0.88	0.16	32,41,48,51	6
6	NO	A	410[A]	2/2	0.88	0.23	21,21,21,24	2
7	NO2	A	418	3/3	0.90	0.09	37,37,45,47	0
3	SO4	A	401	5/5	0.91	0.21	42,48,51,51	5
3	SO4	A	403[A]	5/5	0.91	0.19	37,46,50,58	5
3	SO4	A	404[A]	5/5	0.91	0.15	40,46,50,58	5

*Continued on next page...*

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	SO4	A	412[A]	5/5	0.91	0.17	41,46,47,53	5
3	SO4	A	415[A]	5/5	0.95	0.10	19,29,38,39	5
7	NO2	A	419[A]	3/3	0.95	0.15	26,26,28,44	3
3	SO4	A	413[B]	5/5	0.96	0.17	38,48,49,50	5
3	SO4	A	414	5/5	0.96	0.14	40,41,46,47	5
3	SO4	A	413[A]	5/5	0.96	0.17	13,16,18,20	5
7	NO2	A	420	3/3	0.96	0.07	25,25,26,28	3
3	SO4	A	402	5/5	0.97	0.16	41,42,48,51	5
3	SO4	A	411[A]	5/5	0.98	0.13	35,44,45,47	5
4	CU	A	405	1/1	1.00	0.07	16,16,16,16	0
4	CU	A	406	1/1	1.00	0.07	15,15,15,15	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around CU A 405:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







## 6.5 Other polymers [i](#)

There are no such residues in this entry.