

Full wwPDB X-ray Structure Validation Report (i)

Oct 11, 2023 – 10:12 am BST

PDB ID : 7ZLU

Title: Catalytic domain of UDP-Glucose Glycoprotein Glucosyltransferase from

Chaetomium thermophilum in complex with UDP-2-deoxy-2-fluoro-D-glu

cose

Authors : Roversi, P.; Zitzmann, N.; Bayo, Y.; Ibba, R.

Deposited on : 2022-04-15

Resolution : 2.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.4, CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.35.1

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# \mathrm{Entries})$	$(\# ext{Entries}, ext{ resolution range}(ext{Å}))$
R_{free}	130704	1692 (2.04-2.04)
Clashscore	141614	1773 (2.04-2.04)
Ramachandran outliers	138981	1752 (2.04-2.04)
Sidechain outliers	138945	1752 (2.04-2.04)
RSRZ outliers	127900	1672 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	297	87%	6%	6%
1	В	297	82%	11%	6%
1	С	297	87%	7%	6%



2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 7582 atoms, of which 32 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called UDP-glucose-glycoprotein glucosyltransferase-like protein.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	Λ	278	Total	С	C N O	О	S	0	2	0
1	1 A	210	2354	1520	401	418	15	U	3	
1	D	278	Total	С	N	О	S	0	4	0
1	Ъ	210	2360	1524	400	421	15	0	4	
1	С	278	Total	С	N	О	S	0	6	0
1		210	2371	1530	401	425	15	U	U	

There are 30 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1184	GLU	-	expression tag	UNP G0SB58
A	1185	THR	-	expression tag	UNP G0SB58
A	1186	GLY	-	expression tag	UNP G0SB58
A	1474	GLY	-	expression tag	UNP G0SB58
A	1475	THR	-	expression tag	UNP G0SB58
A	1476	LYS	-	expression tag	UNP G0SB58
A	1477	HIS	-	expression tag	UNP G0SB58
A	1478	HIS	-	expression tag	UNP G0SB58
A	1479	HIS	-	expression tag	UNP G0SB58
A	1480	HIS	-	expression tag	UNP G0SB58
В	1184	GLU	-	expression tag	UNP G0SB58
В	1185	THR	-	expression tag	UNP G0SB58
В	1186	GLY	-	expression tag	UNP G0SB58
В	1474	GLY	-	expression tag	UNP G0SB58
В	1475	THR	-	expression tag	UNP G0SB58
В	1476	LYS	-	expression tag	UNP G0SB58
В	1477	HIS	-	expression tag	UNP G0SB58
В	1478	HIS	-	expression tag	UNP G0SB58
В	1479	HIS	-	expression tag	UNP G0SB58
В	1480	HIS	-	expression tag	UNP G0SB58
С	1184	GLU	-	expression tag	UNP G0SB58
С	1185	THR	-	expression tag	UNP G0SB58
С	1186	GLY	-	expression tag	UNP G0SB58



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
С	1474	GLY	-	expression tag	UNP G0SB58
С	1475	THR	-	expression tag	UNP G0SB58
С	1476	LYS	-	expression tag	UNP G0SB58
С	1477	HIS	-	expression tag	UNP G0SB58
С	1478	HIS	-	expression tag	UNP G0SB58
С	1479	HIS	-	expression tag	UNP G0SB58
С	1480	HIS	-	expression tag	UNP G0SB58


 \bullet Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $\rm C_8H_{15}NO_6).$



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C N O 14 8 1 5	0	0
2	В	1	Total C N O 14 8 1 5	0	0
2	С	1	Total C N O 14 8 1 5	0	0

• Molecule 3 is URIDINE-5'-DIPHOSPHATE-2-DEOXY-2-FLUORO-ALPHA-D-GLUCOS E (three-letter code: U2F) (formula: C₁₅H₂₃FN₂O₁₆P₂) (labeled as "Ligand of Interest" by depositor).






Mol	Chain	Residues	Atoms				ZeroOcc	AltConf			
3	Λ	1	Total	С	F	N	О	Р	0	0	
9	Λ	1	36	15	1	2	16	2	0		
3	В	1	Total	С	F	N	О	Р	0	0	
9	D	1	36	15	1	2	16	2	0	0	
3	С	1	Total	С	F	N	О	Р	0	0	
3		1	36	15	1	2	16	2			

• Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Ca 1 1	0	0
4	В	1	Total Ca 1 1	0	0
4	С	1	Total Ca 1 1	0	0

• Molecule 5 is 1,3-PROPANDIOL (three-letter code: PDO) (formula: $C_3H_8O_2$).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C H O 13 3 8 2	8	0
5	В	1	Total C H O 26 6 16 4	16	1
5	С	1	Total C H O 13 3 8 2	8	0

• Molecule 6 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	100	Total O 100 100	0	0
6	В	97	Total O 97 97	0	0
6	С	95	Total O 95 95	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: UDP-glucose-glycoprotein glucosyltransferase-like protein



• Molecule 1: UDP-glucose-glycoprotein glucosyltransferase-like protein



• Molecule 1: UDP-glucose-glycoprotein glucosyltransferase-like protein





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1	Depositor
Cell constants	68.16Å 72.32Å 72.39Å	Donogitor
a, b, c, α , β , γ	110.48° 108.33° 108.26°	Depositor
Resolution (Å)	41.16 - 2.05	Depositor
Resolution (A)	41.16 - 2.05	EDS
% Data completeness	57.7 (41.16-2.05)	Depositor
(in resolution range)	57.3 (41.16-2.05)	EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.41 (at 2.05Å)	Xtriage
Refinement program	BUSTER 2.10.4 (3-FEB-2022)	Depositor
R, R_{free}	0.226 , 0.270	Depositor
it, it free	0.220 , 0.266	DCC
R_{free} test set	1863 reflections (4.75%)	wwPDB-VP
Wilson B-factor (Å ²)	31.6	Xtriage
Anisotropy	0.035	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.33, 27.5	EDS
L-test for twinning ²	$< L > = 0.51, < L^2> = 0.34$	Xtriage
	0.390 for h,l,-h-k-l	
	0.390 for h,-h-k-l,k	
Estimated twinning fraction	0.000 for -h,-l,-k	Xtriage
	0.000 for -h,-k,h+k+l	
	0.000 for -h,h+k+l,-l	
F_o, F_c correlation	0.94	EDS
Total number of atoms	7582	wwPDB-VP
Average B, all atoms (\mathring{A}^2)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.95% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, U2F, PDO, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
IVIOI		RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.42	0/2437	0.58	0/3308	
1	В	0.44	0/2446	0.61	0/3320	
1	С	0.44	0/2460	0.60	0/3339	
All	All	0.43	0/7343	0.60	0/9967	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2354	0	2268	11	0
1	В	2360	0	2272	18	0
1	С	2371	0	2280	12	0
2	A	14	0	13	0	0
2	В	14	0	13	0	0
2	С	14	0	13	0	0
3	A	36	0	21	0	0
3	В	36	0	21	0	0
3	С	36	0	21	0	0
4	A	1	0	0	0	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	В	1	0	0	0	0
4	С	1	0	0	0	0
5	A	5	8	8	0	0
5	В	10	16	16	2	0
5	С	5	8	8	2	0
6	A	100	0	0	1	0
6	В	97	0	0	0	0
6	С	95	0	0	0	0
All	All	7550	32	6954	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:1342:TRP:HA	1:C:1347:TRP:HB2	1.79	0.64
1:B:1342:TRP:HA	1:B:1347:TRP:HB2	1.83	0.61
1:C:1335:GLU:OE2	5:C:1504:PDO:H31	2.03	0.58
1:A:1381:GLN:HG3	1:A:1403:MET:SD	2.44	0.57
1:C:1332:SER:HB2	1:C:1424:SER:HB3	1.88	0.55
1:A:1295:LEU:O	1:A:1370:ARG:NH1	2.40	0.54
1:B:1333:ARG:HA	5:B:1504[A]:PDO:H32	1.88	0.54
1:A:1197:HIS:HE1	1:A:1294:SER:OG	1.90	0.53
1:B:1415[B]:GLU:H	1:B:1415[B]:GLU:CD	2.14	0.50
1:B:1299:ILE:HD13	1:B:1314:LEU:HD23	1.93	0.49
1:B:1332:SER:HB2	1:B:1424:SER:HB3	1.95	0.49
1:B:1223:MET:HG3	1:B:1254:TYR:HB3	1.94	0.48
1:B:1381:GLN:HG3	1:B:1403:MET:SD	2.53	0.48
1:A:1342:TRP:HA	1:A:1347:TRP:HB2	1.95	0.48
1:B:1295:LEU:O	1:B:1370:ARG:NH1	2.47	0.47
1:B:1385:LEU:HD11	1:B:1402:HIS:CD2	2.49	0.47
1:C:1302:ASP:HB3	1:C:1305:GLN:HG3	1.97	0.47
1:C:1381:GLN:HG3	1:C:1403:MET:SD	2.55	0.46
1:B:1333:ARG:HG3	5:B:1504[A]:PDO:H22	1.97	0.46
1:A:1426:GLU:N	1:A:1426:GLU:CD	2.70	0.45
1:B:1302:ASP:HB3	1:B:1305:GLN:HG3	1.97	0.45
1:C:1454:VAL:HB	1:C:1457:TRP:HB2	1.98	0.45
1:B:1245:ASP:O	1:B:1248:PRO:HD2	2.17	0.44
1:B:1351:LEU:O	1:B:1354:LYS:HB2	2.18	0.44
1:B:1247:ILE:HG23	1:B:1248:PRO:HD3	1.99	0.44



Continued from previous page...

Atom-1	Atom-2	Interatomic	Clash	
Atom-1	Atom-2	${ m distance} ({ m \AA})$	overlap (Å)	
1:B:1328:PRO:HA	1:B:1357:HIS:HA	2.00	0.44	
1:A:1197:HIS:CE1	1:A:1294:SER:OG	2.70	0.43	
1:C:1306:ILE:HD11	1:C:1457:TRP:HD1	1.83	0.43	
1:A:1323:PRO:HD2	1:A:1408:PRO:O	2.18	0.43	
1:B:1336:MET:HG3	1:B:1421:THR:O	2.18	0.43	
1:C:1234:PHE:HB3	1:C:1239:LEU:HD11	1.99	0.43	
1:A:1426:GLU:CD	1:A:1426:GLU:H	2.22	0.42	
1:B:1459:LYS:O	1:B:1463:GLU:HB2	2.18	0.42	
1:A:1336:MET:HG3	1:A:1421:THR:O	2.19	0.42	
1:B:1247:ILE:CG2	1:B:1248:PRO:HD3	2.50	0.41	
1:A:1306:ILE:HG21	1:A:1454:VAL:HG11	2.02	0.41	
1:C:1219:MET:SD	1:C:1234:PHE:HZ	2.44	0.41	
1:A:1237:GLN:HG3	6:A:1617:HOH:O	2.21	0.41	
1:C:1333:ARG:HA	5:C:1504:PDO:H22	2.03	0.40	
1:C:1306:ILE:HD11	1:C:1457:TRP:CD1	2.57	0.40	
1:C:1328:PRO:HA	1:C:1357:HIS:HA	2.03	0.40	

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	A	279/297~(94%)	274 (98%)	3 (1%)	2 (1%)	22	12
1	В	280/297 (94%)	274 (98%)	6 (2%)	0	100	100
1	С	282/297 (95%)	276 (98%)	6 (2%)	0	100	100
All	All	841/891 (94%)	824 (98%)	15 (2%)	2 (0%)	47	39

All (2) Ramachandran outliers are listed below:

\mathbf{Mol}	Chain	Res	Type
1	A	1197	HIS



Continued from previous page...

Mol	Chain	Res	Type
1	A	1198	ALA

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric Outliers		Percentiles	
1	A	252/265~(95%)	247 (98%)	5 (2%)	55 50	
1	В	253/265 (96%)	249 (98%)	4 (2%)	62 59	
1	С	255/265~(96%)	253 (99%)	2 (1%)	81 82	
All	All	760/795 (96%)	749 (99%)	11 (1%)	65 65	

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1196	GLU
1	A	1197	HIS
1	A	1324	TYR
1	A	1335	GLU
1	A	1426	GLU
1	В	1203	PHE
1	В	1324	TYR
1	В	1417	LEU
1	В	1469	ARG
1	С	1203	PHE
1	С	1324	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1197	HIS
1	В	1249	HIS
1	С	1197	HIS
1	С	1249	HIS



5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 13 ligands modelled in this entry, 3 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Mol Type Chain Res		Link	Bond lengths			Bond angles			
MIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	U2F	A	1502	4	35,38,38	0.38	0	51,58,58	0.78	2 (3%)
5	PDO	В	1504[B]	-	4,4,4	0.30	0	3,3,3	0.33	0
3	U2F	В	1501	4	35,38,38	0.32	0	51,58,58	0.83	1 (1%)
2	NAG	С	1502	1	14,14,15	0.27	0	17,19,21	0.65	0
5	PDO	A	1504	-	4,4,4	0.36	0	3,3,3	0.23	0
3	U2F	С	1501	4	35,38,38	0.30	0	51,58,58	0.96	2 (3%)
5	PDO	С	1504	-	4,4,4	0.21	0	3,3,3	0.23	0
5	PDO	В	1504[A]	-	4,4,4	0.27	0	3,3,3	0.27	0
2	NAG	В	1502	1	14,14,15	0.28	0	17,19,21	0.72	1 (5%)
2	NAG	A	1501	1	14,14,15	0.28	0	17,19,21	0.68	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	U2F	A	1502	4	-	4/22/59/59	0/3/3/3
5	PDO	В	1504[B]	-	-	0/2/2/2	-
3	U2F	В	1501	4	-	2/22/59/59	0/3/3/3
2	NAG	С	1502	1	-	1/6/23/26	0/1/1/1
5	PDO	A	1504	-	-	1/2/2/2	-
3	U2F	С	1501	4	-	2/22/59/59	0/3/3/3
5	PDO	С	1504	-	-	0/2/2/2	-
5	PDO	В	1504[A]	-	-	0/2/2/2	-
2	NAG	В	1502	1	-	1/6/23/26	0/1/1/1
2	NAG	A	1501	1	-	1/6/23/26	0/1/1/1

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$Ideal(^{o})$
3	С	1501	U2F	O3A-PB-O1	3.33	109.19	102.48
3	В	1501	U2F	O3A-PB-O1	3.14	108.83	102.48
3	A	1502	U2F	O5-C1-O1	-3.14	107.26	111.36
3	С	1501	U2F	O5-C1-O1	-3.07	107.35	111.36
3	A	1502	U2F	PB-O1-C1	2.56	129.65	119.74
2	В	1502	NAG	C1-O5-C5	2.27	115.27	112.19

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1502	U2F	O5-C1-O1-PB
3	A	1502	U2F	C5'-O5'-PA-O2A
3	С	1501	U2F	O5-C1-O1-PB
2	В	1502	NAG	O5-C5-C6-O6
2	С	1502	NAG	O5-C5-C6-O6
2	A	1501	NAG	O5-C5-C6-O6
3	С	1501	U2F	O5-C5-C6-O6
3	В	1501	U2F	O5-C1-O1-PB
3	A	1502	U2F	C5'-O5'-PA-O3A
3	В	1501	U2F	C1-O1-PB-O3A
3	A	1502	U2F	C5'-O5'-PA-O1A
5	A	1504	PDO	C1-C2-C3-O3


There are no ring outliers.

2 monomers are involved in 4 short contacts:





Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	С	1504	PDO	2	0
5	В	1504[A]	PDO	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\# \mathrm{RSRZ}{>}2$		$OWAB(A^2)$	Q<0.9	
1	A	278/297 (93%)	0.20	7 (2%)	57	61	21, 35, 52, 83	0
1	В	278/297 (93%)	0.20	1 (0%)	92	93	20, 35, 52, 84	0
1	С	278/297 (93%)	0.18	3 (1%)	80	82	19, 35, 50, 82	0
All	All	834/891 (93%)	0.19	11 (1%)	77	79	19, 35, 51, 84	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ	
1	A	1471	VAL	2.6	
1	A	1224[A]	HIS	2.5	
1	С	1447	LEU	2.5	
1	A	1447	LEU	2.4	
1	В	1467	LEU	2.3	
1	С	1416	TRP	2.2	
1	С	1210	LEU	2.1	
1	A	1380[A]	GLN	2.1	
1	A	1261	VAL	2.1	
1	A	1247	ILE	2.0	
1	A	1457	TRP	2.0	

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.




6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
2	NAG	В	1502	14/15	0.63	0.19	64,65,66,66	0
2	NAG	С	1502	14/15	0.70	0.19	62,64,65,65	0
2	NAG	A	1501	14/15	0.75	0.13	59,61,62,62	0
5	PDO	A	1504	5/5	0.85	0.11	36,37,39,39	8
5	PDO	С	1504	5/5	0.88	0.16	40,41,42,42	8
5	PDO	В	1504[B]	5/5	0.89	0.12	35,36,36,36	13
5	PDO	В	1504[A]	5/5	0.89	0.12	18,18,18,18	13
3	U2F	A	1502	36/36	0.90	0.13	36,41,42,43	0
3	U2F	В	1501	36/36	0.91	0.13	33,36,38,39	0
3	U2F	С	1501	36/36	0.91	0.13	35,39,42,42	0
4	CA	В	1503	1/1	0.97	0.07	26,26,26,26	0
4	CA	С	1503	1/1	0.97	0.11	28,28,28,28	0
4	CA	A	1503	1/1	0.97	0.15	25,25,25,25	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.








Electron density around U2F B 1501: 2mF_o-DF_c (at 0.7 rmsd) in gray mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)







Electron density around CA B 1503:

 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around CA C 1503: $2 \mathrm{mF}_o\text{-DF}_c \text{ (at } 0.7 \text{ rmsd) in gray} \\ \mathrm{mF}_o\text{-DF}_c \text{ (at } 3 \text{ rmsd) in purple (negative)}$


and green (positive)











6.5 Other polymers (i)

There are no such residues in this entry.

