



wwPDB X-ray Structure Validation Summary Report ⓘ

May 11, 2021 – 10:15 am BST

PDB ID : 6ZOH
Title : 3-Formylrifamycin SV binding to the access pocket of AcrB-G619P_G621P L and T protomers
Authors : Tam, H.K.; Foong, W.E.; Pos, K.M.
Deposited on : 2020-07-07
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.18
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.18


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1057	 3% 91% 7%
1	B	1057	 3% 93% 5%
1	C	1057	 % 93% 5%
2	D	169	 4% 91% 5%
2	E	169	 30% 90% 1%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	LMT	C	1113	-	-	-	X

2 Entry composition [i](#)

There are 14 unique types of molecules in this entry. The entry contains 26785 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Multidrug efflux pump subunit AcrB.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	1034	7875	5070	1299	1462	44	0	2	0
1	B	1034	7866	5065	1296	1460	45	0	1	0
1	C	1033	7855	5058	1295	1458	44	0	0	0

There are 30 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	619	PRO	GLY	engineered mutation	UNP P31224
A	621	PRO	GLY	engineered mutation	UNP P31224
A	1050	LEU	-	expression tag	UNP P31224
A	1051	GLU	-	expression tag	UNP P31224
A	1052	HIS	-	expression tag	UNP P31224
A	1053	HIS	-	expression tag	UNP P31224
A	1054	HIS	-	expression tag	UNP P31224
A	1055	HIS	-	expression tag	UNP P31224
A	1056	HIS	-	expression tag	UNP P31224
A	1057	HIS	-	expression tag	UNP P31224
B	619	PRO	GLY	engineered mutation	UNP P31224
B	621	PRO	GLY	engineered mutation	UNP P31224
B	1050	LEU	-	expression tag	UNP P31224
B	1051	GLU	-	expression tag	UNP P31224
B	1052	HIS	-	expression tag	UNP P31224
B	1053	HIS	-	expression tag	UNP P31224
B	1054	HIS	-	expression tag	UNP P31224
B	1055	HIS	-	expression tag	UNP P31224
B	1056	HIS	-	expression tag	UNP P31224
B	1057	HIS	-	expression tag	UNP P31224
C	619	PRO	GLY	engineered mutation	UNP P31224
C	621	PRO	GLY	engineered mutation	UNP P31224
C	1050	LEU	-	expression tag	UNP P31224

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	1051	GLU	-	expression tag	UNP P31224
C	1052	HIS	-	expression tag	UNP P31224
C	1053	HIS	-	expression tag	UNP P31224
C	1054	HIS	-	expression tag	UNP P31224
C	1055	HIS	-	expression tag	UNP P31224
C	1056	HIS	-	expression tag	UNP P31224
C	1057	HIS	-	expression tag	UNP P31224

- Molecule 2 is a protein called DARPIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	D	154	Total	C	N	O	S	0	0	0
			1167	736	204	226	1			
2	E	153	Total	C	N	O	S	0	0	0
			1159	732	203	223	1			

- Molecule 3 is DODECYL-BETA-D-MALTOSE (three-letter code: LMT) (formula: $C_{24}H_{46}O_{11}$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
3	A	1	Total	C	O	0	0
			35	24	11		
3	A	1	Total	C	O	0	0
			35	24	11		
3	A	1	Total	C	O	0	0
			35	24	11		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			35	24	11		
3	C	1	Total	C	O	0	0
			35	24	11		
3	C	1	Total	C	O	0	0
			35	24	11		
3	C	1	Total	C	O	0	0
			35	24	11		

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	C	1	Total C O 4 2 2	0	0
4	C	1	Total C O 4 2 2	0	0
4	C	1	Total C O 4 2 2	0	0

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 6 3 3	0	0
5	B	1	Total C O 6 3 3	0	0
5	C	1	Total C O 6 3 3	0	0
5	C	1	Total C O 6 3 3	0	0

- Molecule 6 is DODECANE (three-letter code: D12) (formula: C₁₂H₂₆).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C 12 12	0	0
6	A	1	Total C 12 12	0	0
6	C	1	Total C 12 12	0	0
6	C	1	Total C 12 12	0	0

- Molecule 7 is DECYLAMINE-N,N-DIMETHYL-N-OXIDE (three-letter code: DDQ) (formula: $C_{12}H_{27}NO$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
7	A	1	14	12	1	1	0	0
7	B	1	14	12	1	1	0	0

- Molecule 8 is (2S,12Z,14E,16S,17S,18R,19R,20R,21S,22R,23S,24E)-8-formyl-5,6,9,17,19-pentahydroxy-23-methoxy-2,4,12,16,18,20,22-heptamethyl-1,11-dioxo-1,2-dihydro-2,7-(epoxypentadeca[1,11,13]trienoimino)naphtho[2,1-b]furan-21-yl acetate (three-letter code: 3YI) (formula: C₃₈H₄₇NO₁₃) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
8	A	1	52	38	1	13	0	0
8	B	1	52	38	1	13	0	0

- Molecule 9 is (2S)-3-hydroxypropane-1,2-diyl didecanoate (three-letter code: DDR) (formula: C₂₃H₄₄O₅).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	B	1	Total	C	O	0	0
			28	23	5		

- Molecule 10 is PHOSPHATIDYLETHANOLAMINE (three-letter code: PTY) (formula: $C_{40}H_{80}NO_8P$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
10	C	1	Total	C	N	O	P	0	0
			50	40	1	8	1		

- Molecule 11 is DECANE (three-letter code: D10) (formula: $C_{10}H_{22}$).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
11	C	1	Total C 10 10	0	0

- Molecule 12 is (2S)-3-[[[(R)-(2-aminoethoxy)(hydroxy)phosphoryl]oxy]-2-hydroxypropyl hexadecanoate (three-letter code: LPX) (formula: $C_{21}H_{44}NO_7P$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
12	C	1	Total C N O P 30 21 1 7 1	0	0

- Molecule 13 is HEXANE (three-letter code: HEX) (formula: C_6H_{14}).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
13	C	1	Total C 6 6	0	0
13	C	1	Total C 6 6	0	0


- Molecule 14 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
14	A	78	Total O 78 78	0	0
14	B	67	Total O 69 69	0	2
14	C	68	Total O 69 69	0	1
14	D	17	Total O 18 18	0	1
14	E	8	Total O 10 10	0	2


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Multidrug efflux pump subunit AcrB




- Molecule 1: Multidrug efflux pump subunit AcrB





- Molecule 1: Multidrug efflux pump subunit AcrB








- Molecule 2: DARPIN



- Molecule 2: DARPIN



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	145.14Å 160.13Å 243.59Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.19 – 2.80 49.19 – 2.80	Depositor EDS
% Data completeness (in resolution range)	100.0 (49.19-2.80) 100.0 (49.19-2.80)	Depositor EDS
R_{merge}	0.21	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.45 (at 2.81Å)	Xtrriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.223 , 0.245 0.225 , 0.244	Depositor DCC
R_{free} test set	6765 reflections (4.84%)	wwPDB-VP
Wilson B-factor (Å ²)	57.5	Xtrriage
Anisotropy	0.022	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 55.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	26785	wwPDB-VP
Average B, all atoms (Å ²)	64.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.17% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: DDR, GOL, 3YI, D10, D12, LMT, DDQ, LPX, PTY, EDO, HEX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.66	0/8033	0.71	0/10911
1	B	0.66	0/8021	0.70	0/10895
1	C	0.66	0/8007	0.70	0/10877
2	D	0.67	0/1186	0.71	0/1613
2	E	0.68	0/1178	0.71	0/1602
All	All	0.66	0/26425	0.70	0/35898

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7875	0	8033	39	0
1	B	7866	0	8023	25	0
1	C	7855	0	8009	24	0
2	D	1167	0	1151	0	0
2	E	1159	0	1147	0	0
3	A	105	0	138	0	0
3	B	35	0	46	0	0
3	C	105	0	138	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	16	0	24	0	0
4	B	8	0	12	0	0
4	C	16	0	24	0	0
5	A	6	0	8	0	0
5	B	6	0	8	0	0
5	C	12	0	16	0	0
6	A	24	0	52	0	0
6	C	24	0	52	0	0
7	A	14	0	27	1	0
7	B	14	0	27	0	0
8	A	52	0	0	1	0
8	B	52	0	0	1	0
9	B	28	0	44	0	0
10	C	50	0	79	0	0
11	C	10	0	22	0	0
12	C	30	0	43	0	0
13	C	12	0	28	0	0
14	A	78	0	0	0	0
14	B	69	0	0	0	0
14	C	69	0	0	0	0
14	D	18	0	0	0	0
14	E	10	0	0	0	0
All	All	26785	0	27151	85	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 85 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:209:ALA:HB1	1:B:743:ILE:HG21	1.71	0.72
1:A:968:VAL:HG11	1:A:1023:PRO:HG3	1.73	0.71
1:A:85:THR:HG21	1:A:621:PRO:HD2	1.80	0.64
1:C:372:VAL:HB	1:C:373:PRO:HD3	1.81	0.62
1:C:901:VAL:O	1:C:904:VAL:HG12	2.04	0.57

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1034/1057 (98%)	998 (96%)	33 (3%)	3 (0%)	41	72
1	B	1033/1057 (98%)	1005 (97%)	28 (3%)	0	100	100
1	C	1031/1057 (98%)	1007 (98%)	24 (2%)	0	100	100
2	D	152/169 (90%)	150 (99%)	2 (1%)	0	100	100
2	E	151/169 (89%)	147 (97%)	4 (3%)	0	100	100
All	All	3401/3509 (97%)	3307 (97%)	91 (3%)	3 (0%)	51	81

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	622	GLN
1	A	621	PRO
1	A	126	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	844/865 (98%)	838 (99%)	6 (1%)	84	95
1	B	843/865 (98%)	830 (98%)	13 (2%)	65	89
1	C	841/865 (97%)	832 (99%)	9 (1%)	73	92
2	D	119/132 (90%)	118 (99%)	1 (1%)	81	94
2	E	118/132 (89%)	117 (99%)	1 (1%)	81	94

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	2765/2859 (97%)	2735 (99%)	30 (1%)	73 92

5 of 30 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	784	ASP
1	C	984	LEU
1	B	1030	ARG
2	E	68	LYS
1	C	717	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

35 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
11	D10	C	1105	-	9,9,9	0.09	0	8,8,8	0.05	0
3	LMT	C	1113	-	36,36,36	0.54	1 (2%)	47,47,47	0.74	2 (4%)
4	EDO	C	1110	-	3,3,3	0.07	0	2,2,2	0.18	0
8	3YI	B	1107	-	55,55,55	0.96	2 (3%)	82,83,83	1.48	7 (8%)
3	LMT	A	1202	-	36,36,36	0.52	1 (2%)	47,47,47	0.74	0
6	D12	C	1104	-	11,11,11	0.10	0	10,10,10	0.05	0
4	EDO	A	1212	-	3,3,3	0.06	0	2,2,2	0.21	0
5	GOL	C	1112	-	5,5,5	0.09	0	5,5,5	0.29	0
3	LMT	A	1211	-	36,36,36	0.49	0	47,47,47	0.64	0
4	EDO	A	1203	-	3,3,3	0.06	0	2,2,2	0.15	0
13	HEX	C	1115	-	5,5,5	0.14	0	4,4,4	0.07	0
4	EDO	C	1107	-	3,3,3	0.07	0	2,2,2	0.14	0
7	DDQ	B	1106	-	10,13,13	0.14	0	12,15,15	0.31	0
5	GOL	B	1103	-	5,5,5	0.09	0	5,5,5	0.25	0
10	PTY	C	1101	-	49,49,49	0.26	0	52,54,54	0.33	0
8	3YI	A	1210	-	55,55,55	0.96	2 (3%)	82,83,83	1.44	9 (10%)
13	HEX	C	1111	-	5,5,5	0.14	0	4,4,4	0.09	0
4	EDO	C	1108	-	3,3,3	0.05	0	2,2,2	0.14	0
6	D12	A	1205	-	11,11,11	0.10	0	10,10,10	0.09	0
4	EDO	A	1207	-	3,3,3	0.06	0	2,2,2	0.18	0
12	LPX	C	1106	-	29,29,29	0.28	0	31,33,33	0.36	0
4	EDO	C	1109	-	3,3,3	0.07	0	2,2,2	0.20	0
6	D12	C	1103	-	11,11,11	0.10	0	10,10,10	0.06	0
4	EDO	B	1102	-	3,3,3	0.07	0	2,2,2	0.22	0
3	LMT	C	1114	-	36,36,36	0.51	0	47,47,47	0.75	0
7	DDQ	A	1206	-	10,13,13	0.15	0	12,15,15	0.43	0
9	DDR	B	1105	-	27,27,27	0.26	0	29,29,29	0.24	0
4	EDO	A	1209	-	3,3,3	0.07	0	2,2,2	0.21	0
3	LMT	C	1102	-	36,36,36	0.45	0	47,47,47	0.73	1 (2%)
6	D12	A	1208	-	11,11,11	0.10	0	10,10,10	0.08	0
3	LMT	A	1201	-	36,36,36	0.46	0	47,47,47	0.57	0
3	LMT	B	1101	-	36,36,36	0.48	1 (2%)	47,47,47	0.72	1 (2%)
5	GOL	C	1116	-	5,5,5	0.10	0	5,5,5	0.30	0
4	EDO	B	1104	-	3,3,3	0.07	0	2,2,2	0.19	0
5	GOL	A	1204	-	5,5,5	0.09	0	5,5,5	0.27	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
11	D10	C	1105	-	-	2/7/7/7	-
3	LMT	C	1113	-	-	12/21/61/61	0/2/2/2
4	EDO	C	1110	-	-	1/1/1/1	-
8	3YI	B	1107	-	-	14/57/72/72	0/4/4/4
3	LMT	A	1202	-	-	9/21/61/61	0/2/2/2
6	D12	C	1104	-	-	2/9/9/9	-
4	EDO	A	1212	-	-	0/1/1/1	-
5	GOL	C	1112	-	-	1/4/4/4	-
3	LMT	A	1211	-	-	12/21/61/61	0/2/2/2
4	EDO	A	1203	-	-	0/1/1/1	-
13	HEX	C	1115	-	-	1/3/3/3	-
4	EDO	C	1107	-	-	1/1/1/1	-
7	DDQ	B	1106	-	-	5/11/11/11	-
5	GOL	B	1103	-	-	2/4/4/4	-
10	PTY	C	1101	-	-	29/53/53/53	-
8	3YI	A	1210	-	-	14/57/72/72	0/4/4/4
13	HEX	C	1111	-	-	0/3/3/3	-
4	EDO	C	1108	-	-	1/1/1/1	-
6	D12	A	1205	-	-	6/9/9/9	-
4	EDO	A	1207	-	-	1/1/1/1	-
12	LPX	C	1106	-	-	11/31/31/31	-
4	EDO	C	1109	-	-	1/1/1/1	-
6	D12	C	1103	-	-	5/9/9/9	-
4	EDO	B	1102	-	-	1/1/1/1	-
3	LMT	C	1114	-	-	14/21/61/61	0/2/2/2
7	DDQ	A	1206	-	-	7/11/11/11	-
9	DDR	B	1105	-	-	12/29/29/29	-
4	EDO	A	1209	-	-	1/1/1/1	-
3	LMT	C	1102	-	-	12/21/61/61	0/2/2/2
6	D12	A	1208	-	-	2/9/9/9	-
3	LMT	A	1201	-	-	7/21/61/61	0/2/2/2
3	LMT	B	1101	-	-	7/21/61/61	0/2/2/2
5	GOL	C	1116	-	-	1/4/4/4	-
4	EDO	B	1104	-	-	0/1/1/1	-
5	GOL	A	1204	-	-	4/4/4/4	-

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	B	1107	3YI	C15-N1	4.16	1.44	1.35
8	A	1210	3YI	C15-N1	3.94	1.44	1.35
3	C	1113	LMT	O1'-C1'	2.26	1.44	1.40
3	A	1202	LMT	O1'-C1'	2.15	1.43	1.40
3	B	1101	LMT	O1'-C1'	2.11	1.43	1.40

The worst 5 of 20 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	B	1107	3YI	C2-C3-C4	6.18	123.17	119.20
8	A	1210	3YI	C2-C3-C4	5.92	123.00	119.20
8	B	1107	3YI	C4-C3-C38	-5.03	112.78	119.87
8	A	1210	3YI	C4-C3-C38	-4.63	113.35	119.87
8	B	1107	3YI	C3-C2-C1	-4.01	117.82	120.70

There are no chirality outliers.

5 of 198 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1201	LMT	C2'-C1'-O1'-C1
3	A	1201	LMT	O5'-C1'-O1'-C1
3	A	1202	LMT	O5'-C1'-O1'-C1
3	A	1211	LMT	C2'-C1'-O1'-C1
3	A	1211	LMT	O5'-C1'-O1'-C1


There are no ring outliers.


3 monomers are involved in 3 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	B	1107	3YI	1	0
8	A	1210	3YI	1	0
7	A	1206	DDQ	1	0



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.


The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	1034/1057 (97%)	0.10	34 (3%) 46 36	36, 64, 105, 127	7 (0%)
1	B	1034/1057 (97%)	0.01	33 (3%) 47 37	36, 62, 94, 111	5 (0%)
1	C	1033/1057 (97%)	-0.07	13 (1%) 77 72	37, 54, 79, 101	0
2	D	154/169 (91%)	0.24	7 (4%) 33 23	50, 62, 84, 93	0
2	E	153/169 (90%)	1.36	51 (33%) 0 0	60, 78, 100, 109	0
All	All	3408/3509 (97%)	0.09	138 (4%) 38 28	36, 60, 96, 127	12 (0%)

The worst 5 of 138 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	E	35	ALA	5.7
2	E	37	GLY	5.0
2	E	33	LEU	5.0
1	A	869	SER	4.9
2	E	68	LYS	4.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,




median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	GOL	C	1116	6/6	0.69	0.37	84,86,86,86	0
7	DDQ	A	1206	14/14	0.70	0.36	110,115,121,121	0
3	LMT	C	1114	35/35	0.72	0.34	122,126,138,139	0
6	D12	A	1208	12/12	0.73	0.30	79,83,84,85	0
3	LMT	A	1202	35/35	0.75	0.28	90,116,132,132	0
3	LMT	C	1113	35/35	0.75	0.41	116,128,132,133	0
4	EDO	A	1207	4/4	0.77	0.35	76,77,78,80	0
4	EDO	A	1212	4/4	0.79	0.34	71,71,72,72	0
6	D12	A	1205	12/12	0.80	0.27	93,97,97,98	0
4	EDO	C	1110	4/4	0.80	0.18	88,89,89,90	0
4	EDO	A	1209	4/4	0.80	0.32	67,67,68,68	0
10	PTY	C	1101	50/50	0.80	0.39	84,91,118,122	0
4	EDO	A	1203	4/4	0.81	0.31	66,67,67,68	0
4	EDO	B	1102	4/4	0.81	0.22	82,83,83,83	0
3	LMT	A	1211	35/35	0.83	0.27	87,103,116,117	0
7	DDQ	B	1106	14/14	0.83	0.25	74,86,97,97	0
6	D12	C	1104	12/12	0.83	0.30	73,74,75,75	0
3	LMT	A	1201	35/35	0.85	0.46	104,110,119,119	0
12	LPX	C	1106	30/30	0.85	0.27	88,93,96,97	0
11	D10	C	1105	10/10	0.86	0.49	66,67,68,68	0
13	HEX	C	1111	6/6	0.86	0.43	69,69,70,70	0
3	LMT	B	1101	35/35	0.87	0.33	94,96,106,107	0
5	GOL	A	1204	6/6	0.87	0.32	70,70,71,71	0
6	D12	C	1103	12/12	0.88	0.25	66,68,71,72	0
9	DDR	B	1105	28/28	0.89	0.41	93,104,111,112	0
8	3YI	A	1210	52/52	0.90	0.21	80,86,90,90	0
4	EDO	C	1107	4/4	0.92	0.25	56,57,57,57	0
3	LMT	C	1102	35/35	0.92	0.22	72,74,76,76	0
8	3YI	B	1107	52/52	0.93	0.24	70,77,79,80	0
4	EDO	C	1108	4/4	0.93	0.27	71,71,72,73	0
4	EDO	C	1109	4/4	0.93	0.27	52,53,53,54	0
13	HEX	C	1115	6/6	0.93	0.21	61,62,62,62	0
4	EDO	B	1104	4/4	0.94	0.24	59,60,60,60	0
5	GOL	B	1103	6/6	0.94	0.26	62,63,64,65	0
5	GOL	C	1112	6/6	0.94	0.23	58,59,59,60	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around LMT C 1114:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around LMT A 1202:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around LMT C 1113:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around PTY C 1101:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around LMT A 1211:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around LMT A 1201:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around LPX C 1106:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around LMT B 1101:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around DDR B 1105:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 3YI A 1210:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around LMT C 1102:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.