

wwPDB X-ray Structure Validation Summary Report (i)

Sep 17, 2023 – 06:49 PM EDT

PDB ID	:	4ZU7
Title	:	X-ray structure if the QdtA 3,4-ketoisomerase from Thermoanaerobacterium
		thermosaccharolyticum, double mutant $Y17R/R97H$, in complex with TDP
Authors	:	Thoden, J.B.; Vinogradov, E.; Gilbert, M.; Salinger, A.J.; Holden, H.M.
Deposited on	:	2015-05-15
Resolution	:	2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


Xtriage (Phenix) EDS buster-report Percentile statistics Refmac CCP4 Ideal geometry (proteins) Ideal geometry (DNA, RNA)	: : : : :	20191225.v01 (using entries in the PDB archive December 25th 2019) 5.8.0158 7.0.044 (Gargrove) Engh & Huber (2001) Parkinson et al. (1996)
Ideal geometry (DNA, RNA) Validation Pipeline (wwPDB-VP)		Parkinson et al. (1996) 2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \textbf{Whole archive} \\ \textbf{(\#Entries)} \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
			2%	
1	А	144	83%	11% • 5%
	-		.% ■	
1	В	144	84%	10% •• •
	~		2%	
1	С	144	90%	6% • •
	-		.% ■	
1	D	144	83%	10% • 5%
	_		6%	
1	Е	144	83%	12% 5%

Continued on next page...



Mol	Chain	Length	Quality of chain		
1	F	144	2% 81%	13%	
1	G	144	5%	14%	• 6%
1	Н	144	3% 82%	12%	6%



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 9552 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	А	137	Total	С	Ν	0	S	0	2	0
	A	197	1155	760	186	205	4	0		0
1	В	138	Total	С	Ν	Ο	S	0	0	0
	D	130	1150	752	186	208	4	0	0	0
1	С	138	Total	С	Ν	0	S	0	0	0
		130	1150	752	186	208	4	0	0	0
1	D	137	Total	С	Ν	0	S	0	1	0
	D	137	1149	755	185	205	4	0	T	0
1	Е	137	Total	С	Ν	0	S	0	0	0
	Ľ	157	1135	742	185	205	3	0	0	
1	F	138	Total	С	Ν	Ο	S	0	0	0
	Г	130	1146	750	186	206	4	0	0	0
1	G	136	Total	С	Ν	0	S	0	0	0
	G	130	1133	741	184	204	4	0	0	0
1	Н	136	Total	С	Ν	0	S	0	0	0
	11	130	1130	739	184	204	3		0	0

• Molecule 1 is a protein called QdtA.

There are 112 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	17	ARG	TYR	engineered mutation	UNP Q6TFC5
А	52	SER	ALA	SEE REMARK 999	UNP Q6TFC5
А	94	LEU	PHE	SEE REMARK 999	UNP Q6TFC5
А	95	VAL	LEU	SEE REMARK 999	UNP Q6TFC5
А	97	HIS	ARG	engineered mutation	UNP Q6TFC5
А	116	TYR	CYS	SEE REMARK 999	UNP Q6TFC5
А	137	LEU	-	expression tag	UNP Q6TFC5
А	138	GLU	-	expression tag	UNP Q6TFC5
А	139	HIS	-	expression tag	UNP Q6TFC5
А	140	HIS	-	expression tag	UNP Q6TFC5
А	141	HIS	-	expression tag	UNP Q6TFC5
А	142	HIS	-	expression tag	UNP Q6TFC5
А	143	HIS	-	expression tag	UNP Q6TFC5

Continued on next page...



Chain	Residue	Modelled	Actual	Comment	Reference
А	144	HIS	-	expression tag	UNP Q6TFC5
В	17	ARG	TYR	engineered mutation	UNP Q6TFC5
В	52	SER	ALA	SEE REMARK 999	UNP Q6TFC5
В	94	LEU	PHE	SEE REMARK 999	UNP Q6TFC5
В	95	VAL	LEU	SEE REMARK 999	UNP Q6TFC5
В	97	HIS	ARG	engineered mutation	UNP Q6TFC5
В	116	TYR	CYS	SEE REMARK 999	UNP Q6TFC5
В	137	LEU	-	expression tag	UNP Q6TFC5
В	138	GLU	-	expression tag	UNP Q6TFC5
В	139	HIS	-	expression tag	UNP Q6TFC5
В	140	HIS	-	expression tag	UNP Q6TFC5
В	141	HIS	-	expression tag	UNP Q6TFC5
В	142	HIS	-	expression tag	UNP Q6TFC5
В	143	HIS	-	expression tag	UNP Q6TFC5
В	144	HIS	-	expression tag	UNP Q6TFC5
С	17	ARG	TYR	engineered mutation	UNP Q6TFC5
С	52	SER	ALA	SEE REMARK 999	UNP Q6TFC5
С	94	LEU	PHE	SEE REMARK 999	UNP Q6TFC5
С	95	VAL	LEU	SEE REMARK 999	UNP Q6TFC5
С	97	HIS	ARG	engineered mutation	UNP Q6TFC5
С	116	TYR	CYS	SEE REMARK 999	UNP Q6TFC5
С	137	LEU	-	expression tag	UNP Q6TFC5
С	138	GLU	-	expression tag	UNP Q6TFC5
С	139	HIS	-	expression tag	UNP Q6TFC5
С	140	HIS	-	expression tag	UNP Q6TFC5
С	141	HIS	-	expression tag	UNP Q6TFC5
С	142	HIS	-	expression tag	UNP Q6TFC5
С	143	HIS	-	expression tag	UNP Q6TFC5
С	144	HIS	-	expression tag	UNP Q6TFC5
D	17	ARG	TYR	engineered mutation	UNP Q6TFC5
D	52	SER	ALA	SEE REMARK 999	UNP Q6TFC5
D	94	LEU	PHE	SEE REMARK 999	UNP Q6TFC5
D	95	VAL	LEU	SEE REMARK 999	UNP Q6TFC5
D	97	HIS	ARG	engineered mutation	UNP Q6TFC5
D	116	TYR	CYS	SEE REMARK 999	UNP Q6TFC5
D	137	LEU	-	expression tag	UNP Q6TFC5
D	138	GLU	-	expression tag	UNP Q6TFC5
D	139	HIS	-	expression tag	UNP Q6TFC5
D	140	HIS	-	expression tag	UNP Q6TFC5
D	141	HIS	-	expression tag	UNP Q6TFC5
D	142	HIS	-	expression tag	UNP Q6TFC5
D	143	HIS	-	expression tag	UNP Q6TFC5

Continued on next page...




Chain	Residue	Modelled	Actual	Comment	Reference
D	144	HIS	-	expression tag	UNP Q6TFC5
Е	17	ARG	TYR	engineered mutation	UNP Q6TFC5
Е	52	SER	ALA	SEE REMARK 999	UNP Q6TFC5
Е	94	LEU	PHE	SEE REMARK 999	UNP Q6TFC5
Е	95	VAL	LEU	SEE REMARK 999	UNP Q6TFC5
Е	97	HIS	ARG	engineered mutation	UNP Q6TFC5
Е	116	TYR	CYS	SEE REMARK 999	UNP Q6TFC5
Е	137	LEU	-	expression tag	UNP Q6TFC5
Е	138	GLU	-	expression tag	UNP Q6TFC5
Е	139	HIS	-	expression tag	UNP Q6TFC5
Е	140	HIS	-	expression tag	UNP Q6TFC5
Е	141	HIS	-	expression tag	UNP Q6TFC5
Е	142	HIS	-	expression tag	UNP Q6TFC5
Е	143	HIS	-	expression tag	UNP Q6TFC5
Ε	144	HIS	-	expression tag	UNP Q6TFC5
F	17	ARG	TYR	engineered mutation	UNP Q6TFC5
F	52	SER	ALA	SEE REMARK 999	UNP Q6TFC5
F	94	LEU	PHE	SEE REMARK 999	UNP Q6TFC5
F	95	VAL	LEU	SEE REMARK 999	UNP Q6TFC5
F	97	HIS	ARG	engineered mutation	UNP Q6TFC5
F	116	TYR	CYS	SEE REMARK 999	UNP Q6TFC5
F	137	LEU	-	expression tag	UNP Q6TFC5
F	138	GLU	-	expression tag	UNP Q6TFC5
F	139	HIS	-	expression tag	UNP Q6TFC5
F	140	HIS	-	expression tag	UNP Q6TFC5
F	141	HIS	-	expression tag	UNP Q6TFC5
F	142	HIS	-	expression tag	UNP Q6TFC5
F	143	HIS	-	expression tag	UNP Q6TFC5
F	144	HIS	-	expression tag	UNP Q6TFC5
G	17	ARG	TYR	engineered mutation	UNP Q6TFC5
G	52	SER	ALA	SEE REMARK 999	UNP Q6TFC5
G	94	LEU	PHE	SEE REMARK 999	UNP Q6TFC5
G	95	VAL	LEU	SEE REMARK 999	UNP Q6TFC5
G	97	HIS	ARG	engineered mutation	UNP Q6TFC5
G	116	TYR	CYS	SEE REMARK 999	UNP Q6TFC5
G	137	LEU	-	expression tag	UNP Q6TFC5
G	138	GLU	-	expression tag	UNP Q6TFC5
G	139	HIS	-	expression tag	UNP Q6TFC5
G	140	HIS	-	expression tag	UNP Q6TFC5
G	141	HIS	-	expression tag	UNP Q6TFC5
G	142	HIS	-	expression tag	UNP Q6TFC5
G	143	HIS	-	expression tag	UNP Q6TFC5

Continued on next page...



Chain	Residue	Modelled	Actual	Comment	Reference
G	144	HIS	-	expression tag	UNP Q6TFC5
Н	17	ARG	TYR	engineered mutation	UNP Q6TFC5
Н	52	SER	ALA	SEE REMARK 999	UNP Q6TFC5
Н	94	LEU	PHE	SEE REMARK 999	UNP Q6TFC5
Н	95	VAL	LEU	SEE REMARK 999	UNP Q6TFC5
Н	97	HIS	ARG	engineered mutation	UNP Q6TFC5
Н	116	TYR	CYS	SEE REMARK 999	UNP Q6TFC5
Н	137	LEU	-	expression tag	UNP Q6TFC5
Н	138	GLU	-	expression tag	UNP Q6TFC5
Н	139	HIS	-	expression tag	UNP Q6TFC5
Н	140	HIS	-	expression tag	UNP Q6TFC5
Н	141	HIS	-	expression tag	UNP Q6TFC5
Н	142	HIS	- expression tag		UNP Q6TFC5
Н	143	HIS	-	expression tag	UNP Q6TFC5
Н	144	HIS	-	expression tag	UNP Q6TFC5

• Molecule 2 is THYMIDINE-5'-DIPHOSPHATE (three-letter code: TYD) (formula: $C_{10}H_{16}N_2O_{11}P_2$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	Δ	1	Total	С	Ν	Ο	Р	0	0
2	Л	1	25	10	2	11	2	0	0
2	B	1	Total	С	Ν	Ο	Р	0	0
	D	1	25	10	2	11	2	0	0
9	С	1	Total	С	Ν	Ο	Р	0	0
	U	1	25	10	2	11	2	0	0

Continued on next page...



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	П	1	Total	С	Ν	Ο	Р	0	0
	D	1	25	10	2	11	2	0	0
2	E	1	Total	С	Ν	Ο	Р	0	0
	E	1	25	10	2	11	2	0	0
2	F	1	Total	С	Ν	Ο	Р	0	0
	Г	1	25	10	2	11	2	0	
2	G	1	Total	С	Ν	Ο	Р	0	0
	G	1	25	10	2	11	2	0	0
9	Н	1	Total	С	Ν	Ο	Р	0	0
	11	1	25	10	2	11	2	0	0

• Molecule 3 is (2S)-1-[3-{[(2R)-2-hydroxypropyl]oxy}-2,2-bis({[(2R)-2-hydroxypropyl]oxy}m ethyl)propoxy]propan-2-ol (three-letter code: PXN) (formula: $C_{17}H_{36}O_8$).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	
3	D	1	Total 25	C 17	O 8	0	0

• Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	36	$\begin{array}{cc} \text{Total} & \text{O} \\ 36 & 36 \end{array}$	0	0
4	В	27	Total O 27 27	0	0
4	С	31	Total O 31 31	0	0

Continued on next page...




Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	D	33	Total O 33 33	0	0
4	Е	12	Total O 12 12	0	0
4	F	19	Total O 19 19	0	0
4	G	10	Total O 10 10	0	0
4	Н	11	Total O 11 11	0	0



3 Residue-property plots (i)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: QdtA









4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants	136.56Å 96.17Å 111.75Å	Depositor
a, b, c, α , β , γ	90.00° 114.50° 90.00°	Depositor
Resolution (Å)	29.28 - 2.30	Depositor
Resolution (A)	29.26 - 2.30	EDS
% Data completeness	90.3 (29.28-2.30)	Depositor
(in resolution range)	90.4 (29.26-2.30)	EDS
R _{merge}	0.07	Depositor
R _{sym}	0.07	Depositor
$< I/\sigma(I) > 1$	$1.69 (at 2.31 \text{\AA})$	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
D D.	0.192 , 0.256	Depositor
R, R_{free}	0.196 , 0.257	DCC
R_{free} test set	2660 reflections $(5.03%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	38.9	Xtriage
Anisotropy	0.317	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.31, 28.5	EDS
L-test for twinning ²	$ < L >=0.52, < L^2>=0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	9552	wwPDB-VP
Average B, all atoms $(Å^2)$	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.34% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: TYD, PXN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles	
	Unam	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.78	0/1189	0.93	2/1604~(0.1%)
1	В	0.67	0/1177	0.87	3/1589~(0.2%)
1	С	0.73	0/1177	0.94	3/1589~(0.2%)
1	D	0.73	0/1180	0.91	1/1593~(0.1%)
1	Ε	0.62	0/1162	0.80	0/1570
1	F	0.68	0/1173	0.87	1/1584~(0.1%)
1	G	0.60	0/1160	0.82	0/1566
1	Н	0.63	0/1157	0.85	0/1563
All	All	0.68	0/9375	0.87	10/12658~(0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	1
1	F	0	1
All	All	0	2

There are no bond length outliers.

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	$Observed(^{o})$	$Ideal(^{o})$
1	С	17	ARG	NE-CZ-NH1	7.67	124.13	120.30
1	С	17	ARG	NE-CZ-NH2	-7.00	116.80	120.30
1	F	2	LEU	N-CA-C	-6.34	93.89	111.00
1	В	17	ARG	NE-CZ-NH1	6.16	123.38	120.30
1	В	17	ARG	NE-CZ-NH2	-5.79	117.41	120.30



There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	104	GLU	Peptide
1	F	104	GLU	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	1155	0	1181	7	1
1	В	1150	0	1165	8	0
1	С	1150	0	1165	5	0
1	D	1149	0	1168	8	1
1	Е	1135	0	1140	11	0
1	F	1146	0	1161	10	0
1	G	1133	0	1148	13	0
1	Н	1130	0	1141	7	0
2	А	25	0	13	0	0
2	В	25	0	13	4	0
2	С	25	0	13	1	0
2	D	25	0	13	0	0
2	Е	25	0	13	2	0
2	F	25	0	13	1	0
2	G	25	0	13	0	0
2	Н	25	0	13	1	0
3	D	25	0	36	0	0
4	А	36	0	0	0	0
4	В	27	0	0	0	0
4	С	31	0	0	2	0
4	D	33	0	0	0	0
4	Е	12	0	0	0	0
4	F	19	0	0	0	0
4	G	10	0	0	0	0
4	Н	11	0	0	0	0
All	All	9552	0	9409	63	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:201:TYD:H5'1	2:B:201:TYD:H6	1.60	0.84
1:G:2:LEU:HD12	1:G:78:LYS:HB2	1.66	0.78
1:B:6:ALA:HB2	1:G:6:ALA:HB2	1.74	0.68
2:B:201:TYD:H6	2:B:201:TYD:C5'	2.23	0.68
2:C:201:TYD:O2A	4:C:301:HOH:O	2.14	0.64

The worst 5 of 63 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:126:ASP:OD2	$1:D:72:LYS:NZ[4_455]$	2.18	0.02

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	137/144~(95%)	131~(96%)	6 (4%)	0	100	100
1	В	136/144~(94%)	129~(95%)	7~(5%)	0	100	100
1	С	136/144~(94%)	130~(96%)	6 (4%)	0	100	100
1	D	136/144~(94%)	130~(96%)	6~(4%)	0	100	100
1	Ε	135/144~(94%)	129~(96%)	6 (4%)	0	100	100
1	F	136/144~(94%)	124 (91%)	8 (6%)	4(3%)	4	3
1	G	134/144~(93%)	127~(95%)	7 (5%)	0	100	100
1	Н	134/144~(93%)	125~(93%)	8 (6%)	1 (1%)	22	26
All	All	1084/1152~(94%)	1025 (95%)	54 (5%)	5~(0%)	29	35

All (5) Ramachandran outliers are listed below:



Mol	Chain	Res	Type
1	F	2	LEU
1	F	103	THR
1	F	22	PRO
1	F	102	PHE
1	Н	22	PRO

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the side chain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	128/133~(96%)	124~(97%)	4 (3%)	40 55
1	В	127/133~(96%)	121 (95%)	6 (5%)	26 37
1	С	127/133~(96%)	124 (98%)	3~(2%)	49 66
1	D	127/133~(96%)	122 (96%)	5 (4%)	32 46
1	Ε	124/133~(93%)	119~(96%)	5(4%)	31 44
1	F	126/133~(95%)	120~(95%)	6~(5%)	25 36
1	G	125/133~(94%)	116 (93%)	9~(7%)	14 18
1	Н	124/133~(93%)	118 (95%)	6 (5%)	25 36
All	All	1008/1064~(95%)	964 (96%)	44 (4%)	28 39

5 of 44 residues with a non-rotameric side chain are listed below:

Mol	Chain	Res	Type
1	F	137	LEU
1	G	99	MET
1	G	1	MET
1	G	45	ASP
1	G	133	LYS

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such side chains are listed below:

Mol	Chain	Res	Type					
1	А	64	ASN					
	Continued on next page							



Continued from previous page...

Mol	Chain	Res	Type
1	D	58	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Turne	Chain	Res	Link	Bo	ond leng	ths	В	ond ang	les
IVIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2
2	TYD	С	201	-	21,26,26	0.93	1 (4%)	27,40,40	1.97	2 (7%)
2	TYD	G	201	-	21,26,26	0.98	1 (4%)	27,40,40	1.82	3 (11%)
3	PXN	D	202	-	20,24,24	0.32	0	22,30,30	0.85	0
2	TYD	Е	201	-	21,26,26	1.09	1 (4%)	27,40,40	1.80	3 (11%)
2	TYD	А	201	-	21,26,26	1.16	1 (4%)	27,40,40	2.00	4 (14%)
2	TYD	В	201	-	21,26,26	1.00	1 (4%)	27,40,40	2.08	3 (11%)
2	TYD	D	201	-	21,26,26	1.15	2 (9%)	27,40,40	1.88	1 (3%)
2	TYD	Н	201	-	21,26,26	1.08	1 (4%)	27,40,40	1.76	1 (3%)
2	TYD	F	201	-	21,26,26	1.11	1 (4%)	27,40,40	1.93	3 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral



centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TYD	С	201	-	-	0/13/28/28	0/2/2/2
2	TYD	G	201	-	-	2/13/28/28	0/2/2/2
3	PXN	D	202	-	-	20/28/28/28	-
2	TYD	Е	201	-	-	0/13/28/28	0/2/2/2
2	TYD	А	201	-	-	0/13/28/28	0/2/2/2
2	TYD	В	201	-	-	6/13/28/28	0/2/2/2
2	TYD	D	201	-	-	3/13/28/28	0/2/2/2
2	TYD	Н	201	-	-	1/13/28/28	0/2/2/2
2	TYD	F	201	-	-	5/13/28/28	0/2/2/2

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	201	TYD	C5-C4	3.95	1.49	1.41
2	А	201	TYD	C5-C4	3.94	1.49	1.41
2	Н	201	TYD	C5-C4	3.93	1.49	1.41
2	Е	201	TYD	C5-C4	3.70	1.49	1.41
2	D	201	TYD	C5-C4	3.63	1.49	1.41

The worst 5 of 20 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	С	201	TYD	C2-N3-C4	8.76	122.54	115.14
2	В	201	TYD	C2-N3-C4	8.56	122.37	115.14
2	D	201	TYD	C2-N3-C4	8.41	122.24	115.14
2	Н	201	TYD	C2-N3-C4	8.37	122.21	115.14
2	А	201	TYD	C2-N3-C4	8.01	121.90	115.14

There are no chirality outliers.

5 of 37 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	В	201	TYD	C5'-O5'-PA-O1A
2	В	201	TYD	C5'-O5'-PA-O2A
2	D	201	TYD	C5'-O5'-PA-O1A
2	F	201	TYD	O4'-C1'-N1-C6
3	D	202	PXN	CAY-CAT-CAU-OAV




There are no ring outliers.

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	С	201	TYD	1	0
2	Е	201	TYD	2	0
2	В	201	TYD	4	0
2	Н	201	TYD	1	0
2	F	201	TYD	1	0


5 monomers are involved in 9 short contacts:

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





























5.7 Other polymers (i)

There are no such residues in this entry.



5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle RSRZ \rangle$	#RSRZ>2	$OWAB(Å^2)$	$\mathbf{Q} \! < \! 0.9$
1	А	137/144~(95%)	-0.26	3 (2%) 62 69	20, 29, 49, 88	0
1	В	138/144~(95%)	-0.25	2 (1%) 75 80	22, 35, 57, 84	0
1	С	138/144 (95%)	-0.21	3 (2%) 62 69	21, 33, 55, 88	0
1	D	137/144~(95%)	-0.35	1 (0%) 87 91	20, 31, 54, 87	0
1	Ε	137/144~(95%)	0.15	9 (6%) 18 23	25, 44, 80, 90	0
1	F	138/144~(95%)	-0.10	3 (2%) 62 69	25, 40, 70, 75	0
1	G	136/144~(94%)	0.02	7 (5%) 28 35	30, 48, 79, 87	0
1	Н	136/144 (94%)	0.04	5 (3%) 41 48	29, 46, 71, 88	0
All	All	1097/1152~(95%)	-0.12	33 (3%) 50 57	20, 38, 69, 90	0

The worst 5 of 33 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	Е	134	LYS	3.7
1	Е	133	LYS	3.7
1	С	1	MET	3.6
1	Е	127	PHE	3.4
1	G	127	PHE	3.2

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.




6.4 Ligands (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathrm{\AA}^2)$	Q<0.9
2	TYD	G	201	25/25	0.88	0.16	44,72,93,102	0
2	TYD	Н	201	25/25	0.91	0.17	51,70,82,88	0
2	TYD	F	201	25/25	0.92	0.13	39,54,71,76	0
2	TYD	С	201	25/25	0.92	0.12	34,55,77,78	0
2	TYD	Е	201	25/25	0.92	0.13	42,58,70,71	0
2	TYD	В	201	25/25	0.93	0.12	$42,\!55,\!64,\!69$	0
2	TYD	D	201	25/25	0.94	0.12	32,48,56,58	0
3	PXN	D	202	25/25	0.94	0.15	29,41,48,52	0
2	TYD	А	201	25/25	0.95	0.11	32,42,49,53	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.









































6.5 Other polymers (i)

There are no such residues in this entry.

