



# wwPDB X-ray Structure Validation Summary Report ⓘ

Apr 29, 2024 – 06:14 am BST

PDB ID : 3ZWL  
Title : Structure of eukaryotic translation initiation factor eIF3i complex with eIF3b C-terminus (655-700)  
Authors : Daujotyte, D.; Lukavsky, P.J.  
Deposited on : 2011-08-01  
Resolution : 2.20 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.36.2  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	369	 80% 11% 8%
1	D	369	 84% 11% 5%
2	E	50	 4% 66% 12% 22%
2	F	50	 4% 70% 8% 22%

## 2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 6381 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called EUKARYOTIC TRANSLATION INITIATION FACTOR 3 SUBUNIT I.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	B	339	2683	1709	441	524	9	0	0	0
1	D	352	2774	1762	457	546	9	0	0	0

There are 44 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-21	MET	-	expression tag	UNP P40217
B	-20	GLY	-	expression tag	UNP P40217
B	-19	SER	-	expression tag	UNP P40217
B	-18	SER	-	expression tag	UNP P40217
B	-17	HIS	-	expression tag	UNP P40217
B	-16	HIS	-	expression tag	UNP P40217
B	-15	HIS	-	expression tag	UNP P40217
B	-14	HIS	-	expression tag	UNP P40217
B	-13	HIS	-	expression tag	UNP P40217
B	-12	HIS	-	expression tag	UNP P40217
B	-11	SER	-	expression tag	UNP P40217
B	-10	SER	-	expression tag	UNP P40217
B	-9	GLY	-	expression tag	UNP P40217
B	-8	GLU	-	expression tag	UNP P40217
B	-7	ASN	-	expression tag	UNP P40217
B	-6	LEU	-	expression tag	UNP P40217
B	-5	TYR	-	expression tag	UNP P40217
B	-4	PHE	-	expression tag	UNP P40217
B	-3	GLN	-	expression tag	UNP P40217
B	-2	GLY	-	expression tag	UNP P40217
B	-1	SER	-	expression tag	UNP P40217
B	0	HIS	-	expression tag	UNP P40217
D	-21	MET	-	expression tag	UNP P40217
D	-20	GLY	-	expression tag	UNP P40217

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
D	-19	SER	-	expression tag	UNP P40217
D	-18	SER	-	expression tag	UNP P40217
D	-17	HIS	-	expression tag	UNP P40217
D	-16	HIS	-	expression tag	UNP P40217
D	-15	HIS	-	expression tag	UNP P40217
D	-14	HIS	-	expression tag	UNP P40217
D	-13	HIS	-	expression tag	UNP P40217
D	-12	HIS	-	expression tag	UNP P40217
D	-11	SER	-	expression tag	UNP P40217
D	-10	SER	-	expression tag	UNP P40217
D	-9	GLY	-	expression tag	UNP P40217
D	-8	GLU	-	expression tag	UNP P40217
D	-7	ASN	-	expression tag	UNP P40217
D	-6	LEU	-	expression tag	UNP P40217
D	-5	TYR	-	expression tag	UNP P40217
D	-4	PHE	-	expression tag	UNP P40217
D	-3	GLN	-	expression tag	UNP P40217
D	-2	GLY	-	expression tag	UNP P40217
D	-1	SER	-	expression tag	UNP P40217
D	0	HIS	-	expression tag	UNP P40217

- Molecule 2 is a protein called EUKARYOTIC TRANSLATION INITIATION FACTOR 3 SUBUNIT B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	39	Total	C	N	O	S	0	0	0
			344	219	60	63	2			
2	F	39	Total	C	N	O	S	0	0	0
			344	219	60	63	2			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	651	GLY	-	expression tag	UNP P06103
E	652	SER	-	expression tag	UNP P06103
E	653	HIS	-	expression tag	UNP P06103
F	651	GLY	-	expression tag	UNP P06103
F	652	SER	-	expression tag	UNP P06103
F	653	HIS	-	expression tag	UNP P06103


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	106	Total 106	O 106	0	0
3	D	124	Total 124	O 124	0	0
3	E	2	Total 2	O 2	0	0
3	F	4	Total 4	O 4	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: EUKARYOTIC TRANSLATION INITIATION FACTOR 3 SUBUNIT I




- Molecule 1: EUKARYOTIC TRANSLATION INITIATION FACTOR 3 SUBUNIT I



- Molecule 2: EUKARYOTIC TRANSLATION INITIATION FACTOR 3 SUBUNIT B



- Molecule 2: EUKARYOTIC TRANSLATION INITIATION FACTOR 3 SUBUNIT B



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	126.47Å 126.47Å 105.74Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	109.53 – 2.20 41.40 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.8 (109.53-2.20) 99.8 (41.40-2.20)	Depositor EDS
$R_{merge}$	0.14	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.18 (at 2.20Å)	Xtrriage
Refinement program	REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.189 , 0.247 0.183 , 0.240	Depositor DCC
$R_{free}$ test set	2526 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	38.2	Xtrriage
Anisotropy	0.268	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 34.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.033 for -h,-k,l	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	6381	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	39.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.13% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	B	1.10	3/2749 (0.1%)	0.93	5/3720 (0.1%)
1	D	1.14	3/2841 (0.1%)	0.97	6/3846 (0.2%)
2	E	0.98	0/350	0.90	0/466
2	F	0.91	0/350	1.00	2/466 (0.4%)
All	All	1.10	6/6290 (0.1%)	0.95	13/8498 (0.2%)

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	67	CYS	CB-SG	-8.37	1.68	1.82
1	B	104	CYS	CB-SG	7.54	1.95	1.82
1	D	196	GLU	CG-CD	6.83	1.62	1.51
1	B	67	CYS	CB-SG	-6.54	1.71	1.82
1	D	196	GLU	CB-CG	5.65	1.62	1.52

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	24	LEU	CB-CG-CD1	7.88	124.39	111.00
2	F	678	ARG	NE-CZ-NH2	-7.46	116.57	120.30
2	F	678	ARG	CG-CD-NE	-7.23	96.62	111.80
1	D	129	ARG	NE-CZ-NH2	-7.15	116.72	120.30
1	B	129	ARG	NE-CZ-NH2	-6.95	116.83	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen



atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	2683	0	2585	22	0
1	D	2774	0	2676	21	0
2	E	344	0	342	4	0
2	F	344	0	342	1	0
3	B	106	0	0	1	0
3	D	124	0	0	4	0
3	E	2	0	0	0	0
3	F	4	0	0	0	0
All	All	6381	0	5945	48	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 48 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:307:SER:HB2	1:D:320:HIS:O	1.93	0.68
1:D:221:PHE:CE1	1:D:233:LYS:HG3	2.31	0.65
1:B:177:ILE:HD12	1:B:194:LEU:HD22	1.80	0.62
1:D:30:LYS:HA	1:D:54:THR:HG23	1.82	0.61
1:B:307:SER:HB2	1:B:320:HIS:O	2.01	0.61

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	335/369 (91%)	321 (96%)	13 (4%)	1 (0%)	41	46
1	D	350/369 (95%)	332 (95%)	17 (5%)	1 (0%)	41	46

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	E	37/50 (74%)	33 (89%)	3 (8%)	1 (3%)	5	2
2	F	37/50 (74%)	35 (95%)	2 (5%)	0	100	100
All	All	759/838 (91%)	721 (95%)	35 (5%)	3 (0%)	34	37

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	8	GLY
2	E	688	SER
1	D	8	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	295/320 (92%)	288 (98%)	7 (2%)	49	62
1	D	305/320 (95%)	296 (97%)	9 (3%)	41	53
2	E	38/46 (83%)	38 (100%)	0	100	100
2	F	38/46 (83%)	37 (97%)	1 (3%)	46	58
All	All	676/732 (92%)	659 (98%)	17 (2%)	47	60

5 of 17 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	302	SER
2	F	679	GLU
1	D	-1	SER
1	D	5	LYS
1	D	65	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	85	GLN
1	D	269	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	B	339/369 (91%)	-0.34	5 (1%) 73 72	22, 35, 56, 98	0
1	D	352/369 (95%)	-0.31	5 (1%) 75 73	21, 34, 57, 101	0
2	E	39/50 (78%)	0.27	2 (5%) 28 26	37, 55, 73, 83	0
2	F	39/50 (78%)	0.29	2 (5%) 28 26	37, 55, 74, 82	0
All	All	769/838 (91%)	-0.26	14 (1%) 68 66	21, 36, 64, 101	0

The worst 5 of 14 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	293	PHE	4.1
1	D	341	GLU	3.6
2	F	661	ARG	3.4
1	B	342	HIS	2.9
1	D	342	HIS	2.9

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.