



wwPDB X-ray Structure Validation Summary Report ⓘ

Apr 30, 2024 – 11:18 pm BST

PDB ID : 3ZXU
Title : Crystal structure of the Ctf19-Mcm21 kinetochore heterodimer from yeast
Authors : Schmitzberger, F.; Harrison, S.C.
Deposited on : 2011-08-15
Resolution : 3.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1049 (3.88-3.52)
Clashscore	141614	1027 (3.86-3.54)
Ramachandran outliers	138981	1069 (3.88-3.52)
Sidechain outliers	138945	1065 (3.88-3.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	296	43% (green), 26% (yellow), 29% (grey)
1	C	296	43% (green), 25% (yellow), 29% (grey)
2	B	270	50% (green), 23% (yellow), 27% (grey)
2	D	270	48% (green), 25% (yellow), 27% (grey)

2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 13474 atoms, of which 6763 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MCM21.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	211	Total	C	H	N	O	S	0	0	0
			3487	1127	1745	290	318	7			
1	C	211	Total	C	H	N	O	S	0	0	0
			3488	1127	1746	290	318	7			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	SER	-	expression tag	UNP Q6CVQ9
A	-1	ASN	-	expression tag	UNP Q6CVQ9
A	0	ALA	-	expression tag	UNP Q6CVQ9
C	-2	SER	-	expression tag	UNP Q6CVQ9
C	-1	ASN	-	expression tag	UNP Q6CVQ9
C	0	ALA	-	expression tag	UNP Q6CVQ9

- Molecule 2 is a protein called CTF19.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
2	B	197	Total	C	H	N	O	S	0	0	0
			3216	1013	1632	265	299	7			
2	D	198	Total	C	H	N	O	S	0	0	0
			3230	1017	1638	267	301	7			

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total	0	0
			Ca		
3	C	1	Total	0	0
			Ca		
3	D	1	Total	0	0
			Ca		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	13	Total O 13 13	0	0
4	B	6	Total O 6 6	0	0
4	C	18	Total O 18 18	0	0
4	D	11	Total H O 13 2 11	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: MCM21

Chain A:  43% 26% 29%




- Molecule 1: MCM21


Chain C:  43% 25% 29%




- Molecule 2: CTF19

Chain B:  50% 23% 27%





• Molecule 2: CTF19



4 Data and refinement statistics

Property	Value	Source
Space group	P 62 2 2	Depositor
Cell constants a, b, c, α , β , γ	239.04Å 239.04Å 179.49Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	47.49 – 3.70 47.53 – 3.70	Depositor EDS
% Data completeness (in resolution range)	96.7 (47.49-3.70) 96.6 (47.53-3.70)	Depositor EDS
R_{merge}	0.19	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.45 (at 3.67Å)	Xtrriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.183 , 0.209 0.182 , 0.203	Depositor DCC
R_{free} test set	2208 reflections (6.97%)	wwPDB-VP
Wilson B-factor (Å ²)	158.3	Xtrriage
Anisotropy	0.290	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 231.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	13474	wwPDB-VP
Average B, all atoms (Å ²)	168.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.02% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.52	0/1779	0.77	0/2396
1	C	0.53	0/1779	0.79	1/2396 (0.0%)
2	B	0.48	0/1607	0.79	1/2176 (0.0%)
2	D	0.48	0/1615	0.78	0/2187
All	All	0.50	0/6780	0.78	2/9155 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3
1	C	0	4
2	B	0	1
2	D	0	1
All	All	0	9

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	C	117	VAL	CG1-CB-CG2	5.48	119.67	110.90
2	B	190	ARG	NE-CZ-NH1	5.13	122.87	120.30

There are no chirality outliers.

5 of 9 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	145	LEU	Peptide
1	A	163	ARG	Peptide
1	A	57	MET	Peptide
2	B	230	ASN	Peptide
1	C	57	MET	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1742	1745	1741	81	1
1	C	1742	1746	1741	76	1
2	B	1584	1632	1631	65	1
2	D	1592	1638	1637	75	3
3	A	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	13	0	0	3	0
4	B	6	0	0	0	0
4	C	18	0	0	1	0
4	D	11	2	0	0	0
All	All	6711	6763	6750	265	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 20.

The worst 5 of 265 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:54:SER:OG	2:B:149:GLU:OE1	1.89	0.91
2:D:198:MET:SD	2:D:258:ILE:CD1	2.59	0.89
2:D:198:MET:SD	2:D:258:ILE:HD13	2.20	0.81
2:B:72:LEU:HD21	2:B:106:LEU:HD12	1.63	0.80
1:A:49:GLU:O	1:A:118:ARG:NH2	2.15	0.79

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:173:GLU:OE2	2:D:176:GLN:NE2[11_655]	2.03	0.17
1:A:235:LEU:O	2:D:256:TYR:OH[4_665]	2.05	0.15
2:B:256:TYR:OH	1:C:235:LEU:O[7_554]	2.08	0.12
2:D:173:GLU:OE2	2:D:176:GLN:HE21[11_655]	1.59	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	207/296 (70%)	185 (89%)	21 (10%)	1 (0%)	29	66
1	C	207/296 (70%)	184 (89%)	22 (11%)	1 (0%)	29	66
2	B	193/270 (72%)	173 (90%)	20 (10%)	0	100	100
2	D	194/270 (72%)	173 (89%)	21 (11%)	0	100	100
All	All	801/1132 (71%)	715 (89%)	84 (10%)	2 (0%)	47	78

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	291	ILE
1	C	51	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	197/269 (73%)	189 (96%)	8 (4%)	30	59

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	197/269 (73%)	189 (96%)	8 (4%)	30	59
2	B	183/254 (72%)	180 (98%)	3 (2%)	62	80
2	D	184/254 (72%)	180 (98%)	4 (2%)	52	72
All	All	761/1046 (73%)	738 (97%)	23 (3%)	41	66

5 of 23 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	178	ILE
1	C	259	ASP
1	C	212	VAL
1	C	261	ASP
1	A	259	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	172	HIS
1	A	289	GLN
2	B	213	ASN
1	C	172	HIS
2	D	166	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers

Unable to reproduce the depositors R factor - this section is therefore empty.