



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 14, 2022 – 04:09 pm BST

PDB ID : 7ZXF
Title : Pfs48/45 bound to monoclonal antibodies 10D8 and 85RF45.1
Authors : Ko, K.T.; Lennartz, F.L.; Higgins, M.K.
Deposited on : 2022-05-20
Resolution : 3.72 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.28.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0267
CCP4 : 7.1.010 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.28.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.72 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1089 (3.90-3.54)
Clashscore	141614	1012 (3.88-3.56)
Ramachandran outliers	138981	1114 (3.90-3.54)
Sidechain outliers	138945	1110 (3.90-3.54)
RSRZ outliers	127900	1020 (3.90-3.54)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	448	 4% 59% 16% 25%
2	B	445	 2% 37% 9% 54%
3	C	216	 3% 81% 19%
4	D	466	 2% 35% 11% 54%
5	E	240	 4% 75% 14% 10%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
6	F	4	 25% 75%
6	G	4	 50% 50%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 9384 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Gametocyte surface protein P45/48.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	337	2738	1765	432	524	17	0	0	0

- Molecule 2 is a protein called 85RF45.1 heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	206	1573	1002	259	304	8	85	0	0

- Molecule 3 is a protein called 85RF45.1 light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	216	1657	1025	284	339	9	0	0	0

- Molecule 4 is a protein called 10D8 heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	D	214	1626	1023	273	320	10	34	0	0

- Molecule 5 is a protein called 10D8 light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
5	E	217	1690	1061	278	344	7	50	0	0

- Molecule 6 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
6	F	4	Total	C	N	O	0	0	0
			50	28	2	20			
6	G	4	Total	C	N	O	0	0	0
			50	28	2	20			



- Molecule 6: alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G:  50% 50%



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	156.88Å 156.88Å 148.76Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	70.16 – 3.72 70.16 – 3.72	Depositor EDS
% Data completeness (in resolution range)	84.6 (70.16-3.72) 84.6 (70.16-3.72)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.27 (at 3.77Å)	Xtrriage
Refinement program	BUSTER 1.19_4092, PHENIX 1.19_4092	Depositor
R, R_{free}	0.281 , 0.298 0.289 , 0.316	Depositor DCC
R_{free} test set	790 reflections (4.66%)	wwPDB-VP
Wilson B-factor (Å ²)	189.9	Xtrriage
Anisotropy	0.003	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	(Not available) , (Not available)	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	9384	wwPDB-VP
Average B, all atoms (Å ²)	216.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.71% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, NAG, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.25	0/2798	0.47	0/3783
2	B	0.24	0/1614	0.48	0/2206
3	C	0.24	0/1696	0.48	0/2311
4	D	0.24	0/1667	0.49	0/2275
5	E	0.25	0/1731	0.47	0/2350
All	All	0.25	0/9506	0.48	0/12925

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2738	0	2672	44	0
2	B	1573	0	1535	29	0
3	C	1657	0	1578	25	0
4	D	1626	0	1586	34	0
5	E	1690	0	1623	24	0
6	F	50	0	43	0	0
6	G	50	0	43	0	0
All	All	9384	0	9080	149	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 149 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:36:TRP:HB2	3:C:49:ILE:HB	1.69	0.74
1:A:229:LEU:HB2	1:A:232:LYS:HD2	1.70	0.73
4:D:37:VAL:HG12	4:D:47:TRP:HA	1.72	0.71
3:C:155:ASN:HD22	3:C:191:HIS:HB3	1.56	0.70
4:D:148:LEU:HD13	4:D:203:VAL:HG21	1.73	0.69

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	327/448 (73%)	292 (89%)	35 (11%)	0	100	100
2	B	202/445 (45%)	186 (92%)	16 (8%)	0	100	100
3	C	214/216 (99%)	205 (96%)	9 (4%)	0	100	100
4	D	210/466 (45%)	195 (93%)	15 (7%)	0	100	100
5	E	215/240 (90%)	203 (94%)	12 (6%)	0	100	100
All	All	1168/1815 (64%)	1081 (93%)	87 (7%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	319/419 (76%)	314 (98%)	5 (2%)	62	79
2	B	177/400 (44%)	174 (98%)	3 (2%)	60	78
3	C	193/193 (100%)	193 (100%)	0	100	100
4	D	186/417 (45%)	186 (100%)	0	100	100
5	E	194/214 (91%)	190 (98%)	4 (2%)	53	74
All	All	1069/1643 (65%)	1057 (99%)	12 (1%)	73	85

5 of 12 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	104	ASP
5	E	34	ASN
5	E	173	ASP
5	E	44	GLN
1	A	244	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
3	C	82	GLN
3	C	155	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

8 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	NAG	F	1	1,6	14,14,15	0.60	0	17,19,21	0.87	0
6	NAG	F	2	6	14,14,15	1.16	1 (7%)	17,19,21	1.32	2 (11%)
6	BMA	F	3	6	11,11,12	1.18	1 (9%)	15,15,17	1.45	2 (13%)
6	MAN	F	4	6	11,11,12	0.77	0	15,15,17	1.47	2 (13%)
6	NAG	G	1	1,6	14,14,15	1.61	1 (7%)	17,19,21	1.14	2 (11%)
6	NAG	G	2	6	14,14,15	0.24	0	17,19,21	0.67	0
6	BMA	G	3	6	11,11,12	0.78	0	15,15,17	0.88	0
6	MAN	G	4	6	11,11,12	0.62	0	15,15,17	0.95	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	NAG	F	1	1,6	-	2/6/23/26	0/1/1/1
6	NAG	F	2	6	-	3/6/23/26	0/1/1/1
6	BMA	F	3	6	-	1/2/19/22	0/1/1/1
6	MAN	F	4	6	-	0/2/19/22	1/1/1/1
6	NAG	G	1	1,6	-	2/6/23/26	0/1/1/1
6	NAG	G	2	6	-	3/6/23/26	0/1/1/1
6	BMA	G	3	6	-	2/2/19/22	0/1/1/1
6	MAN	G	4	6	-	2/2/19/22	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	G	1	NAG	O5-C1	-5.83	1.34	1.43
6	F	2	NAG	O5-C1	-4.10	1.37	1.43
6	F	3	BMA	C1-C2	2.60	1.58	1.52

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	F	4	MAN	C1-O5-C5	4.54	118.35	112.19
6	F	2	NAG	C4-C3-C2	3.43	116.04	111.02
6	F	3	BMA	C1-C2-C3	-3.22	105.71	109.67
6	F	3	BMA	O5-C1-C2	-2.85	106.37	110.77
6	F	2	NAG	C3-C4-C5	2.48	114.65	110.24

There are no chirality outliers.

5 of 15 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
6	G	3	BMA	O5-C5-C6-O6
6	G	3	BMA	C4-C5-C6-O6
6	G	2	NAG	O5-C5-C6-O6
6	G	2	NAG	C4-C5-C6-O6
6	G	4	MAN	C4-C5-C6-O6


All (1) ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	F	4	MAN	C1-C2-C3-C4-C5-O5

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	337/448 (75%)	0.28	19 (5%) 24 20	158, 209, 264, 280	0
2	B	194/445 (43%)	0.02	8 (4%) 37 31	146, 201, 246, 256	1 (0%)
3	C	216/216 (100%)	0.07	7 (3%) 47 39	177, 223, 253, 260	0
4	D	209/466 (44%)	0.15	11 (5%) 26 23	156, 209, 258, 295	0
5	E	211/240 (87%)	-0.02	10 (4%) 31 26	180, 231, 261, 282	0
All	All	1167/1815 (64%)	0.12	55 (4%) 31 26	146, 217, 257, 295	1 (0%)

The worst 5 of 55 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
5	E	43	GLN	6.4
2	B	184	THR	5.5
5	E	110	LEU	4.4
5	E	88	ASP	4.1
3	C	121	MET	4.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	BMA	F	3	11/12	0.69	0.19	247,252,259,262	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
6	NAG	G	2	14/15	0.82	0.15	226,261,274,277	0
6	BMA	G	3	11/12	0.83	0.13	251,258,266,270	0
6	NAG	G	1	14/15	0.84	0.12	216,245,270,277	0
6	NAG	F	1	14/15	0.86	0.19	231,240,245,249	0
6	NAG	F	2	14/15	0.88	0.18	238,249,256,256	0
6	MAN	G	4	11/12	0.91	0.18	261,265,270,271	0
6	MAN	F	4	11/12	0.95	0.15	253,257,266,269	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.