



wwPDB X-ray Structure Validation Summary Report ⓘ

May 17, 2020 – 07:11 pm BST

PDB ID : 5ZYT
Title : Crystal structure of human MGME1 with 3' overhang double strand DNA3
Authors : Yang, C.; Gan, J.
Deposited on : 2018-05-28
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	324	 44% 23% 32%
1	B	324	 45% 21% 32%
1	C	324	 45% 22% 32%
1	D	324	 47% 20% 32%
2	E	18	 11% 28% 6% 56%
2	F	18	 17% 44% 39%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	G	18	 44% 44% 6% 6%
2	H	18	 17% 28% 6% 50%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 8036 atoms, of which 0 are hydrogens and 0 are deuteriums.


In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Mitochondrial genome maintenance exonuclease 1.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	220	Total 1787	C 1153	N 299	O 326	S 9	0	0	0
1	B	219	Total 1782	C 1150	N 299	O 324	S 9	0	0	0
1	C	220	Total 1802	C 1162	N 303	O 328	S 9	0	0	0
1	D	220	Total 1793	C 1156	N 302	O 326	S 9	0	0	0

- Molecule 2 is a DNA chain called DNA (5'-D(P*CP*TP*TP*CP*TP*TP*CP*C)-3').


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	E	8	Total 141	C 67	N 17	O 49	P 8	0	0	0
2	F	11	Total 216	C 106	N 26	O 73	P 11	0	0	0
2	G	17	Total 336	C 164	N 49	O 107	P 16	0	0	0
2	H	9	Total 179	C 87	N 30	O 54	P 8	0	0	0




• Molecule 1: Mitochondrial genome maintenance exonuclease 1



• Molecule 2: DNA (5'-D(P*CP*TP*TP*CP*TP*TP*CP*C)-3')





• Molecule 2: DNA (5'-D(P*CP*TP*TP*CP*TP*TP*CP*C)-3')



• Molecule 2: DNA (5'-D(P*CP*TP*TP*CP*TP*TP*CP*C)-3')



• Molecule 2: DNA (5'-D(P*CP*TP*TP*CP*TP*TP*CP*C)-3')



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	146.50Å 77.49Å 164.71Å 90.00° 112.40° 90.00°	Depositor
Resolution (Å)	29.96 – 2.70 29.96 – 2.70	Depositor EDS
% Data completeness (in resolution range)	89.4 (29.96-2.70) 73.6 (29.96-2.70)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.86 (at 2.72Å)	Xtrriage
Refinement program	PHENIX (1.10.1_2155: ???)	Depositor
R, R_{free}	0.249 , 0.296 0.250 , 0.299	Depositor DCC
R_{free} test set	2150 reflections (5.11%)	wwPDB-VP
Wilson B-factor (Å ²)	40.9	Xtrriage
Anisotropy	0.141	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 22.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.41$, $\langle L^2 \rangle = 0.24$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.86	EDS
Total number of atoms	8036	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 14.44% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.59	2/1831 (0.1%)	0.71	2/2481 (0.1%)
1	B	0.64	3/1826 (0.2%)	0.75	3/2473 (0.1%)
1	C	0.58	0/1846	0.74	2/2498 (0.1%)
1	D	0.55	1/1837 (0.1%)	0.68	0/2487
2	E	0.98	0/154	1.27	1/235 (0.4%)
2	F	0.93	0/237	1.22	0/362
2	G	1.03	0/371	1.16	1/567 (0.2%)
2	H	1.00	1/199 (0.5%)	1.09	0/305
All	All	0.65	7/8301 (0.1%)	0.80	9/11408 (0.1%)

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	156	MET	CG-SD	6.39	1.97	1.81
1	B	248	CYS	CB-SG	-6.11	1.71	1.82
1	B	279	MET	CG-SD	6.00	1.96	1.81
1	B	309	MET	CG-SD	5.93	1.96	1.81
1	D	276	MET	CG-SD	5.50	1.95	1.81

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	101	PHE	C-N-CD	-9.86	98.90	120.60
1	C	188	SER	C-N-CD	-8.41	102.10	120.60
1	A	237	LEU	CA-CB-CG	7.85	133.36	115.30
2	E	13	DT	O5'-P-OP2	-6.31	100.02	105.70
2	G	8	DT	O4'-C1'-N1	6.19	112.33	108.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1787	0	1766	58	0
1	B	1782	0	1768	62	0
1	C	1802	0	1794	62	0
1	D	1793	0	1778	47	0
2	E	141	0	81	4	0
2	F	216	0	129	10	0
2	G	336	0	198	7	0
2	H	179	0	104	11	0
All	All	8036	0	7618	222	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 222 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:264:SER:OG	2:F:17:DC:O2	1.81	0.98
2:H:1:DG:H5''	2:H:2:DG:OP2	1.68	0.93
2:E:9:DC:H2''	2:E:10:DT:H5''	1.55	0.89
1:B:220:ARG:HB2	1:B:240:VAL:HG23	1.57	0.86
1:B:262:ILE:O	1:B:265:THR:OG1	1.95	0.83

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	214/324 (66%)	206 (96%)	8 (4%)	0	100	100
1	B	213/324 (66%)	206 (97%)	6 (3%)	1 (0%)	29	54
1	C	214/324 (66%)	206 (96%)	7 (3%)	1 (0%)	29	54
1	D	214/324 (66%)	204 (95%)	10 (5%)	0	100	100
All	All	855/1296 (66%)	822 (96%)	31 (4%)	2 (0%)	47	73

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	102	PRO
1	C	106	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	198/294 (67%)	190 (96%)	8 (4%)	31	60
1	B	198/294 (67%)	186 (94%)	12 (6%)	18	41
1	C	201/294 (68%)	194 (96%)	7 (4%)	36	65
1	D	199/294 (68%)	192 (96%)	7 (4%)	36	65
All	All	796/1176 (68%)	762 (96%)	34 (4%)	29	57

5 of 34 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	188	SER
1	B	279	MET
1	D	249	VAL
1	B	227	GLN
1	A	248	CYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 15 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	210	GLN
1	C	228	HIS
1	D	99	ASN
1	B	210	GLN
1	C	316	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	220/324 (67%)	0.12	2 (0%) 84 85	25, 41, 64, 84	0
1	B	219/324 (67%)	0.02	1 (0%) 91 92	28, 42, 62, 72	0
1	C	220/324 (67%)	0.07	2 (0%) 84 85	24, 39, 63, 77	0
1	D	220/324 (67%)	0.13	4 (1%) 68 70	34, 46, 60, 69	0
2	E	8/18 (44%)	0.44	1 (12%) 3 3	37, 61, 72, 94	0
2	F	11/18 (61%)	0.50	2 (18%) 1 1	38, 51, 115, 116	0
2	G	17/18 (94%)	0.46	0 100 100	37, 57, 97, 99	0
2	H	9/18 (50%)	0.01	0 100 100	42, 50, 71, 74	0
All	All	924/1368 (67%)	0.10	12 (1%) 77 78	24, 43, 64, 116	0

The worst 5 of 12 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	106	PRO	3.8
2	F	7	DT	3.1
2	F	8	DC	2.7
1	D	198	GLU	2.6
1	B	329	THR	2.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.