



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 9, 2026 – 08:05 AM EDT

PDB ID : 10BR / pdb_000010br
Title : Crystal structure of B. burgdorferi HtrA PDZ1-2 domains
Authors : Shakya, A.K.; Herzberg, O.; Gallagher, D.T.
Deposited on : 2026-01-09
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4037 (1.50-1.50)
Clashscore	190562	4235 (1.50-1.50)
Ramachandran outliers	187476	4153 (1.50-1.50)
Sidechain outliers	187428	4150 (1.50-1.50)
RSRZ outliers	180081	4039 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	204	 88% 11% .
1	B	204	 87% 9% ..
1	C	204	 85% 12% ..
1	D	204	 89% 11%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 7263 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Periplasmic serine protease DO.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	204	Total	C	N	O	S	0	4	0
			1559	1002	255	298	4			
1	B	199	Total	C	N	O	S	0	3	0
			1522	975	248	295	4			
1	C	199	Total	C	N	O	S	0	0	0
			1514	969	247	294	4			
1	D	204	Total	C	N	O	S	0	1	0
			1549	994	253	298	4			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	271	GLY	-	expression tag	UNP O51131
A	272	PRO	-	expression tag	UNP O51131
A	273	LEU	-	expression tag	UNP O51131
A	274	GLY	-	expression tag	UNP O51131
A	275	SER	-	expression tag	UNP O51131
B	271	GLY	-	expression tag	UNP O51131
B	272	PRO	-	expression tag	UNP O51131
B	273	LEU	-	expression tag	UNP O51131
B	274	GLY	-	expression tag	UNP O51131
B	275	SER	-	expression tag	UNP O51131
C	271	GLY	-	expression tag	UNP O51131
C	272	PRO	-	expression tag	UNP O51131
C	273	LEU	-	expression tag	UNP O51131
C	274	GLY	-	expression tag	UNP O51131
C	275	SER	-	expression tag	UNP O51131
D	271	GLY	-	expression tag	UNP O51131
D	272	PRO	-	expression tag	UNP O51131
D	273	LEU	-	expression tag	UNP O51131
D	274	GLY	-	expression tag	UNP O51131
D	275	SER	-	expression tag	UNP O51131

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	272	Total 272	O 272	0	0
2	B	289	Total 289	O 289	0	0
2	C	259	Total 259	O 259	0	0
2	D	299	Total 299	O 299	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Periplasmic serine protease DO

Chain A:  88% 11%




- Molecule 1: Periplasmic serine protease DO

Chain B:  87% 9%




- Molecule 1: Periplasmic serine protease DO

Chain C:  85% 12%



- Molecule 1: Periplasmic serine protease DO

Chain D:  89% 11%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	38.02Å 72.28Å 76.97Å 65.35° 89.96° 89.19°	Depositor
Resolution (Å)	20.00 – 1.50 20.00 – 1.50	Depositor EDS
% Data completeness (in resolution range)	98.6 (20.00-1.50) 94.4 (20.00-1.50)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.44 (at 1.50Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.182 , 0.230 0.183 , 0.231	Depositor DCC
R_{free} test set	5996 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	19.1	Xtriage
Anisotropy	0.145	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 38.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.178 for h,-k,-l	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	7263	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 19.62% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.73	0/1596	1.21	6/2162 (0.3%)
1	B	0.70	1/1556 (0.1%)	1.17	0/2109
1	C	0.70	0/1538	1.18	3/2086 (0.1%)
1	D	0.75	0/1577	1.27	4/2138 (0.2%)
All	All	0.72	1/6267 (0.0%)	1.21	13/8495 (0.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	276	ILE	N-CA	8.20	1.61	1.46

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	474	PHE	CA-CB-CG	6.54	120.34	113.80
1	A	427	ASN	CA-CB-CG	-6.03	106.57	112.60
1	C	405	GLN	CB-CG-CD	5.89	122.61	112.60
1	C	423	ASN	CB-CA-C	5.86	120.16	110.90
1	D	418	ASP	CA-CB-CG	5.68	118.28	112.60
1	C	291	ARG	CG-CD-NE	-5.60	99.68	112.00
1	A	465	ARG	CB-CG-CD	5.52	123.99	111.30
1	D	428	ILE	CA-C-N	-5.45	113.93	122.65
1	D	428	ILE	C-N-CA	-5.45	113.93	122.65
1	A	421	ASP	CA-CB-CG	5.28	117.88	112.60
1	A	467	ASN	CA-CB-CG	-5.11	107.49	112.60
1	A	316	PRO	N-CD-CG	-5.07	95.60	103.20
1	A	315	TYR	O-C-N	-5.06	116.06	121.42

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1559	0	1582	20	0
1	B	1522	0	1528	18	0
1	C	1514	0	1510	16	0
1	D	1549	0	1559	14	0
2	A	272	0	0	3	0
2	B	289	0	0	3	0
2	C	259	0	0	4	0
2	D	299	0	0	2	0
All	All	7263	0	6179	64	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (64) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:276:ILE:HD12	1:B:276:ILE:N	1.85	0.92
1:D:399:VAL:HG22	1:D:402:ILE:HD12	1.68	0.76
1:B:277:GLU:HG2	1:B:350:SER:HB2	1.71	0.73
1:D:307:SER:HB2	1:D:339:MET:O	1.95	0.67
1:B:440:ASN:O	1:B:441:SER:HB2	1.96	0.66
1:C:424:LEU:O	1:C:428:ILE:HG12	1.96	0.66
1:A:411:TRP:NE1	1:B:405:GLN:HE22	1.93	0.65
1:B:276:ILE:HD13	2:B:547:HOH:O	1.97	0.65
1:C:368:LYS:NZ	2:C:502:HOH:O	2.29	0.65
1:A:271:GLY:N	2:A:501:HOH:O	2.30	0.64
1:A:411:TRP:HE1	1:B:405:GLN:HE22	1.45	0.64
1:D:430:MET:HG2	1:D:463:ILE:HD11	1.79	0.64
1:A:325:LEU:HD23	1:A:371:ILE:HD12	1.81	0.63
1:B:384:LEU:C	1:B:384:LEU:HD13	2.25	0.62
1:C:399:VAL:HG22	1:C:402:ILE:HD12	1.85	0.58
1:D:291:ARG:NH2	1:D:304:ASN:HA	2.21	0.56
1:A:411:TRP:CD1	1:B:405:GLN:HE22	2.22	0.56
1:C:401:ASP:O	1:C:405:GLN:HG3	2.07	0.55
1:D:291:ARG:HH22	1:D:304:ASN:HA	1.72	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:411:TRP:HE1	1:B:405:GLN:NE2	2.05	0.53
1:A:429:LYS:CB	1:A:472:ILE:HG21	2.39	0.53
1:A:274:GLY:HA3	1:A:350:SER:OG	2.09	0.53
1:A:399:VAL:HG22	1:A:402:ILE:HD12	1.90	0.53
1:B:276:ILE:N	1:B:276:ILE:CD1	2.61	0.53
1:B:284:SER:HB3	1:B:313[B]:SER:OG	2.11	0.50
1:C:403:LYS:HE3	2:C:666:HOH:O	2.12	0.49
1:C:456:VAL:HG12	2:C:747:HOH:O	2.12	0.48
1:D:403:LYS:HA	1:D:408:LEU:HD12	1.95	0.48
1:A:402:ILE:HG22	1:A:406:LEU:HD12	1.97	0.47
1:C:426:SER:OG	1:C:427:ASN:ND2	2.47	0.47
1:A:402:ILE:CG2	1:A:406:LEU:HD12	2.45	0.47
1:C:396:TYR:CG	1:C:397:PRO:HD2	2.49	0.46
1:D:274:GLY:HA3	1:D:350:SER:OG	2.15	0.46
1:A:463:ILE:CD1	1:A:463:ILE:N	2.79	0.46
1:A:304:ASN:OD1	1:A:304:ASN:C	2.59	0.45
1:A:456:VAL:HG23	2:A:717:HOH:O	2.18	0.45
1:B:369:LYS:NZ	2:B:509:HOH:O	2.50	0.44
1:C:285:PHE:CG	1:C:342:PHE:HB2	2.52	0.44
1:A:430:MET:HE2	1:A:463:ILE:HD11	1.99	0.43
1:D:402:ILE:HG22	1:D:406:LEU:HD12	2.00	0.43
1:C:399:VAL:HG22	1:C:402:ILE:CD1	2.47	0.43
1:A:463:ILE:N	1:A:463:ILE:HD13	2.34	0.43
1:B:438:SER:HB2	1:B:462:LYS:HB3	2.01	0.43
1:A:304:ASN:OD1	1:A:305:ASP:N	2.51	0.42
1:A:470:PHE:HE2	1:A:472:ILE:CG1	2.33	0.42
1:C:325:LEU:HD22	1:C:362:ILE:HG21	2.01	0.42
1:D:430:MET:HG2	1:D:463:ILE:CD1	2.46	0.42
2:A:687:HOH:O	1:D:388:LYS:HD2	2.18	0.42
1:B:463:ILE:O	1:B:469:SER:HA	2.19	0.42
1:C:424:LEU:HG	1:C:428:ILE:HD13	2.00	0.42
1:A:384:LEU:HD23	1:A:384:LEU:HA	1.92	0.42
1:D:297:LYS:HG2	2:D:770:HOH:O	2.18	0.42
1:B:424:LEU:O	1:B:428:ILE:HG12	2.20	0.42
1:C:384:LEU:HD13	1:C:384:LEU:C	2.45	0.42
1:D:399:VAL:HG22	1:D:402:ILE:CD1	2.44	0.41
1:B:285:PHE:CG	1:B:342:PHE:HB2	2.55	0.41
1:D:310:ILE:HG12	1:D:330:ILE:HG12	2.03	0.41
1:C:390:LEU:HA	1:C:391:PRO:HA	1.90	0.41
1:B:471:LYS:NZ	2:B:514:HOH:O	2.54	0.41
1:C:384:LEU:HD12	2:C:510:HOH:O	2.21	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:312:ALA:HB3	2:D:589:HOH:O	2.21	0.41
1:A:470:PHE:CE2	1:A:472:ILE:CG1	3.04	0.40
1:C:387:SER:HB2	1:C:423:ASN:HB2	2.04	0.40
1:B:396:TYR:CG	1:B:397:PRO:HD2	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	206/204 (101%)	205 (100%)	1 (0%)	0	100	100
1	B	200/204 (98%)	195 (98%)	3 (2%)	2 (1%)	12	2
1	C	197/204 (97%)	193 (98%)	4 (2%)	0	100	100
1	D	203/204 (100%)	201 (99%)	2 (1%)	0	100	100
All	All	806/816 (99%)	794 (98%)	10 (1%)	2 (0%)	36	22

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	277	GLU
1	B	278	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	172/178 (97%)	170 (99%)	2 (1%)	63	38
1	B	169/178 (95%)	167 (99%)	2 (1%)	63	38
1	C	166/178 (93%)	163 (98%)	3 (2%)	51	24
1	D	170/178 (96%)	167 (98%)	3 (2%)	51	24
All	All	677/712 (95%)	667 (98%)	10 (2%)	57	31

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	304	ASN
1	A	463	ILE
1	B	276	ILE
1	B	405	GLN
1	C	276	ILE
1	C	290	THR
1	C	431	LYS
1	D	277	GLU
1	D	307	SER
1	D	409	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	407	ASN
1	B	370	ASN
1	B	405	GLN
1	B	423	ASN
1	C	427	ASN
1	D	410	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	204/204 (100%)	-1.16	0 100 100	8, 20, 39, 63	4 (1%)
1	B	199/204 (97%)	-1.12	0 100 100	12, 23, 41, 70	3 (1%)
1	C	199/204 (97%)	-1.12	0 100 100	14, 23, 46, 62	0
1	D	204/204 (100%)	-1.19	0 100 100	8, 20, 36, 48	1 (0%)
All	All	806/816 (98%)	-1.15	0 100 100	8, 21, 41, 70	8 (0%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.