Full wwPDB X-ray Structure Validation Report ⓘ

Nov 2, 2026 – 10:17 AM JST

PDB ID	:	44IB / pdb_000044ib
Title	:	Crystal structure of human glutaminyl cyclase in complex with Inhibitor LL1260
Authors	:	Li, G.-B.; Chen, Z.-Y.; Li, R.
Deposited on	:	2026-07-30
Resolution	:	1.89 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references ⓘ) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.2
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	2.2
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.51

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.89 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	329	7% 86% 12%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	MG	A	409	-	-	-	X

2 Entry composition


There are 7 unique types of molecules in this entry. The entry contains 5607 atoms, of which 2566 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glutaminyl-peptide cyclotransferase.


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	322	5155	1676	2539	453	478	9	0	1	0

- Molecule 2 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (CCD ID: TRS) (formula: $C_4H_{12}NO_3$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	N	O		
2	A	1	14	4	6	1	3	0	0

- Molecule 3 is {N}-[2-(5-methylimidazol-1-yl)ethyl]-4-(4-methylsulfonylphenyl)-5-(trifluoromethyl)pyrimidin-2-amine (CCD ID: A1FDO) (formula: $C_{18}H_{18}F_3N_5O_2S$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms							ZeroOcc	AltConf
3	A	1	Total	C	F	H	N	O	S	0	0
			47	18	3	18	5	2	1		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	H	O	0	0
			9	3	3	3		

- Molecule 5 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total Zn 1 1	0	0

- Molecule 6 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	5	Total Mg 5 5	0	0

- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	376	Total O 376 376	0	0


3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Glutaminyl-peptide cyclotransferase

Chain A:





4 Data and refinement statistics (i)

Property	Value	Source
Space group	F 41 3 2	Depositor
Cell constants a, b, c, α, β, γ	272.38Å 272.38Å 272.38Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	96.30 – 1.89 96.30 – 1.89	Depositor EDS
% Data completeness (in resolution range)	96.8 (96.30-1.89) 96.8 (96.30-1.89)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.02 (at 1.90Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.171 , 0.195 0.171 , 0.196	Depositor DCC
R_{free} test set	1996 reflections (2.88%)	wwPDB-VP
Wilson B-factor (Å^2)	33.9	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.46 , 62.3	EDS
L-test for twinning[2]	$< \|L\| > = 0.50$, $< L^2 > = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.97	EDS
Total number of atoms	5607	wwPDB-VP
Average B, all atoms (Å^2)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.94% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: TRS, MG, GOL, A1FDO, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#$\|Z\|>5$	RMSZ	#$\|Z\|>5$
1	A	0.55	1/2693 (0.0%)	0.64	0/3663

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	223	MET	SD-CE	-5.37	1.66	1.79

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2616	2539	2536	26	1
2	A	8	6	12	2	0
3	A	29	18	0	0	0
4	A	6	3	8	1	0
5	A	1	0	0	0	0
6	A	5	0	0	0	0
7	A	376	0	0	11	7
All	All	3041	2566	2556	28	7

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 5.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:401:TRS:H12	7:A:781:HOH:O	1.67	0.95
1:A:327:GLU:OE2	7:A:501:HOH:O	1.96	0.83
1:A:176:LYS:NZ	7:A:502:HOH:O	2.05	0.78
1:A:92:MET:SD	1:A:106:ILE:HD11	2.31	0.70
1:A:301:GLY:N	7:A:512:HOH:O	2.32	0.61
1:A:101:ASP:OD1	7:A:504:HOH:O	2.16	0.59
1:A:44:GLN:OE1	7:A:505:HOH:O	2.16	0.59
1:A:88:ARG:HG2	1:A:92:MET:HE2	1.90	0.54
1:A:177:LYS:O	1:A:180:SER:HB3	2.09	0.52
1:A:297:TYR:OH	7:A:503:HOH:O	2.15	0.52
1:A:89:GLN:O	1:A:93:GLN:HG3	2.10	0.51
1:A:36:TRP:N	1:A:37:PRO:CD	2.74	0.50
1:A:35:ALA:O	1:A:38:GLU:HG2	2.13	0.49
1:A:295:GLN:HB3	1:A:297:TYR:CE2	2.47	0.49
1:A:205:LEU:O	1:A:206:HIS:CD2	2.65	0.49
2:A:401:TRS:O3	7:A:506:HOH:O	2.19	0.48
1:A:205:LEU:O	1:A:206:HIS:HD2	1.96	0.48
1:A:238:LEU:HD12	1:A:241:MET:SD	2.54	0.47
1:A:41:ASN:HB2	7:A:573:HOH:O	2.15	0.46
1:A:224:ALA:HB2	1:A:238:LEU:HD22	1.98	0.45
1:A:233:ARG:NH2	7:A:525:HOH:O	2.48	0.45
1:A:94:ARG:O	1:A:97:ARG:HG2	2.16	0.45
1:A:327:GLU:HG3	7:A:680:HOH:O	2.16	0.44
1:A:308:ILE:HB	1:A:309:PRO:HD3	2.00	0.43
1:A:74:LEU:HD13	1:A:336:GLU:HB2	2.01	0.43
1:A:109:PHE:CZ	1:A:120:PHE:HB2	2.55	0.42
1:A:270:ARG:HH12	4:A:403:GOL:H31	1.84	0.41
1:A:76:GLU:O	1:A:83:GLY:HA3	2.21	0.41

All (7) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:A:755:HOH:O	7:A:846:HOH:O[59_555]	1.88	0.32
7:A:573:HOH:O	7:A:636:HOH:O[80_555]	1.98	0.22
1:A:269:GLU:OE1	7:A:501:HOH:O[80_555]	1.99	0.21
7:A:737:HOH:O	7:A:802:HOH:O[9_555]	2.02	0.18

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:A:501:HOH:O	7:A:625:HOH:O[59_555]	2.08	0.12
7:A:697:HOH:O	7:A:795:HOH:O[80_555]	2.12	0.08
7:A:524:HOH:O	7:A:671:HOH:O[59_555]	2.18	0.02

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	319/329 (97%)	310 (97%)	8 (2%)	1 (0%)	36 29

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	212	SER

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	285/290 (98%)	283 (99%)	2 (1%)	76 78

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	180	SER
1	A	242	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	67	GLN
1	A	68	ASN
1	A	89	GLN
1	A	99	GLN
1	A	150	ASN
1	A	194	GLN
1	A	206	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

Of 9 ligands modelled in this entry, 6 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
4	GOL	A	403	-	5,5,5	0.97	0	5,5,5	0.81	0
3	A1FDO	A	402	5	31,31,31	2.65	7 (22%)	41,46,46	3.56	21 (51%)
2	TRS	A	401	-	7,7,7	0.22	0	9,9,9	0.90	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	403	-	-	2/4/4/4	-
3	A1FDO	A	402	5	-	8/22/22/22	0/3/3/3
2	TRS	A	401	-	-	0/9/9/9	-

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	402	A1FDO	C10-N09	9.33	1.48	1.34
3	A	402	A1FDO	C22-S23	8.12	1.85	1.77
3	A	402	A1FDO	C19-C18	4.46	1.54	1.49
3	A	402	A1FDO	C02-N06	-3.36	1.32	1.38
3	A	402	A1FDO	C14-C13	2.71	1.56	1.50
3	A	402	A1FDO	C12-C13	2.07	1.41	1.38
3	A	402	A1FDO	C24-S23	2.01	1.83	1.75

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
3	A	402	A1FDO	C08-C07-N06	9.41	123.84	112.57
3	A	402	A1FDO	O26-S23-O25	-7.57	104.89	117.92
3	A	402	A1FDO	O25-S23-C22	6.97	113.93	108.25
3	A	402	A1FDO	N11-C10-N29	-6.19	120.68	126.55
3	A	402	A1FDO	N06-C05-N04	-5.41	108.13	113.51
3	A	402	A1FDO	C07-N06-C05	-5.15	118.44	126.64
3	A	402	A1FDO	C20-C19-C18	-4.79	113.02	120.61
3	A	402	A1FDO	C24-S23-C22	4.67	110.10	104.58
3	A	402	A1FDO	C19-C18-C13	4.25	128.73	123.69
3	A	402	A1FDO	C01-C02-N06	4.21	127.84	122.80
3	A	402	A1FDO	C27-C22-S23	4.11	123.39	119.58
3	A	402	A1FDO	C13-C12-N11	-3.77	118.97	123.95
3	A	402	A1FDO	C12-N11-C10	3.76	121.55	115.88
3	A	402	A1FDO	C08-N09-C10	3.73	130.19	123.75
3	A	402	A1FDO	C13-C18-N29	-3.31	119.45	122.78
3	A	402	A1FDO	C28-C19-C18	3.18	125.65	120.61
3	A	402	A1FDO	O26-S23-C24	3.10	112.94	108.49
3	A	402	A1FDO	C19-C18-N29	-3.05	111.03	115.09
3	A	402	A1FDO	C05-N06-C02	2.96	110.10	106.37
3	A	402	A1FDO	C21-C22-S23	-2.92	116.88	119.58
3	A	402	A1FDO	F16-C14-C13	-2.74	107.93	112.70

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	402	A1FDO	N06-C07-C08-N09
3	A	402	A1FDO	C08-C07-N06-C02
3	A	402	A1FDO	C08-C07-N06-C05
3	A	402	A1FDO	N11-C10-N09-C08
3	A	402	A1FDO	N29-C10-N09-C08
4	A	403	GOL	O1-C1-C2-C3
3	A	402	A1FDO	C21-C22-S23-O25
4	A	403	GOL	O1-C1-C2-O2
3	A	402	A1FDO	C27-C22-S23-O25
3	A	402	A1FDO	C21-C22-S23-C24


There are no ring outliers.

2 monomers are involved in 3 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	403	GOL	1	0
2	A	401	TRS	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand A1FDO A 402




Bond lengths



Bond angles



Torsions



Rings

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	322/329 (97%)	0.03	24 (7%)	20	21	20, 38, 59, 78	1 (0%)

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	189	PRO	6.4
1	A	149	TRP	5.9
1	A	117	TYR	4.4
1	A	312[A]	ARG	4.1
1	A	34	SER	4.1
1	A	35	ALA	4.0
1	A	182	LYS	3.9
1	A	150	ASN	3.5
1	A	300	GLY	3.4
1	A	233	ARG	2.9
1	A	296	ASN	2.7
1	A	302	VAL	2.7
1	A	301	GLY	2.6
1	A	297	TYR	2.5
1	A	290	GLU	2.5
1	A	41	ASN	2.5
1	A	210	GLN	2.4
1	A	176	LYS	2.4
1	A	206	HIS	2.3
1	A	42	TYR	2.3
1	A	97	ARG	2.3
1	A	181	LEU	2.2
1	A	179	LEU	2.2
1	A	58	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no oligosaccharides in this entry.

6.4 Ligands (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
6	MG	A	409	1/1	0.67	0.41	75,75,75,75	0
6	MG	A	407	1/1	0.78	0.22	65,65,65,65	0
2	TRS	A	401	8/8	0.82	0.21	48,66,78,87	0
6	MG	A	408	1/1	0.88	0.18	73,73,73,73	0
6	MG	A	405	1/1	0.88	0.19	59,59,59,59	0
6	MG	A	406	1/1	0.90	0.18	62,62,62,62	0
4	GOL	A	403	6/6	0.90	0.18	48,64,77,79	0
3	A1FDO	A	402	29/29	0.93	0.13	28,44,67,77	0
5	ZN	A	404	1/1	1.00	0.01	28,28,28,28	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around A1FDO A 402:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.