Full wwPDB X-ray Structure Validation Report (i)

Nov 4, 2026 – 02:12 PM JST

PDB ID	:	44IY / pdb_000044iy
Title	:	KIR2DS4 bound to HLA-C*05:01 presenting the HECTD1 peptide
Authors	:	Faoro, C.; Rossjohn, J.
Deposited on	:	2026-07-31
Resolution	:	2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.2
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	2.2
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.51

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3538 (2.70-2.70)
Clashscore	190562	3843 (2.70-2.70)
Ramachandran outliers	187476	3778 (2.70-2.70)
Sidechain outliers	187428	3778 (2.70-2.70)
RSRZ outliers	180081	3538 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	279	3% 89% 10%
1	E	279	2% 84% 14%
2	B	100	4% 84% 16%
2	F	100	3% 87% 12%
3	C	9	11% 100%
3	G	9	11% 67% 33%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	D	198	9% 86% 13%
4	H	198	19% 87% 13%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	SO4	D	302	-	-	X	-

2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 9498 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HLA class I histocompatibility antigen, C alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	275	Total	C	N	O	S	0	0	0
			2250	1403	414	426	7			
1	E	275	Total	C	N	O	S	0	1	0
			2258	1408	417	426	7			

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	initiating methionine	UNP P10321
A	1	MET	-	expression tag	UNP P10321
A	9	TYR	ASP	variant	UNP P10321
A	24	ALA	SER	variant	UNP P10321
A	35	GLN	ARG	variant	UNP P10321
A	73	THR	ALA	variant	UNP P10321
A	77	ASN	SER	variant	UNP P10321
A	80	LYS	ASN	variant	UNP P10321
A	90	ALA	ASP	variant	UNP P10321
A	99	TYR	SER	variant	UNP P10321
A	114	ASN	ASP	variant	UNP P10321
A	116	PHE	SER	variant	UNP P10321
A	138	LYS	THR	variant	UNP P10321
A	147	TRP	LEU	variant	UNP P10321
A	152	GLU	ALA	variant	UNP P10321
A	156	ARG	LEU	variant	UNP P10321
A	177	LYS	GLU	variant	UNP P10321
A	184	HIS	PRO	variant	UNP P10321
A	194	VAL	LEU	variant	UNP P10321
A	253	GLU	GLN	variant	UNP P10321
A	261	VAL	MET	variant	UNP P10321
A	267	PRO	GLN	variant	UNP P10321
A	273	ARG	SER	variant	UNP P10321
A	275	GLY	GLU	variant	UNP P10321
E	0	MET	-	initiating methionine	UNP P10321

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	1	MET	-	expression tag	UNP P10321
E	9	TYR	ASP	variant	UNP P10321
E	24	ALA	SER	variant	UNP P10321
E	35	GLN	ARG	variant	UNP P10321
E	73	THR	ALA	variant	UNP P10321
E	77	ASN	SER	variant	UNP P10321
E	80	LYS	ASN	variant	UNP P10321
E	90	ALA	ASP	variant	UNP P10321
E	99	TYR	SER	variant	UNP P10321
E	114	ASN	ASP	variant	UNP P10321
E	116	PHE	SER	variant	UNP P10321
E	138	LYS	THR	variant	UNP P10321
E	147	TRP	LEU	variant	UNP P10321
E	152	GLU	ALA	variant	UNP P10321
E	156	ARG	LEU	variant	UNP P10321
E	177	LYS	GLU	variant	UNP P10321
E	184	HIS	PRO	variant	UNP P10321
E	194	VAL	LEU	variant	UNP P10321
E	253	GLU	GLN	variant	UNP P10321
E	261	VAL	MET	variant	UNP P10321
E	267	PRO	GLN	variant	UNP P10321
E	273	ARG	SER	variant	UNP P10321
E	275	GLY	GLU	variant	UNP P10321

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	100	Total	C	N	O	S	0	0	0
			826	526	140	157	3			
2	F	100	Total	C	N	O	S	0	0	0
			822	523	139	157	3			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	MET	-	initiating methionine	UNP P61769
F	0	MET	-	initiating methionine	UNP P61769


- Molecule 3 is a protein called HECTD1 peptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	9	Total	C	N	O	0	0	0
			78	48	13	17			
3	G	9	Total	C	N	O	0	0	0
			78	48	13	17			

- Molecule 4 is a protein called Killer cell immunoglobulin-like receptor 2DS4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	H	198	Total	C	N	O	S	0	1	0
			1505	953	260	282	10			
4	D	198	Total	C	N	O	S	0	2	0
			1545	978	269	288	10			

- Molecule 5 is SULFATE ION (CCD ID: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	D	1	Total	O	S	0	0
			5	4	1		
5	D	1	Total	O	S	0	0
			5	4	1		
5	D	1	Total	O	S	0	0
			5	4	1		
5	D	1	Total	O	S	0	0
			5	4	1		
5	D	1	Total	O	S	0	0
			5	4	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	28	Total O 28 28	0	0
6	B	11	Total O 11 11	0	0
6	E	20	Total O 20 20	0	0
6	F	8	Total O 8 8	0	0
6	G	2	Total O 2 2	0	0
6	H	19	Total O 19 19	0	0
6	D	23	Total O 23 23	0	0


3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: HLA class I histocompatibility antigen, C alpha chain




• Molecule 1: HLA class I histocompatibility antigen, C alpha chain



• Molecule 2: Beta-2-microglobulin




• Molecule 2: Beta-2-microglobulin




• Molecule 3: HECTD1 peptide

Chain C:




• Molecule 3: HECTD1 peptide

Chain G:




• Molecule 4: Killer cell immunoglobulin-like receptor 2DS4

Chain H:



• Molecule 4: Killer cell immunoglobulin-like receptor 2DS4

Chain D:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α, β, γ	69.75Å 96.40Å 230.82Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.18 – 2.70 47.18 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.7 (47.18-2.70) 99.7 (47.18-2.70)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	2.26 (at 2.69Å)	Xtriage
Refinement program	PHENIX (1.21rc1_5058: ???)	Depositor
R, R_{free}	0.210 , 0.254 0.209 , 0.251	Depositor DCC
R_{free} test set	2219 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å^2)	45.4	Xtriage
Anisotropy	0.660	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.32 , 35.9	EDS
L-test for twinning[2]	$< \|L\| > = 0.50$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.94	EDS
Total number of atoms	9498	wwPDB-VP
Average B, all atoms (Å^2)	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 42.67 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.9719e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| > 5$	RMSZ	$\#\|Z\| > 5$
1	A	0.09	0/2312	0.25	0/3138
1	E	0.09	0/2323	0.25	0/3152
2	B	0.09	0/849	0.26	0/1149
2	F	0.08	0/845	0.27	0/1145
3	C	0.08	0/80	0.23	0/106
3	G	0.07	0/80	0.25	0/106
4	D	0.12	0/1595	0.31	0/2169
4	H	0.11	0/1552	0.30	0/2113
All	All	0.10	0/9636	0.27	0/13078

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2250	0	2109	15	0
1	E	2258	0	2122	21	0
2	B	826	0	782	9	0
2	F	822	0	771	8	0
3	C	78	0	64	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	G	78	0	64	2	0
4	D	1545	0	1477	16	0
4	H	1505	0	1429	11	0
5	D	25	0	0	2	0
6	A	28	0	0	0	0
6	B	11	0	0	0	0
6	D	23	0	0	0	0
6	E	20	0	0	0	0
6	F	8	0	0	0	0
6	G	2	0	0	0	0
6	H	19	0	0	0	0
All	All	9498	0	8818	77	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (77) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:H:131:ARG:NH1	4:H:157:ASN:O	2.23	0.71
4:D:41[B]:ARG:NH1	4:D:45:PHE:O	2.30	0.65
2:B:36:GLU:HB2	2:B:83:ASN:HB3	1.78	0.64
1:E:14:ARG:NH1	1:E:39:ASP:OD2	2.31	0.64
2:F:36:GLU:HB2	2:F:83:ASN:HB3	1.80	0.63
1:A:14:ARG:NH1	1:A:39:ASP:OD2	2.32	0.62
1:E:44:ARG:HD2	1:E:64:THR:HG21	1.80	0.62
4:H:83:VAL:HG13	4:H:84:PRO:HD3	1.82	0.61
1:E:220:ASP:OD2	1:E:256:ARG:NH2	2.28	0.61
2:F:29:GLY:HA2	2:F:61:SER:HB3	1.85	0.58
2:B:7:ILE:HG12	2:B:82:VAL:HG21	1.86	0.57
1:A:15:PRO:HG3	1:A:91:GLY:H	1.69	0.56
4:D:24:VAL:HG11	4:D:99:MET:HE2	1.86	0.56
1:E:9:TYR:HB2	1:E:97:ARG:HB3	1.88	0.55
4:H:24:VAL:HG11	4:H:99:MET:HE2	1.89	0.55
4:D:149:ARG:NH2	5:D:302:SO4:O3	2.34	0.54
4:H:100:VAL:HG11	4:H:188:TRP:CD2	2.43	0.54
2:B:73:THR:OG1	2:B:76:ASP:OD2	2.25	0.53
1:E:15:PRO:HG2	1:E:91:GLY:H	1.73	0.52
4:D:100:VAL:HG11	4:D:188:TRP:CD2	2.45	0.52
4:D:72:VAL:HG13	4:D:73:LEU:HD22	1.91	0.52
2:B:29:GLY:HA2	2:B:61:SER:HB3	1.92	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:230:LEU:HD22	1:E:243:LYS:HE3	1.93	0.51
4:D:36:HIS:NE2	4:D:85:HIS:HB2	2.25	0.51
1:A:103:LEU:HD13	1:A:168:LEU:HD23	1.92	0.51
4:D:149:ARG:NH2	5:D:302:SO4:O4	2.44	0.51
1:A:214:THR:HB	1:A:262:GLN:HB2	1.93	0.50
1:E:55:GLU:HG2	1:E:59:TYR:CG	2.47	0.49
1:A:201:LEU:HD12	1:A:249:VAL:HG21	1.95	0.49
1:E:201:LEU:HD12	1:E:249:VAL:HG21	1.95	0.49
2:F:7:ILE:HD12	2:F:91:LYS:HE3	1.95	0.49
1:A:44:ARG:HD2	1:A:64:THR:HG21	1.94	0.49
4:H:14:PRO:HD2	4:H:24:VAL:HG23	1.96	0.48
4:H:84:PRO:HB2	4:H:87:PRO:HD2	1.95	0.48
4:H:72:VAL:HG13	4:H:73:LEU:HD13	1.96	0.48
1:E:35:GLN:HG3	2:F:53:ASP:CG	2.39	0.48
1:A:9:TYR:HB2	1:A:97:ARG:HB3	1.95	0.47
4:D:126:LEU:HD12	4:D:139:LEU:HD21	1.95	0.47
4:H:183:ASP:OD1	4:H:184:ALA:N	2.47	0.47
1:E:10:THR:HG21	2:F:62:PHE:HE1	1.81	0.46
1:A:55:GLU:HG2	1:A:59:TYR:CG	2.51	0.46
1:A:263:HIS:HB3	1:A:266:LEU:HD22	1.98	0.46
2:B:22:PHE:CE2	2:B:69:GLU:HG2	2.51	0.46
1:E:177:LYS:HD2	1:E:177:LYS:HA	1.71	0.45
4:D:163:ASP:O	4:D:165:PRO:HD3	2.17	0.45
1:A:162:GLY:O	1:A:166:GLU:HG2	2.16	0.45
2:B:25:CYS:HB2	2:B:39:LEU:HD21	1.97	0.45
4:D:83:VAL:HB	4:D:84:PRO:HD3	1.98	0.45
4:H:86:SER:HB3	4:H:87:PRO:HD3	1.98	0.44
4:D:41[B]:ARG:HH12	4:D:45:PHE:H	1.65	0.44
4:H:53:GLY:HA2	4:H:61:LYS:O	2.16	0.44
1:A:117:ALA:HB2	2:B:60:TRP:CE2	2.53	0.44
1:E:103:LEU:HD13	1:E:168:LEU:HD23	2.00	0.44
2:F:22:PHE:CE2	2:F:69:GLU:HG2	2.53	0.44
1:E:124:ILE:HD11	1:E:133:TRP:HB3	1.98	0.44
1:E:117:ALA:HB2	2:F:60:TRP:CE2	2.53	0.43
1:A:35:GLN:HG3	2:B:53:ASP:CG	2.44	0.43
1:E:162:GLY:O	1:E:166:GLU:HG2	2.19	0.42
2:B:17:ASN:HA	2:B:72:PRO:O	2.18	0.42
2:F:1:ILE:HD13	2:F:2:GLN:N	2.34	0.42
3:G:4:ASP:OD1	3:G:5:LYS:N	2.46	0.42
1:E:5:MET:HB2	1:E:168:LEU:HD13	2.01	0.42
1:E:199:ALA:N	1:E:249:VAL:O	2.52	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:230:LEU:HD22	1:A:243:LYS:HE3	2.01	0.41
1:E:0:MET:HA	1:E:105:PRO:HB3	2.02	0.41
1:E:196:ASP:OD1	1:E:196:ASP:N	2.53	0.41
4:H:91:SER:OG	4:H:92:ALA:N	2.53	0.41
4:D:102:ILE:HG22	4:D:188:TRP:HB2	2.02	0.41
1:A:185:PRO:HB3	1:A:208:PHE:HB3	2.01	0.41
1:E:187:THR:HB	1:E:272:LEU:HD11	2.01	0.41
1:E:21:ARG:HE	1:E:39:ASP:HB2	1.86	0.41
4:D:4:VAL:HB	4:D:5:HIS:H	1.71	0.41
1:A:196:ASP:OD1	1:A:196:ASP:N	2.51	0.41
4:D:23:THR:HG23	4:D:67:GLY:HA3	2.02	0.41
4:D:100:VAL:HG11	4:D:188:TRP:CE2	2.55	0.41
3:G:3:ASP:OD1	3:G:4:ASP:N	2.52	0.41
4:D:176:ARG:CZ	4:D:194:PRO:HG3	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	273/279 (98%)	262 (96%)	11 (4%)	0	100	100
1	E	274/279 (98%)	264 (96%)	10 (4%)	0	100	100
2	B	98/100 (98%)	97 (99%)	1 (1%)	0	100	100
2	F	98/100 (98%)	98 (100%)	0	0	100	100
3	C	7/9 (78%)	7 (100%)	0	0	100	100
3	G	7/9 (78%)	7 (100%)	0	0	100	100
4	D	198/198 (100%)	191 (96%)	7 (4%)	0	100	100
4	H	197/198 (100%)	186 (94%)	11 (6%)	0	100	100
All	All	1152/1172 (98%)	1112 (96%)	40 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	229/234 (98%)	227 (99%)	2 (1%)	70 87
1	E	230/234 (98%)	222 (96%)	8 (4%)	32 61
2	B	92/95 (97%)	92 (100%)	0	100 100
2	F	91/95 (96%)	90 (99%)	1 (1%)	65 85
3	C	8/8 (100%)	8 (100%)	0	100 100
3	G	8/8 (100%)	8 (100%)	0	100 100
4	D	171/170 (101%)	166 (97%)	5 (3%)	37 67
4	H	163/170 (96%)	155 (95%)	8 (5%)	22 49
All	All	992/1014 (98%)	968 (98%)	24 (2%)	43 72

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	76	VAL
1	A	266	LEU
1	E	10	THR
1	E	76	VAL
1	E	156	ARG
1	E	177	LYS
1	E	194	VAL
1	E	202	ARG
1	E	225	THR
1	E	258	THR
2	F	1	ILE
4	H	4	VAL
4	H	5	HIS
4	H	33	MET
4	H	42	GLU
4	H	47	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
4	H	51	LEU
4	H	83	VAL
4	H	113	GLN
4	D	4	VAL
4	D	57	ASP
4	D	59	VAL
4	D	73	LEU
4	D	157	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	54	GLN
2	B	13	HIS
2	B	31	HIS
1	E	54	GLN
1	E	72	GLN
1	E	224	GLN
2	F	13	HIS
2	F	17	ASN
2	F	31	HIS
4	H	113	GLN
4	H	123	ASN
4	D	27	GLN
4	D	40	HIS
4	D	46	ASN
4	D	63	ASN
4	D	89	GLN

5.3.3 RNA

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
5	SO4	D	302	-	4,4,4	0.60	0	6,6,6	0.06	0
5	SO4	D	305	-	4,4,4	0.59	0	6,6,6	0.08	0
5	SO4	D	303	-	4,4,4	0.59	0	6,6,6	0.04	0
5	SO4	D	304	-	4,4,4	0.60	0	6,6,6	0.08	0
5	SO4	D	301	-	4,4,4	0.60	0	6,6,6	0.05	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	D	302	SO4	2	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	275/279 (98%)	0.31	7 (2%)	58	55	29, 50, 93, 129	0
1	E	275/279 (98%)	0.41	6 (2%)	62	59	31, 52, 92, 121	1 (0%)
2	B	100/100 (100%)	0.34	4 (4%)	42	38	31, 49, 83, 109	0
2	F	100/100 (100%)	0.34	3 (3%)	52	49	35, 52, 87, 116	0
3	C	9/9 (100%)	0.10	1 (11%)	10	8	38, 41, 45, 48	0
3	G	9/9 (100%)	0.61	1 (11%)	10	8	42, 47, 54, 55	0
4	D	198/198 (100%)	0.53	18 (9%)	15	12	27, 50, 91, 142	2 (1%)
4	H	198/198 (100%)	1.00	38 (19%)	3	3	28, 60, 130, 156	1 (0%)
All	All	1164/1172 (99%)	0.49	78 (6%)	24	21	27, 52, 103, 156	4 (0%)

All (78) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	H	45	PHE	5.4
4	H	54	GLU	5.0
4	D	45	PHE	4.9
4	H	47	ASN	4.1
4	H	52	ILE	4.1
4	D	4	VAL	4.1
4	D	143	GLY	3.9
4	H	83	VAL	3.8
2	B	1	ILE	3.7
1	A	0	MET	3.7
4	H	57	ASP	3.7
4	H	44	LYS	3.7
1	E	74	ASP	3.6
4	H	34	PHE	3.5
3	G	9	PHE	3.5
4	H	36	HIS	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
4	H	81	GLY	3.4
4	H	37	PHE	3.4
2	F	1	ILE	3.3
4	D	79	CYS	3.2
4	D	85	HIS	3.2
4	H	86	SER	3.0
4	H	88	TYR	3.0
4	H	51	LEU	2.9
4	H	5	HIS	2.9
4	H	87	PRO	2.9
4	H	85	HIS	2.9
2	F	80	CYS	2.9
1	A	1	MET	2.8
2	B	0	MET	2.8
4	H	58	GLY	2.8
2	F	0	MET	2.8
4	H	201	GLY	2.7
4	H	56	HIS	2.7
1	A	54	GLN	2.7
1	E	54	GLN	2.7
4	H	49	LEU	2.6
4	H	4	VAL	2.6
4	H	28	CYS	2.6
4	D	42	GLU	2.6
1	A	41	ALA	2.6
4	H	11	LEU	2.6
4	H	55	HIS	2.6
4	H	38	LEU	2.5
2	B	48	LYS	2.5
4	H	64	PHE	2.5
4	H	62	ALA	2.5
4	D	44	LYS	2.5
1	E	17	ARG	2.5
4	H	53	GLY	2.5
4	H	89	GLN	2.5
4	D	201	GLY	2.4
4	H	6	ARG	2.4
4	D	43	GLY	2.3
4	H	59	VAL	2.3
1	A	262	GLN	2.3
4	D	11	LEU	2.3
4	H	50	HIS	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	E	19	GLU	2.2
4	H	29	TRP	2.2
4	H	61	LYS	2.2
4	D	115	GLY	2.2
1	A	194	VAL	2.2
1	E	264	GLU	2.2
4	D	47	ASN	2.2
4	D	48	THR	2.1
4	D	5	HIS	2.1
4	D	156	ILE	2.1
4	D	87	PRO	2.1
4	H	32	VAL	2.1
2	B	74	GLU	2.1
4	H	128	CYS	2.1
3	C	9	PHE	2.1
1	A	42	SER	2.0
4	H	92	ALA	2.0
1	E	252	GLY	2.0
4	D	86	SER	2.0
4	D	41[A]	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no oligosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
5	SO4	D	305	5/5	0.82	0.13	67,69,89,106	0
5	SO4	D	303	5/5	0.86	0.11	68,71,95,120	0
5	SO4	D	301	5/5	0.90	0.14	74,82,96,108	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
5	SO4	D	304	5/5	0.93	0.09	61,65,68,83	0
5	SO4	D	302	5/5	0.94	0.09	64,70,81,82	0

6.5 Other polymers (i)

There are no such residues in this entry.