



wwPDB EM Validation Summary Report ⓘ

Sep 24, 2026 – 02:24 PM EDT

PDB ID : 38RH / pdb_000038rh
EMDB ID : EMD-79079
Title : Structure of the DAB1 Complex in State A
Authors : Phillips, N.R.; Oltrogge, L.M.; Remis, J.P.; Savage, D.F.
Deposited on : 2026-09-15
Resolution : 2.29 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:



EMDB validation analysis : 0.0.1.dev133
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.50

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 2.29 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	3699 (1.79 - 2.79)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1031	
2	B	773	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 27007 atoms, of which 13221 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Probable inorganic carbon transporter subunit DabA.

Mol	Chain	Residues	Atoms						AltConf	Trace
1	A	907	Total	C	H	N	O	S	0	0
			14426	4740	7164	1208	1291	23		

There are 32 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1000	GLY	-	expression tag	UNP D3SP01
A	1001	GLY	-	expression tag	UNP D3SP01
A	1002	SER	-	expression tag	UNP D3SP01
A	1003	GLY	-	expression tag	UNP D3SP01
A	1004	TRP	-	expression tag	UNP D3SP01
A	1005	SER	-	expression tag	UNP D3SP01
A	1006	HIS	-	expression tag	UNP D3SP01
A	1007	PRO	-	expression tag	UNP D3SP01
A	1008	GLN	-	expression tag	UNP D3SP01
A	1009	PHE	-	expression tag	UNP D3SP01
A	1010	GLU	-	expression tag	UNP D3SP01
A	1011	LYS	-	expression tag	UNP D3SP01
A	1012	GLY	-	expression tag	UNP D3SP01
A	1013	GLY	-	expression tag	UNP D3SP01
A	1014	GLY	-	expression tag	UNP D3SP01
A	1015	SER	-	expression tag	UNP D3SP01
A	1016	GLY	-	expression tag	UNP D3SP01
A	1017	GLY	-	expression tag	UNP D3SP01
A	1018	GLY	-	expression tag	UNP D3SP01
A	1019	SER	-	expression tag	UNP D3SP01
A	1020	GLY	-	expression tag	UNP D3SP01
A	1021	GLY	-	expression tag	UNP D3SP01
A	1022	SER	-	expression tag	UNP D3SP01
A	1023	ALA	-	expression tag	UNP D3SP01
A	1024	TRP	-	expression tag	UNP D3SP01
A	1025	SER	-	expression tag	UNP D3SP01
A	1026	HIS	-	expression tag	UNP D3SP01
A	1027	PRO	-	expression tag	UNP D3SP01

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	1028	GLN	-	expression tag	UNP D3SP01
A	1029	PHE	-	expression tag	UNP D3SP01
A	1030	GLU	-	expression tag	UNP D3SP01
A	1031	LYS	-	expression tag	UNP D3SP01

- Molecule 2 is a protein called NADH/Ubiquinone/plastoquinone (Complex I).

Mol	Chain	Residues	Atoms					AltConf	Trace
2	B	740	Total	C	H	N	O	S	
			11934	3964	6057	912	976	25	0
									0

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	760	SER	-	expression tag	UNP D3SP02
B	761	GLY	-	expression tag	UNP D3SP02
B	762	SER	-	expression tag	UNP D3SP02
B	763	SER	-	expression tag	UNP D3SP02
B	764	ALA	-	expression tag	UNP D3SP02
B	765	HIS	-	expression tag	UNP D3SP02
B	766	HIS	-	expression tag	UNP D3SP02
B	767	HIS	-	expression tag	UNP D3SP02
B	768	HIS	-	expression tag	UNP D3SP02
B	769	HIS	-	expression tag	UNP D3SP02
B	770	HIS	-	expression tag	UNP D3SP02
B	771	HIS	-	expression tag	UNP D3SP02
B	772	HIS	-	expression tag	UNP D3SP02
B	773	HIS	-	expression tag	UNP D3SP02

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		AltConf
3	A	1	Total	Zn	
			1	1	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		AltConf
4	A	421	Total	O	
			421	421	0
4	B	225	Total	O	
			225	225	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Probable inorganic carbon transporter subunit DabA



• Molecule 2: NADH/Ubiquinone/plastoquinone (Complex I)





4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	164827	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION; Patch CTF used in cryoSPARC	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	50	Depositor
Minimum defocus (nm)	500	Depositor
Maximum defocus (nm)	1800	Depositor
Magnification	Not provided	
Image detector	GATAN K3 BIOQUANTUM (6k x 4k)	Depositor
Maximum map value	4.671	Depositor
Minimum map value	-2.609	Depositor
Average map value	-0.000	Depositor
Map value standard deviation	0.323	Depositor
Recommended contour level	0.205	Depositor
Map size ($\text{\AA}$)	90.299995, 118.649994, 133.875	wwPDB
Map dimensions	255, 226, 172	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing ($\text{\AA}$)	0.525, 0.525, 0.525	Depositor

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.88	43/7442 (0.6%)	0.96	53/10084 (0.5%)
2	B	1.03	58/6036 (1.0%)	1.20	72/8217 (0.9%)
All	All	0.95	101/13478 (0.7%)	1.08	125/18301 (0.7%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	12
2	B	0	2
All	All	0	14

The worst 5 of 101 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	906	ILE	C-O	-13.58	1.07	1.23
1	A	458	ARG	C-O	-11.36	1.10	1.24
2	B	156	ASP	C-O	-11.32	1.10	1.24
2	B	152	HIS	C-O	-10.04	1.12	1.24
1	A	829	TYR	C-O	-9.72	1.14	1.24

The worst 5 of 125 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	341	PRO	CB-CA-C	9.08	119.93	111.39
2	B	576	PHE	N-CA-CB	-8.86	97.63	110.56
2	B	609	LEU	N-CA-CB	-8.83	96.62	110.22
1	A	724	PHE	CA-C-O	-8.74	110.79	121.02

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	476	PHE	CA-CB-CG	-8.38	105.42	113.80

There are no chirality outliers.

5 of 14 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	144	ARG	Sidechain
1	A	27	ARG	Sidechain
1	A	271	ARG	Sidechain
1	A	32	ARG	Sidechain
1	A	66	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7262	7164	7175	86	0
2	B	5877	6057	6078	126	0
3	A	1	0	0	0	0
4	A	421	0	0	15	0
4	B	225	0	0	27	0
All	All	13786	13221	13253	207	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 207 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:604:HIS:HB3	4:B:943:HOH:O	1.70	0.89
2:B:101:MET:HE2	2:B:127:ILE:HG22	1.66	0.77
1:A:752:ILE:HD11	1:A:771:LEU:HD13	1.69	0.74
2:B:627:SER:HB3	4:B:865:HOH:O	1.86	0.74
2:B:375:THR:HG21	2:B:387:LYS:HG3	1.71	0.72

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	901/1031 (87%)	859 (95%)	39 (4%)	3 (0%)	36	46
2	B	734/773 (95%)	695 (95%)	37 (5%)	2 (0%)	36	46
All	All	1635/1804 (91%)	1554 (95%)	76 (5%)	5 (0%)	37	46

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	711	ALA
2	B	577	TRP
1	A	480	VAL
1	A	566	ALA
2	B	341	VAL

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	746/896 (83%)	718 (96%)	28 (4%)	29	44
2	B	633/681 (93%)	609 (96%)	24 (4%)	29	44
All	All	1379/1577 (87%)	1327 (96%)	52 (4%)	30	44

5 of 52 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	10	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	239	ILE
2	B	683	MET
2	B	11	PRO
2	B	154	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	129	ASN
2	B	89	GLN
2	B	249	HIS
2	B	289	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-79079. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 86




Y Index: 113




Z Index: 127


6.2.2 Raw map



X Index: 288



Y Index: 288




Z Index: 288


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 73




Y Index: 83




Z Index: 127


6.3.2 Raw map



X Index: 291



Y Index: 265




Z Index: 283


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.205. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.

7.2 Volume estimate [i](#)



The volume at the recommended contour level is 84 nm³; this corresponds to an approximate mass of 76 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.


7.3 Rotationally averaged power spectrum [i](#)

This section was not generated. The rotationally averaged power spectrum is only generated for cubic maps.

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.437 $\text{\AA}^{-1}$

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	2.29	-	-
Author-provided FSC curve	2.29	2.61	2.32
Unmasked-calculated*	2.90	3.56	3.00

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 2.90 differs from the reported value 2.29 by more than 10 %


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-79079 and PDB model 38RH. Per-residue inclusion information can be found in section [3](#) on page [5](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.205 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.205).

9.4 Atom inclusion [i](#)



At the recommended contour level, 95% of all backbone atoms, 95% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.205) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.9520	0.7150
A	0.9510	0.7150
B	0.9570	0.7150

